

An approximate version of the Loebel-Komlós-Sós conjecture

Diana Piguet* and Maya Jakobine Stein†

Abstract

Loebel, Komlós, and Sós conjectured that if at least half of the vertices of a graph G have degree at least some $k \in \mathbb{N}$, then every tree with at most k edges is a subgraph of G . Our main result is an approximate version of this conjecture for large enough $n = |V(G)|$, assumed that $n = O(k)$.

Our result implies an asymptotic bound for the Ramsey number of trees. We prove that $r(\mathcal{T}_k, \mathcal{T}_m) \leq k + m + o(k + m)$, as $k + m \rightarrow \infty$

1 Introduction

We explore how certain global assumptions on a graph G ensure the existence of specific subgraphs. More precisely, we are interested in finding trees as (not necessarily induced) subgraphs. The main conjecture in our investigations makes, to this end, assumptions on the median degree of G .

Conjecture 1 (Loebel, Komlós, Sós [6]). *Let $k > 0$. Then every graph on $n \in \mathbb{N}$ vertices of which at least $n/2$ have degree at least k , contains as subgraphs all trees with at most k edges.*

The original version for $k = n/2$ was formulated by Loebel, the generalisation to arbitrary k is due to Komlós and Sós (see [6]).

Our main result is an approximate version of Conjecture 1 for $n = O(k)$.

Theorem 2. *For every $\eta, q > 0$ there is an $n_0 \in \mathbb{N}$ such that for every graph G on $n \geq n_0$ vertices and every $k \geq qn$ the following is true.*

If at least $n/2$ vertices of G have degree at least $(1 + \eta)k$, then G contains all trees with at most k edges.

For arbitrary k , this has been conjectured by Ajtai, Komlós and Szemerédi in [1]. They gave a proof for the special case $k = n/2$.

The exact version, Conjecture 1, is trivial for stars, and for trees that consist of two stars with adjacent centres. Bazgan, Li, and Woźniak [2] have proved the conjecture for paths. The authors of the present paper proved in [10] the Loebel–Komlós–Sós conjecture for trees of diameter at most 5.

In Loebel’s version with $k = n/2$, the conjecture has recently been proved by Zhao [14] for large enough graphs. Extending the methods of Zhao, and of

*Supported by ITI grant no. 1M00216220808.

†Supported by FAPESP grant no. 05/54051-9.

the present paper, the full Loeb–Komlós–Sós conjecture has been proved very recently for $n = O(k)$ by Hladký together with the first author [8], and independently, by Cooley [4].

A generalisation of an example due to Zhao [14] shows that the bound for the number of vertices of high degree in Conjecture 1 is asymptotically best possible. It cannot be replaced by $n/2 - n/(k+1)$, whenever $k+1$ is even and divides n (for bounds in other cases, see [10]).

To see this, construct a graph G on n vertices as follows. Divide $V(G)$ into $2n/(k+1)$ sets A_i, B_i , so that $|A_i| = (k-1)/2$, and $|B_i| = (k+3)/2$, for $i = 1, \dots, n/(k+1)$. Insert all edges inside each A_i , and insert all edges between each pair A_i, B_i . Now, consider the tree T we obtain from a star with $(k+1)/2$ edges by subdividing each edge but one. Clearly, T is not a subgraph of G .

An interesting folklore observation is the following. Assume that there is a counterexample to Conjecture 1 for the dense case, i.e., there exists a graph G of order n with half of its vertices of degree k , where $n = O(k)$, that does not contain some tree of order $k+1$. By taking many copies of G , we could then construct a counterexample to Conjecture 1 for the sparse case.

The Ramsey number $r(H, H')$ of two graphs, H and H' , is defined as the minimum integer n such for every graph G of order at least n either H is a subgraph of G , or H' is a subgraph of the complement \bar{G} of G . Extending this definition, we denote by $r(\mathcal{H}, \mathcal{H}')$ the Ramsey number of two classes of graphs, \mathcal{H} and \mathcal{H}' , that is, $r(\mathcal{H}, \mathcal{H}')$ is the minimum integer n such for every graph G of order at least n either each graph $H \in \mathcal{H}$ is a subgraph of G , or each graph $H' \in \mathcal{H}'$ is a subgraph of the complement \bar{G} of G . We write $r(\mathcal{H})$ as shorthand for $r(\mathcal{H}, \mathcal{H})$. For $i \in \mathbb{N}$, let \mathcal{T}_i denote the class of all trees of order i . Zhao’s result implies that the Ramsey number $r(\mathcal{T}_{k+1}) \leq 2k$, for large k . Bounds for Ramsey numbers of trees have been studied for instance in [7].

In the same way as the bound on $r(\mathcal{T}_{k+1})$ follows from the Loeb conjecture, one can deduce from Conjecture 1, if true, a bound on $r(\mathcal{T}_{k+1}, \mathcal{T}_{m+1})$. Namely, for any colouring of the edges of the complete graph K_{m+k} with two colours, either half of the vertices have degree k in the subgraph induced by the first colour, or half of the vertices have degree m in the subgraph induced by the second colour. So the Loeb–Komlós–Sós conjecture would then imply that $r(\mathcal{T}_{k+1}, \mathcal{T}_{m+1}) \leq k+m$. This upper bound has been conjectured in [6], and it is not difficult to see that the bound is best possible.

Using Theorem 2, we prove this to be asymptotically true.

Proposition 3. $r(\mathcal{T}_{k+1}, \mathcal{T}_{m+1}) \leq k+m+o(k+m)$, as $k+m \rightarrow \infty$.

It is not difficult to see that the exact bound of $r(\mathcal{T}_{k+1}, \mathcal{T}_{m+1}) \leq k+m$ also follows from a positive answer to the Erdős–Sós conjecture. This well-known conjecture states that each graph with average degree greater than $k-1$ contains all trees with at most k edges as subgraphs. For partial results on the Erdős–Sós conjecture, see e.g. [3, 11, 13]. Ajtai, Komlós, Simonovits and Szemerédi proved the Erdős–Sós conjecture for large n . (Unfortunately, a manuscript is not available yet.)

Our proof of Theorem 2 is inspired by the proof of the approximate version of the Loeb conjecture by Ajtai, Komlós and Szemerédi [1]. Here also, we use the regularity lemma followed by a Gallai–Edmonds decomposition of the reduced cluster graph. This enables us to find a certain substructure in the cluster graph,

which contains a large matching, and captures the degree condition on G . The tree is then embedded mainly into the matching edges.

We shall see that in the case that $k \geq n/2$, it is not difficult to obtain the same structure as in [1]. Our proof then follows [1], providing all details.

In the case that $k < n/2$, however, the situation is more complex. We will have to content ourselves with a less favourable structure in the cluster graph, which complicates the embedding of the tree. For a brief outline of the crucial ideas we then employ, see Section 3.1. The full proof is given in the remainder of Section 3.

Using similar ideas of proof, we extend Theorem 2 in a different direction. We pursue the question which other subgraphs are contained in our graph G from Theorem 2.

Our third result asserts that we can replace the trees with bipartite graphs that may have a few more edges than trees.

Theorem 4. *For every $\eta, q > 0$ and for every $c \in \mathbb{N}$ there is an $n_0 \in \mathbb{N}$ so that for each graph G on $n \geq n_0$ vertices and each $k \geq qn$ the following is true.*

If at least $n/2$ vertices of G have degree at least $(1 + \eta)k$, then each connected bipartite graph Q on $k + 1$ vertices with at most $k + c$ edges is a subgraph of G .

In particular, the condition of Theorem 2 allows for embedding even cycles in G :

Corollary 5. *For every $\eta, q > 0$ there is an $n_0 \in \mathbb{N}$ so that for all graphs G on $n \geq n_0$ vertices and each $k \geq qn$ the following is true.*

If at least $n/2$ vertices of G have degree at least $(1 + \eta)k$, then G contains all even cycles of length at most $k + 1$.

Observe that a sharp version of Theorem 4 does not hold, as is witnessed by the following example. Take the complete graph on k vertices and the empty graph on k vertices. Connect these two graphs with a matching of order k . The graph we obtain satisfies the condition of the sharp version of Theorem 4, but does not contain the cycle of length $k + 1$.

Also, the condition that Q is bipartite is necessary. This can be seen by considering the complete bipartite graph $K_{(1+\eta)k, (1+\eta)k}$. This graph satisfies the condition of Theorem 4, but all its subgraphs are bipartite.

Our paper is organised as follows. In Section 2.1, we introduce the regularity lemma and discuss some basic properties of regularity. Our tool for finding the desired structure of the cluster graph, Lemma 8, will be proved in Section 2.2. All of Section 3 is dedicated to the proof of our main result, Theorem 2.

In Section 4, we explore applications and generalisations of Theorem 2. Our asymptotic bound for Ramsey numbers of trees (Proposition 3) will be derived in Section 4.1. In Section 4.2, we prove Theorem 4.

2 Preliminaries

The purpose of this section is to introduce the two main tools used in the proofs of Theorem 2 and Theorem 4. The first of these tools is the well-known regularity lemma. The second is Lemma 8, which will give structural information on our graph G from Theorem 2 (and Theorem 4). We derive it from the Gallai-Edmonds matching theorem.

2.1 Regularity

In this subsection, we introduce the notion of regularity, state Szemerédi's regularity lemma, and review a few useful properties of regularity. All of this is well-known, so the advanced reader is invited to skip this section. For an instructive survey on the regularity lemma and its applications, consult [9].

Let us first go through some necessary notation. For a graph $G = (V, E)$, with $W \subseteq E$ and $S \subseteq V$, we will write $G - W$ for the subgraph $(V, E \setminus W)$ of G , and $G - S$ the subgraph of G which is obtained by deleting all vertices of S and all incident edges. For subsets X and Y of the vertex set $V(G)$, define $N_Y(X)$ as the set of all neighbours of X in $Y \setminus X$. If X and Y are disjoint, then let $e(X, Y)$ denote the number of edges between X and Y . The *density* of the pair (X, Y) is $d(X, Y) := \frac{e(X, Y)}{|X||Y|}$.

A bipartite graph G with partition classes C_1 and C_2 is called ε -regular if for all subsets $C'_1 \subseteq C_1$, $C'_2 \subseteq C_2$ with $|C'_1| \geq \varepsilon|C_1|$ and $|C'_2| \geq \varepsilon|C_2|$, it is true that $|d(C_1, C_2) - d(C'_1, C'_2)| < \varepsilon$.

A partition $C_0 \cup C_1 \cup \dots \cup C_N$ of $V(G)$ is called (ε, N) -regular, if

- $|C_0| \leq \varepsilon n$ and $|C_i| = |C_j|$ for $i, j = 1, \dots, N$,
- all but at most εN^2 pairs (C_i, C_j) with $i \neq j$ are ε -regular.

We are now ready to state Szemerédi's regularity lemma.

Theorem 6 (Regularity lemma, Szemerédi [12]). *For every $\varepsilon > 0$ and $m_0 \in \mathbb{N}$, there exist $M_0, N_0 \in \mathbb{N}$ so that every graph G of order $n \geq N_0$ admits an (ε, N) -regular partition of its vertex set $V(G)$ with $m_0 \leq N \leq M_0$.*

Call the partition classes C_i of G *clusters*. Now, for each graph G , for each (ε, N) -regular partition of $V(G)$, and for any density p define the *cluster graph* (sometimes called *reduced graph*) in the following standard way.

First, we construct an auxiliary graph G_p obtained from G by deleting all edges inside the clusters C_i , all edges that are incident with C_0 , all edges between irregular pairs, and all edges between regular pairs (C_i, C_j) of density $d(C_i, C_j) < p$. Set $s := |C_i|$, and observe that

$$|E(G - G_p)| \leq N \frac{s^2}{2} + \varepsilon n^2 + \varepsilon N^2 s^2 + \frac{N^2}{2} p s^2 \leq \left(\frac{1}{2m} + 2\varepsilon + \frac{p}{2} \right) n^2. \quad (1)$$

Now, the *cluster graph* $H = H_p$ on the vertex set $\{C_i\}_{1 \leq i \leq N}$ has an edge $C_i C_j$ for each pair (C_i, C_j) of clusters that has positive density in G_p . We shall prefer to work with the *weighted cluster graph* $\bar{H} = \bar{H}_p$ which we obtain from H by assigning weights

$$w(C_i C_j) := d(C_i, C_j) s$$

to the edges $C_i C_j \in E(H)$.

In the setting of weighted graphs, the (weighted) *degree* of a vertex v is defined as

$$\text{d\bar{e}g}(v) := \sum_{u \in N(v)} w(vu),$$

and the degree into a subset $U \subseteq V(\bar{H})$, where we only count the weights of $v-U$ edges, is denoted by $\text{d\bar{e}g}_U(v)$. We shall adopt this notation for our weighted cluster graph \bar{H} . For a subset $X \subseteq C_j$, we write

$$\text{d\bar{e}g}_X(C_i) := \frac{e(X, C_i)}{s}.$$

For a set \mathcal{Y} of subsets of distinct clusters from $G_p - C_i$, we shall write $\text{d\bar{e}g}_{\mathcal{Y}}(C_i)$ for $\sum_{Y \in \mathcal{Y}} \text{d\bar{e}g}_Y(C_i)$.

We shall often use edges of \bar{H} to represent the respective subgraph of G_p , or its vertex set. For example, an edge $e = CD \in E(\bar{H})$, might refer to the subgraph of G_p induced by $C \cup D$, or to $C \cup D$ itself. And for a set $U \subseteq C \cup D$, we sometimes use the shorthand $e \cap U$ for $(C \cup D) \cap U$.

Let us review some basic properties of G_p and \bar{H} . Let $C, D \in V(\bar{H})$: We call a set $D' \subseteq D$ *significant*, if $|D'| \geq \varepsilon s$. A vertex $v \in C$ is called *typical* to a significant set D' if $\text{deg}_{D'}(v) \geq \text{deg}_{D'}(C) - 2\varepsilon s$. Observe that

$$\text{At most } \varepsilon s \text{ vertices of } C \text{ are not typical to a given significant set } D'. \quad (2)$$

Similarly, we have that

$$\text{all but at most } \varepsilon s \text{ vertices } v \text{ of } C \text{ have degree } \text{deg}_{G_p}(v) \leq \text{d\bar{e}g}(C) + 2\varepsilon s. \quad (3)$$

Also, almost all vertices of any cluster $C \in V(\bar{H})$ are *typical* to almost all significant sets, in the following sense.

If \mathcal{Y} is a set of significant subsets of clusters in $V(\bar{H})$, then

$$|\{Y \in \mathcal{Y} : \text{deg}_Y(v) \geq \text{d\bar{e}g}_Y(C) - 2\varepsilon s\}| \geq (1 - \sqrt{\varepsilon})|\mathcal{Y}|, \quad (4)$$

for all but at most $\sqrt{\varepsilon}s$ vertices $v \in C$.

To see this, assume that the set $C' \subseteq C$ of vertices not satisfying (4) is larger than $\sqrt{\varepsilon}s$. Then

$$\begin{aligned} \sum_{Y \in \mathcal{Y}} |\{v \in C : v \text{ is not typical to } Y\}| &\geq \sum_{v \in C'} |\{Y \in \mathcal{Y} : v \text{ is not typical to } Y\}| \\ &\geq |C'| \sqrt{\varepsilon} |\mathcal{Y}| \\ &> \varepsilon s |\mathcal{Y}|. \end{aligned}$$

Thus there is a $Y \in \mathcal{Y}$ such that more than $\varepsilon|C|$ vertices in C are not typical to Y , a contradiction to (2).

2.2 The matching

The main interest in this subsection is Lemma 8, which will give us important structural information on the cluster graph H that corresponds to the graph G from Theorem 2 (or later Theorem 4). A weaker variant of this lemma, Lemma 9 below, appeared in [1].

For the proof of Lemma 8, we need a simplified version of the Gallai-Edmonds matching theorem, a proof of which can be found for example in [5].

A 1-*factor*, or *perfect matching*, of a graph G is a 1-regular spanning subgraph of G . We call G *factor-critical*, if for each $v \in V(G)$, there exists a perfect matching of $G - v$.




Figure 1: The graph \bar{H} with the matching M , and sets L , S and Y .

Theorem 7 (Gallai, Edmonds). *Every graph G contains a set $S \subseteq V(G)$ so that each component of $G - S$ is factor-critical, and so that there is a matching in G that matches the vertices of S to vertices of different components of $G - S$.*

We are now ready for one of the key tools in the proof of Theorem 2. (Recall that we write $\text{d\ddot{e}g}_{M \cup L}(v)$ for $\text{d\ddot{e}g}_{V(M) \cup L}(v)$.)

Lemma 8. *Let \bar{H} be a weighted graph on N vertices, and let $K \in \mathbb{R}$. Let L be the set of those vertices $v \in V(\bar{H})$ with $\text{d\ddot{e}g}(v) \geq K$. If $|L| > N/2$, then there are two adjacent vertices $v_A, v_B \in L$, and a matching M in \bar{H} such that one of the following holds.*

- (a) M covers $N(\{v_A, v_B\})$,
- (b) M covers $N(v_A)$, and $\text{d\ddot{e}g}_{M \cup L}(v_B) \geq K/2$. Moreover, each edge in M has at most one endvertex in $N(v_A)$.

Proof. Observe that we may assume that $Y := V(\bar{H}) \setminus L$ is independent. (In fact, otherwise we simply delete the edges in $E(Y)$, which will not affect the degree of the vertices in L .)

Theorem 7 applied to the unweighted version of \bar{H} yields a set $S \subseteq V(\bar{H})$. Among all matchings M' satisfying the conclusion of Theorem 7, choose M' so that it contains a maximal number of vertices of Y . Extend M' to a maximal matching M of \bar{H} .

Set $L' := L \setminus S$. Clearly, if there is an edge $v_A v_B$ with endvertices $v_A, v_B \in L'$, then (a) holds. Therefore, we may assume that L' is independent.

Then, each edge of \bar{H} that is not incident with S has one endvertex in L' , and one in Y . Consider any component C of $\bar{H} - S$. Since C is factor-critical, we have that $|(C - u) \cap Y| = |(C - u) \cap L'|$, for every $u \in V(C)$. Hence, C consists of only one vertex, and so must every component of $\bar{H} - S$.

Denote by X the subset of Y that is not covered by M . Set $\tilde{L} := N(L') \cap L \subseteq S$ (see Figure 1). Now, if there is a vertex $v_B \in \tilde{L}$ whose weighted degree into $\bar{H} - X$ is at least $K/2$, then v_B , together with any of its neighbours v_A in L' , satisfies (b). So, we may assume that for each $u \in \tilde{L}$,

$$\text{d\ddot{e}g}_{\bar{H} - X}(u) < K/2, \quad (5)$$

and hence $\text{d\ddot{e}g}_X(u) \geq K/2$.

On the other hand, $\text{d\ddot{e}g}_{\tilde{L}}(w) < K$ for each $w \in X$. Thus, by double (weighted) edge-counting, it follows that

$$|X| \geq \frac{|\tilde{L}|}{2}. \quad (6)$$

Set $S' := S \cap Y$. By (5), the total weight of the edges in $E(\tilde{L} \cup S', L')$ is less than $|\tilde{L}|K/2 + |S'|K$, while each vertex of L' has weighted degree at least K into $\tilde{L} \cup S'$. Thus, again by double edge-counting, and by (6),

$$|X| + |S'| \geq \frac{|\tilde{L}|}{2} + |S'| > |L'|. \quad (7)$$

Furthermore, since Y is independent, M matches $S' \subseteq Y$ to L' . Thus $|L'| \geq |S'| + |L \setminus M|$, and so, by (7),

$$|X| > |L \setminus M|.$$

Since $|L| > \frac{N}{2}$, this implies that M contains an edge uv with both $u, v \in L$. We may assume that $v \in L'$ and $u \in \tilde{L}$. By (5), u has a neighbour w in X . Hence, the matching $M' \cup \{uw\} \setminus \{uv\}$ covers more vertices of Y than M' does, a contradiction to the choice of M' . \square

Note that in the case $K \geq N/2$ the situation in Lemma 8 is less complicated. In that case, observe that clearly $|S| \leq |V(\bar{H} - S)|$. So, either $|S| = |V(\bar{H} - S)|$ (in which case conclusion (a) of Lemma 8 holds), or there is a component C of $\bar{H} - S$ that has more than one vertex. Thus, as C is factor-critical, there exists an L' - L' edge in C , and (a) holds again.

This proves the following lemma, which appeared in [1].

Lemma 9. *If $K \geq N/2$, then Lemma 8 always yields Case (a).*

In the case $k \geq n/2$, this observation simplifies our proof of Theorem 2 considerably, as then only the simplest case needs to be treated. We shall not make use of Lemma 9 in our proof of Theorem 2.

3 Proof of Theorem 2

The organisation of this section is as follows. The first subsection is devoted to an outline of our proof, highlighting the main ideas, leaving out all details. In Subsection 3.2, assuming that we are given a host graph G and a tree T^* as in Theorem 2, we shall first apply the regularity lemma to G . We then use Lemma 8 to find a substructure of the corresponding weighted cluster graph \bar{H} , which will facilitate the embedding of T^* .

We shall prepare T^* for this by cutting it into small pieces in Subsections 3.3 and 3.4. Then, in Subsection 3.5, we partition the matching given by Lemma 8, according to the decomposition of the tree T^* . In Subsection 3.6, we expose tools that we need for our embedding. What remains is the actual embedding procedure, which we divide into the two cases given by Lemma 8, and treat separately in Subsections 3.7 and 3.8.

3.1 Overview

In this subsection, we shall give an outline of our proof of Theorem 2. So, assume that we are given $\eta > 0$ and $q > 0$. The regularity lemma applied to parameters depending on η and q yields an $n_0 \in \mathbb{N}$. Now, let $n \geq n_0$, let $k \geq qn$, let G be a graph of order n that satisfies the condition of Theorem 2, and let T^* be a tree with k edges. We wish to find a subgraph of G that is isomorphic to T^* , i.e. we would like to *embed* T^* in G .

In order to do so, consider the weighted cluster graph \bar{H} corresponding to G that is given by the regularity lemma. Denote by $L \subseteq V(\bar{H})$ the set of those clusters that have degree at least $(1 + \pi')k$ in \bar{H} , where $\pi' = \pi'(\eta, q) > 0$. Apply Lemma 8 to \bar{H} and $K := (1 + \pi')k$ which yields vertices $A, B \in V(\bar{H})$ and a matching M . The rest of our proof will be divided into two cases, corresponding to the two possible conclusions (a) and (b) of Lemma 8.

As the technical details for these two cases overlap, we will not completely separate them later on in the proof. In this outline, however, we think it is more instructive to present first the easier proof for Case (a), and then turn our attention to Case (b).

If the output of Lemma 8 is Case (a), then we shall decompose T^* into small subtrees (of order much below ηk) and a small set SD of vertices (of constant order in n), so that between any two of our subtrees lies a vertex from SD (the name SD stands for ‘seeds’). In fact, SD is the disjoint union of two sets SD^A and SD^B , and each tree of $T^* - SD$ is adjacent to only one of these two sets. Denote the set of trees adjacent to SD^A by \mathcal{T}_A , and the set of trees adjacent to SD^B by \mathcal{T}_B . The formal definition of SD , \mathcal{T}_A and \mathcal{T}_B can be found in Section 3.3.

Next, in Section 3.5, we partition the matching M from Lemma 8 into M_A and M_B . This is done in a way so that $\text{d\ddot{e}g}_{M_A}(A)$ is large enough so that $F_A := \bigcup \mathcal{T}_A$ fits into M_A , and $\text{d\ddot{e}g}_{M_B}(B)$ is large enough so that $F_B := \bigcup \mathcal{T}_B$ fits into M_B .

Finally, in Section 3.7, we embed SD^A in A and SD^B in B and use the regularity of the edges in \bar{H} to embed the small trees of $\mathcal{T}_A \cup \mathcal{T}_B$, one after the other, levelwise, into $M_A \cup M_B$. The order of this embedding procedure will be such that the already embedded part of T^* is always connected.

Moreover, the structure of our decomposition of T^* , and the fact that we embed the trees from $\mathcal{T}_A \cup \mathcal{T}_B$ in the matching edges, ensures that the predecessor of any vertex $r \in SD^A \cup SD^B$ is embedded in a cluster that is adjacent to A , respectively to B (in which we wish embed r). This enables us to embed all of SD in $A \cup B$, as planned.

An important detail of our embedding technique is that we shall always try to *balance* the embedding in the matching edges, in the sense that the used part of either side should have about the same size. We only allow for an unbalanced embedding if the degree of A resp. B into one of the endclusters of the concerned edge is already ‘exhausted’ (cf. Property (\diamond) in Section 3.6). In practice, this means that whenever we have the choice into which endcluster of an edge $e \in M$ we embed the root of some tree of $\mathcal{T}_A \cup \mathcal{T}_B$, we shall choose the side carefully. In this manner, we can ensure that all of T^* will fit into M (or more precisely into the corresponding subgraph of G). This finishes the embedding of T^* in Case (a) of Lemma 8.

In Case (b) of Lemma 8, it is not possible to partition the matching M into M_A and M_B so that F_A fits into M_A and F_B fits into M_B , as in Case (a). More precisely, for any partition of M into M_A and M_B , if $\text{d\bar{e}g}_{M_A}(A)$ allows for the embedding of a forest of order t , say, in M_A , then $\text{d\bar{e}g}_{M_B \cup L}(B)$ only guarantees for the embedding of a forest of order at most $(k-t)/2$ in the subgraph of G_p induced by M_B and the edges incident with L' , where $L' := L \setminus M$. For more details on this, see Lemma 10.

We use a combination of two strategies to overcome this problem. Firstly, we shall embed T^* in two phases, leaving for the second phase some subtrees that are (each) adjacent to only one vertex from SD . Secondly, we shall embed some of the trees from \mathcal{T}_B in part of the matching reserved for F_A . This means that we ‘switch’ some of our trees to \mathcal{T}_A .

Let us explain the two strategies in more detail. We modify our sets $\mathcal{T}_A \cup \mathcal{T}_B$, in the following way. Denote by $\bar{\mathcal{T}}_A$ the set of those trees from \mathcal{T}_A that are adjacent to only one vertex from SD^A , and similarly define $\bar{\mathcal{T}}_B$. (Observe that thus the deletion of any tree in $\bar{\mathcal{T}}_A \cup \bar{\mathcal{T}}_B$ leaves T^* connected.)

We may assume that

$$|V(\bigcup \bar{\mathcal{T}}_A)| \geq |V(\bigcup \bar{\mathcal{T}}_B)|.$$

Finally, set $\mathcal{T}' := (\mathcal{T}_A \cup \mathcal{T}_B) \setminus (\bar{\mathcal{T}}_A \cup \bar{\mathcal{T}}_B)$. Our plan now is to first embed the trees from $\mathcal{T}' \cup \bar{\mathcal{T}}_B$ together with the vertices from SD and to postpone the embedding of $\bar{F}_A := \bigcup \bar{\mathcal{T}}_A$ to a later stage. As the part of the tree embedded in the first phase is connected, we avoid the difficulty of having to connect already embedded parts of T^* in the second phase.

Now, we shall partition M into M_F and \bar{M}_B so that $\text{d\bar{e}g}_{M_F}(A)$ allows for the embedding of $\bigcup \mathcal{T}'$, and $\text{d\bar{e}g}_{\bar{M}_B \cup L}(B)$ allows for the embedding of $\bar{F}_B := \bigcup \bar{\mathcal{T}}_B$. This actually means that the place we reserved for the embedding of $F_B - V(\bar{F}_B)$ lies in M_F . Therefore, we shall ‘switch’ this forest to \mathcal{T}_A (which is the second of our strategies).

Let us explain what we mean by *switching*. For each tree $T \in \mathcal{T}_B \setminus \bar{\mathcal{T}}_B$, delete all vertices from T that are adjacent to SD^B in T^* and add them to SD^A . Put the components of what remains of T into \mathcal{T}_A . Denote the thus enlarged SD^A by $\bar{S}D^A$ and set $\bar{S}D := \bar{S}D^A \cup SD^B$.

After switching all trees $T \in \mathcal{T}_B \setminus \bar{\mathcal{T}}_B$, denote by \mathcal{T}_F the (enlarged) set $\mathcal{T}_A \setminus \bar{\mathcal{T}}_A$. That is, \mathcal{T}_F consists of all trees from the original $\mathcal{T}_A \setminus \bar{\mathcal{T}}_A$, together with all trees we generated by switching. It will be easy to verify that the switching procedure does not increase too much the number of seeds.

Also, each tree from \mathcal{T}_F and $\bar{\mathcal{T}}_A$ is adjacent only to the enlarged $\bar{S}D^A$, and each tree from $\bar{\mathcal{T}}_B$ is still adjacent only to SD^B . For details on the switching procedure, consult Section 3.4.

It remains to embed T^* in G , which is done in Section 3.8. We first embed the vertices from $\bar{S}D^A \cup SD^B$ in $A \cup B$, embed $F_F := \bigcup \mathcal{T}_F$ in M_F , and embed part of $\bar{\mathcal{T}}_B$ in \bar{M}_B , in the same way as in Case (a). In a second phase, we embed the remaining trees from $\bar{\mathcal{T}}_B$ into edges of H that are incident with L' . For each tree, we are able to find a free space in a suitable edge because of the high degree of the clusters from L' .

In the remaining third phase we wish to embed \bar{F}_A . We shall now use all of M , forgetting about the partition into M_F and \bar{M}_B . The neighbours of the trees from $\bar{\mathcal{T}}_A$ in $\bar{S}D^A$ have already been embedded in the first phase. Having chosen

their images carefully then, ensures that now they have still large enough degree into what is not yet used of M . Hence, there is enough place for \bar{F}_A in M . Also, it is essential here that each edge of M meets $N(A)$ in at most one cluster. The reason is that parts of these clusters might have been used in the first and second phases of the embedding. So, some of the edges involved might be unbalanced, in the sense above, because e.g. the degree of B was such that we were not able to choose the endcluster in which we embedded the roots of the trees from \bar{T}_B . However, as each edge of M has at most one endcluster in $N(A)$, it is irrelevant whether the embedding is balanced or not in these edges. The embedding itself of \bar{F}_A is done as before. This finishes the sketch of our proof in Case (b).

3.2 Preparations

We shall now prove Theorem 2. First of all, we fix a few constants depending on η and q . Set

$$\pi := \min\{\eta, q\}, \quad \varepsilon := \frac{\pi^7 q}{25 \cdot 10^7} \quad \text{and} \quad m_0 := \frac{500}{q\pi^3}.$$

The regularity lemma (Theorem 6) applied to ε , and m_0 yields natural numbers M_0 and N_0 .

Fix

$$\beta := \frac{\varepsilon}{M_0}, \quad p := \frac{\pi^3 q}{250} \quad \text{and} \quad n_0 := \max\left\{N_0, \frac{8M_0}{\beta} \cdot \frac{8}{p}\right\}.$$

Thus our constants satisfy the following relations

$$\frac{1}{n_0} \ll \beta \ll \varepsilon \ll \frac{1}{m_0} < p \ll \pi \leq q,$$

where $a \ll b$ stands for the fact that $a < \frac{\pi}{100}b$.

In particular, p satisfies

$$4\varepsilon + \frac{1}{m_0} < p = \frac{\pi^3 q}{250}. \tag{8}$$

Let $n \geq n_0$, let $k \geq qn$, and let G be a graph of order n which has at least $\frac{n}{2}$ vertices of degree at least $(1 + \eta)k$. Suppose T^* is a tree of order $k + 1$. Our aim is to find an embedding $\varphi : V(T^*) \rightarrow V(G)$ that preserves adjacency.

Now, by Theorem 6 there exists an (ε, N) -regular partition of $V(G)$, with $m_0 \leq N \leq M_0$. As in Section 2.1, let G_p be the subgraph of G that preserves exactly the edges between regular pairs of density at least p .

By (1) and by (8),

$$|E(G - G_p)| < pn^2 \leq \frac{\pi^3}{250}kn.$$

Thus, for all but at most $\frac{\pi^2}{50}n$ vertices v , we have that $\deg_{G_p}(v) \geq \deg_G(v) - \frac{\pi}{5}k$. Hence,

$$G_p \text{ has at least } \left(1 - \frac{\pi^2}{25}\right)\frac{n}{2} \text{ vertices of degree at least } \left(1 + \frac{4\pi}{5}\right)k.$$

Let $\bar{H} = \bar{H}_p$ be the weighted cluster graph corresponding to G_p . Denote by L the set of those clusters in $V(\bar{H})$ that contain more than εn vertices of degree at least $(1 + \frac{4\pi}{5})k$ in G_p . A simple calculation shows that $|L| > (1 - \frac{\pi^2}{5})\frac{N}{2}$.

Now, delete $\min\{\pi^2 N/5, |V(\bar{H}) \setminus L|\}$ clusters in $V(\bar{H}) \setminus L$ to obtain a subgraph of the cluster graph \bar{H} . As this subgraph is very similar (or identical) to \bar{H} , in the rest of the text we shall denote it as well by \bar{H} . So from now on, by \bar{H} , we shall always refer to this subgraph. Each cluster in L drops its degree by at most $\frac{\pi^2}{5}Ns \leq \frac{\pi k}{5}$. Thus, by (3), each cluster X in L has degree

$$\text{d\bar{e}g}_{\bar{H}}(X) > (1 + \frac{3\pi}{5})k - 2\varepsilon n > (1 + \frac{\pi}{5})k. \quad (9)$$

Then Lemma 8 applied to \bar{H} and $K := (1 + \frac{\pi}{5})k$ yields an edge $AB \in E(\bar{H})$ with $A, B \in L$, together with a matching M' of \bar{H} , which satisfy (a) or (b) of Lemma 8. Obtain M from M' by deleting all edges that are incident with A or with B . In Case (a) of Lemma 8, we calculate that

$$\begin{aligned} \min\{\text{d\bar{e}g}_M(A), \text{d\bar{e}g}_M(B)\} &\geq (1 + \frac{\pi}{5})k - \frac{3n}{N} \\ &\geq (1 + \frac{\pi}{5} - \frac{3}{qm_0})k \\ &\geq (1 + \frac{\pi}{10})k. \end{aligned} \quad (10)$$

Similarly, in Case (b) it follows that

$$\text{d\bar{e}g}_M(A) \geq (1 + \frac{\pi}{10})k \quad \text{and} \quad \text{d\bar{e}g}_{M \cup L}(B) \geq (1 + \frac{\pi}{10})\frac{k}{2}. \quad (11)$$

Thus, for the remainder of our proof of Theorem 2 we shall work with the assumption that there is a matching M of \bar{H} and vertices $A, B \notin V(M)$ so that

1. $\text{d\bar{e}g}_M(A), \text{d\bar{e}g}_M(B) \geq (1 + \frac{\pi}{10})k$, or
2. $\text{d\bar{e}g}_M(A) \geq (1 + \frac{\pi}{10})k$, $\text{d\bar{e}g}_{M \cup L}(B) \geq (1 + \frac{\pi}{10})\frac{k}{2}$, and each cluster in $N(A)$ meets a different edge of M .

We shall refer to these two cases as ‘Case 1’ and ‘Case 2’, respectively. We will embed the tree T^* in the subgraph $G'_p \subseteq G_p$ corresponding to \bar{H} , using two different strategies in Case 1 and in Case 2.

3.3 Partitioning the tree

In this section, we shall cut our tree into small pieces. More precisely, we shall define a set $SD \subseteq V(T^*)$, and sets \mathcal{T}_A and \mathcal{T}_B of disjoint small subtrees of T^* which are connected through the vertices from SD . Moreover, SD together with the union of all trees from $\mathcal{T}_A \cup \mathcal{T}_B$ will span T^* .

Fix a root R of T^* , and regard T^* as a poset having R as the minimal element. For a vertex x of a subtree $T \subseteq T^*$, denote by $T(x)$ the subtree of T induced by x and all vertices y greater than x in the tree-order of T^* . (That is, $T(x)$ contains all vertices y such that the path between the root R and y contains the vertex x .) If $R \notin V(T)$, then define the *seed* $sd(T)$ of T as the maximal vertex of T^* which is smaller than every vertex of T .

Our sets $SD = SD^A \cup SD^B$, \mathcal{T}_A and \mathcal{T}_B will satisfy:

$$(I) \quad SD^A \cap SD^B = \emptyset,$$

- (II) $r \in SD$ lies at even distance to R if and only if $r \in SD^A$, and $R \in SD^A$,
- (III) $\mathcal{T}_A \cup \mathcal{T}_B$ consists of the components of $T^* - SD$,
- (IV) $|V(T)| \leq \beta k$, and $sd(T) \in SD$, for each $T \in \mathcal{T}_A \cup \mathcal{T}_B$,
- (V) $\max\{|SD^A|, |SD^B|\} \leq \frac{2}{\beta}$, and
- (VI) $e_{T^*}(V(F_A), SD^B) = 0$, and $e_{T^*}(V(F_B), SD^A) = 0$,

where $F_A := \bigcup_{T \in \mathcal{T}_A} T$ and $F_B := \bigcup_{T \in \mathcal{T}_B} T$ are the forests spanned by \mathcal{T}_A and \mathcal{T}_B .

Let us first define SD . To this end, we shall inductively find vertices x_i , and define auxiliary trees $T^i \subseteq T^*$. Set $T^0 := T^*$.

In step $i \geq 1$, let $x_i \in V(T^*)$ be the maximal vertex in the tree-order of $V(T^{i-1})$ with

$$|V(T^{i-1}(x_i))| > \beta k, \quad (12)$$

as illustrated in Figure 2(a), and define

$$T^i := T^{i-1} - (T^{i-1}(x_i) - x_i).$$




Figure 2: Phases of the partition of T^* .

If there is no vertex satisfying (12), then set $x_i := R$, and stop the definition process.

Say our process stops in some step j . Let A' be the set of all x_i , $i \leq j$, with even distance to the root R , and let B' be the set of all other x_i .

Note that, because of (12), at each step $i \leq j - 1$,

$$|V(T_i)| \leq |V(T_{i-1})| - (\beta k - 1),$$

and thus, by the definition of n_0 ,

$$j - 1 \leq \frac{|V(T^*)|}{\beta k - 1} \leq \frac{k + 1}{\beta k - 1} \leq \frac{3}{2\beta}.$$

Hence,

$$|A' \cup B'| \leq \frac{2}{\beta}. \quad (13)$$

For the sake of condition (VI), we shall now add a few more vertices to our sets A' and B' , which will result in the desired SD .

Let \mathcal{C} be the set of the components of $T^* - (A' \cup B')$. For each $T \in \mathcal{C}$ with $sd(T) \in A'$, denote by $A(T)$ the set of vertices adjacent to B' . Similarly, if $sd(T) \in B'$, then denote by $B(T)$ the set of vertices adjacent to A' (cf. Figure 2(b)). Set

$$SD^A := A' \cup \bigcup_{T \in \mathcal{C}} A(T), \quad \text{and} \quad SD^B := B' \cup \bigcup_{T \in \mathcal{C}} B(T)$$

and set $SD := SD^A \cup SD^B$.

Since each vertex in B' has at most one neighbour in the union of the $A(T)$, it follows that

$$|SD^A \setminus A'| \leq |B'|,$$

and analogously,

$$|SD^B \setminus B'| \leq |A'|.$$

Thus,

$$\max\{|SD^A|, |SD^B|\} \leq |A' \cup B'|. \quad (14)$$

Finally, we shall define \mathcal{T}_A and \mathcal{T}_B . Let \mathcal{C}' be the set of the components of $T^* - SD$. Set

$$\mathcal{T}_A := \{T \in \mathcal{C}' : sd(T) \in SD^A\} \quad \text{and} \quad \mathcal{T}_B := \{T \in \mathcal{C}' : sd(T) \in SD^B\},$$

as shown in Figure 2(c), and define the forests

$$F_A := \bigcup_{T \in \mathcal{T}_A} T \quad \text{and} \quad F_B := \bigcup_{T \in \mathcal{T}_B} T.$$

Observe that Conditions (I)–(IV) and (VI) are clearly met and that (V) holds because of (13) and (14).

This finishes our manipulation of the tree T^* in Case 1.

3.4 The switching

In Case 2 from Section 3.2, we shall not only cut our tree to small pieces (cf. Section 3.3), but also switch some of our small subtrees from one of the two sets $\mathcal{T}_A, \mathcal{T}_B$ to the other. We achieve this by adding some more vertices to SD , thus naturally refining our partition of T^* .

Set

$$\begin{aligned}\bar{\mathcal{T}}_A &:= \{T \in \mathcal{T}_A : e(V(T), SD - sd(T)) = \emptyset\}, \text{ and} \\ \bar{\mathcal{T}}_B &:= \{T \in \mathcal{T}_B : e(V(T), SD - sd(T)) = \emptyset\}.\end{aligned}$$

We may assume that

$$\left| \bigcup_{T \in \bar{\mathcal{T}}_A} V(T) \right| \geq \left| \bigcup_{T \in \bar{\mathcal{T}}_B} V(T) \right|. \quad (15)$$

Now, consider a tree $T \in \mathcal{T}_B \setminus \bar{\mathcal{T}}_B$ as in Figure 3(a). Denote by $S(T)$ the set of all vertices in $V(T)$ that in T^* are adjacent to some vertex of SD^B . For illustration see Figure 3(b).




Figure 3: The switching procedure.

Set

$$\bar{SD}^A := SD^A \cup \bigcup_{T \in \mathcal{T}_B \setminus \bar{\mathcal{T}}_B} S(T) \quad \text{and} \quad \bar{SD} := \bar{SD}^A \cup SD^B.$$

Finally, define

$$\mathcal{T}'_A := \bigcup_{T \in \mathcal{T}_B \setminus \bar{\mathcal{T}}_B} \{C : C \text{ is a component of } T - S(T)\}$$

and

$$\mathcal{T}_F := (\mathcal{T}_A \setminus \bar{\mathcal{T}}_A) \cup \mathcal{T}'_A.$$

(The F in \mathcal{T}_F stands for ‘first’, as this part of the tree is to be embedded first.)
 Finally, set

$$F_F := \bigcup_{T \in \mathcal{T}_F} T,$$

$$\bar{F}_A := \bigcup_{T \in \bar{\mathcal{T}}_A} T \quad \text{and} \quad \bar{F}_B := \bigcup_{T \in \bar{\mathcal{T}}_B} T.$$

Observe that our sets $\bar{S}D = \bar{S}D^A \cup SD^B$, $\mathcal{T}_F \cup \bar{\mathcal{T}}_A$, and $\bar{\mathcal{T}}_B$ still satisfy conditions (I)-(IV) and (VI) from Section 3.3 (with SD , SD^A , \mathcal{T}_A , \mathcal{T}_B , F_A , and F_B replaced by $\bar{S}D$, $\bar{S}D^A$, $\mathcal{T}_F \cup \bar{\mathcal{T}}_A$, $\bar{\mathcal{T}}_B$, \bar{F}_A , and \bar{F}_B , respectively). Instead of (V), we now have the similar

$$(V)' \quad |\bar{S}D| \leq \frac{8}{\beta},$$

since by the definition of $\bar{S}D^A$ we know that for each vertex x of SD^B , we have created at most 2 vertices of $\bar{S}D^A \setminus SD^A$ (between x and the next vertex of SD^B in direction of the root R). Thus,

$$|\bar{S}D^A| \leq |SD^A| + 2|SD^B| \leq \frac{6}{\beta},$$

as needed for (V)'.

3.5 Partitioning the matching

In this subsection, we shall divide the matching M into two parts, into which we will later embed the two forests F_A , F_B , respectively F_F and \bar{F}_B , of T^* that we defined in Subsection 3.3, resp. in Subsection 3.4. (The forest \bar{F}_A will be embedded later).

For this, we will need the following number-theoretic lemma, which appeared also in [1]. We give a short proof.

Lemma 10. *Let I be a finite set, and let $a, b, \Delta > 0$. For $i \in I$, let $a_i, b_i \in (0, \Delta]$. Suppose that*

$$\frac{a}{\sum_{i \in I} a_i} + \frac{b}{\sum_{i \in I} b_i} \leq 1. \quad (16)$$

Then there is a partition of I into I_a and I_b such that $\sum_{i \in I_a} a_i > a - \Delta$ and $\sum_{i \in I_b} b_i \geq b$.

Proof. Define a total order \preceq on I in a way that $i \preceq j$ implies $\frac{a_i}{b_i} \leq \frac{a_j}{b_j}$ for all $i, j \in I$. Let $\ell \in I$ be minimal in this order with $a \geq \sum_{i \succ \ell} a_i$.

Set $I_a := \{i \in I : i \succ \ell\}$ and set $I_b := I \setminus I_a$. It is clear that $\sum_{i \in I_a} a_i > a - \Delta$, by the definition of ℓ and as $a_\ell \leq \Delta$. So, all we have to show is that $\sum_{i \in I_b} b_i \geq b$. Indeed, suppose otherwise. Then by (16), and by the definition of ℓ , we have

that

$$\begin{aligned}
\frac{\sum_{i \in I_b} b_i}{\sum_{i \in I} b_i} &< \frac{b}{\sum_{i \in I} b_i} \\
&\leq \frac{a - \sum_{i \in I_a} a_i}{\sum_{i \in I} a_i} + \frac{b}{\sum_{i \in I} b_i} \\
&\leq 1 - \frac{\sum_{i \in I_a} a_i}{\sum_{i \in I} a_i} \\
&= \frac{\sum_{i \in I_b} a_i}{\sum_{i \in I} a_i}.
\end{aligned}$$

Multiply the two sides of this inequality with $\sum_{i \in I} a_i \cdot \sum_{i \in I} b_i$, subtract the term $\sum_{i \in I_b} a_i \cdot \sum_{i \in I_b} b_i$, and divide by $\sum_{i \in I_a} b_i \sum_{i \in I_b} b_i$ to obtain

$$\frac{a_\ell}{b_\ell} \leq \frac{\sum_{i \in I_a} a_i}{\sum_{i \in I_a} b_i} < \frac{\sum_{i \in I_b} a_i}{\sum_{i \in I_b} b_i} \leq \frac{a_\ell}{b_\ell},$$

(where the first and last inequality follow from the definition of \preceq). This yields the desired contradiction. \square

We shall now apply Lemma 10 to partition our matching $M = \{e_i\}_{i \leq |M|}$. We do this separately for the two cases from Section 3.2.

In Case 1, we set

$$a := |V(F_A)| + \frac{\pi k}{20}, \quad b := |V(F_B)| + \frac{\pi k}{20}, \quad \text{and} \quad \Delta := 2s.$$

For $i \leq |M|$, set $a_i := \text{d}\bar{\text{e}}\text{g}_{e_i}(A) \leq \Delta$, and $b_i := \text{d}\bar{\text{e}}\text{g}_{e_i}(B) \leq \Delta$. Now, (10) implies that

$$\frac{a}{\sum_{i=1}^{|M|} a_i} + \frac{b}{\sum_{i=1}^{|M|} b_i} \leq \frac{|V(F_A)| + |V(F_B)| + \frac{\pi k}{10}}{(1 + \frac{\pi}{10})k} \leq 1.$$

Hence, Lemma 10 yields a partition of M into M_A and M_B such that

$$\text{d}\bar{\text{e}}\text{g}_{M_A}(A) > |V(F_A)| + \frac{\pi k}{40} \quad \text{and} \quad \text{d}\bar{\text{e}}\text{g}_{M_B}(B) > |V(F_B)| + \frac{\pi k}{40}. \quad (17)$$

In Case 2, set

$$a := |V(F_F)| + \frac{\pi k}{20}, \quad b := |V(\bar{F}_B)| + \frac{\pi k}{40}, \quad \text{and} \quad \Delta := 2s.$$

For $i = 1, \dots, |M|$, again set $a_i := \text{d}\bar{\text{e}}\text{g}_{e_i}(A)$, and $b_i := \text{d}\bar{\text{e}}\text{g}_{e_i}(B)$. Set $L' := L \setminus M$. For $i = |M| + 1, \dots, |M| + |L'|$, set $a_i := 0$, and set $b_i := \text{d}\bar{\text{e}}\text{g}_{C_i}(B)$, where C_i is the i th cluster in L' .

Observe that by (15),

$$|V(\bar{F}_B)| \leq \frac{k - |V(F_F)|}{2}. \quad (18)$$

Now, let us check that the conditions of Lemma 10 hold. Clearly, $a_i, b_i \leq \Delta$ for all $i \leq |M| + |L'|$.

Moreover, Condition (16) holds since (11) and (18) imply that

$$\begin{aligned} \frac{a}{\sum_{i=1}^{|M|+|L'|} a_i} + \frac{b}{\sum_{i=1}^{|M|+|L'|} b_i} &\leq \frac{|V(F_F)| + \frac{\pi k}{20}}{(1 + \frac{\pi}{10})k} + \frac{|V(\bar{F}_B)| + \frac{\pi}{40}}{(1 + \frac{\pi}{10})\frac{k}{2}} \\ &\leq \frac{|V(F_F)| + 2|V(\bar{F}_B)| + \frac{\pi k}{10}}{(1 + \frac{\pi}{10})k} \\ &\leq 1. \end{aligned}$$

We thus obtain a partition of M into M_F and \bar{M}_B such that

$$\text{d\bar{e}g}_{M_F}(A) > |V(F_F)| + \frac{\pi k}{40} \quad \text{and} \quad \text{d\bar{e}g}_{\bar{M}_B \cup L'}(B) \geq |V(\bar{F}_B)| + \frac{\pi k}{40}. \quad (19)$$

Let $\mathcal{T}_B^M \subseteq \bar{\mathcal{T}}_B$ be maximal with

$$\text{d\bar{e}g}_{\bar{M}_B}(B) \geq \left| \bigcup_{T \in \mathcal{T}_B^M} V(T) \right| + \frac{\pi k}{40N} |\bar{M}_B|. \quad (20)$$

Set $\mathcal{T}_B^L := \bar{\mathcal{T}}_B \setminus \mathcal{T}_B^M$. Let $F_B^M := \bigcup_{T \in \mathcal{T}_B^M} T$ and let $F_B^L := \bar{F}_B - V(F_B^M)$. Observe that if $\mathcal{T}_B^M \neq \bar{\mathcal{T}}_B$, then the maximality of \mathcal{T}_B^M ensures that

$$\text{d\bar{e}g}_{\bar{M}_B}(B) < |V(F_B^M)| + \frac{\pi k}{40N} |\bar{M}_B| + \beta k.$$

Hence, by (19), either $\mathcal{T}_B^L = \emptyset$, or

$$\text{d\bar{e}g}_{L'}(B) \geq |V(F_B^L)| + \frac{\pi k}{80N} |L'|. \quad (21)$$

3.6 Embedding lemmas for trees

In this section, we shall prove some preparatory lemmas on embedding trees in regular pairs of \bar{H} .

Let $C, D \in V(\bar{H})$, and let $U, N \subseteq C \cup D$. We say that U has Property (\star) in CD for N if it satisfies the following.

- (\star) If $||C \cap U| - |D \cap U|| > \beta k + \varepsilon s$, then $\min\{|N \cap C|, |N \cap D|\} \leq \min\{|C \cap U|, |D \cap U|\} + 2\varepsilon s + \beta k$.

Lemma 11. *Let (T, r) be a rooted tree of order at most βk . Let $CD \in E(\bar{H})$. Suppose that $U, N \subseteq C \cup D$ are such that*

$$\min\{|N \cap C \setminus U|, |D \setminus U|\} > \frac{2}{p}(\varepsilon s + \beta k).$$

Then there is an embedding φ of T in $(C \cup D) \setminus U$ such that $\varphi(r) \in N$ and such that the following holds.

- ($\star\star$) *If U has Property (\star) in CD for N , then also $U_\varphi := U \cup \varphi(V(T))$ has Property (\star) in CD for N .*

Proof. Write $V(T) = r \cup L_1 \cup L_2 \cup \dots$, where L_ℓ is the ℓ th level of T (i. e. the set of vertices at distance ℓ to r).

First, suppose that $|N \cap D \setminus U| \leq \varepsilon s$. In this case, choose $\varphi(r) \in N \cap C \setminus U$ typical w. r. t. $D \setminus U$. (This is possible, as by (2), there are at most εs vertices that are not typical.)

Embed the rest of $V(T)$ levelwise, choosing for $\varphi(L_\ell)$ unused vertices of $D \setminus U$ that are typical with respect to $C \setminus U$, if ℓ is odd; and choosing vertices of $C \setminus U$ that are typical with respect to $D \setminus U$, if ℓ is even.

Now, suppose that $|N \cap D \setminus U| > \varepsilon s$. In this case, we may alternatively wish to embed r in $N \cap D$. We do so in either of the following cases

1. $|\bigcup_{\ell \in \mathbb{N}} L_{2\ell-1}| > |\bigcup_{\ell \in \mathbb{N}} L_{2\ell}|$ and $|C \setminus U| \geq |D \setminus U|$, or
2. $|\bigcup_{\ell \in \mathbb{N}} L_{2\ell-1}| < |\bigcup_{\ell \in \mathbb{N}} L_{2\ell}|$ and $|C \setminus U| \leq |D \setminus U|$,

and otherwise embed r in $N \cap C$, as before. After having thus chosen a place for the root r , the rest of T is embedded analogously as above (possibly swapping the roles of C and D). We have thus completed the embedding of T .

It remains to prove $(\star\star)$. So assume that U_φ has Property (\star) for N in some edge CD . Furthermore, assume that

$$\left| |C \cap U_\varphi| - |D \cap U_\varphi| \right| > \beta k + \varepsilon s. \quad (22)$$

Now, if $\left| |C \cap U| - |D \cap U| \right| > \beta k + \varepsilon s$, then Property (\star) for U_φ follows from Property (\star) for U . So suppose otherwise, that is

$$\left| |C \cap U| - |D \cap U| \right| \leq \beta k + \varepsilon s. \quad (23)$$

By (22), this means that we could not choose into which of $N \cap C$ and $N \cap D$ we would embed the root of T . Hence,

$$\min_{Y=C,D} \{|N \cap Y \setminus U|\} \leq |N \cap D \setminus U| \leq \varepsilon s.$$

Using (23), this gives

$$\begin{aligned} \min\{|N \cap C|, |N \cap D|\} &\leq \max\{|C \cap U|, |D \cap U|\} + \min_{Y=C,D} \{|N \cap Y \setminus U|\} \\ &\leq \max\{|C \cap U|, |D \cap U|\} + \varepsilon s \\ &\leq \min\{|C \cap U|, |D \cap U|\} + 2\varepsilon s + \beta k \\ &\leq \min\{|C \cap U_\varphi|, |D \cap U_\varphi|\} + 2\varepsilon s + \beta k, \end{aligned}$$

as desired. \square

Let $C, D, X \in V(\bar{H})$, let $X' \subseteq X$, let $\mathcal{Z} \subseteq V(\bar{H})$, let $U \subseteq \bigcup V(\bar{H})$, let $m \in \mathbb{N}$, and let (T, r) be a rooted tree.

We say that U has Property (\diamond) in (C, D) with respect to X if it satisfies the following.

- (\diamond) If $\left| |C \cap U| - |D \cap U| \right| > \beta k$, then $\min\{\text{d}\bar{\text{e}}\text{g}_C(X), \text{d}\bar{\text{e}}\text{g}_D(X)\} \leq \min\{|C \cap U|, |D \cap U|\} + 5\varepsilon s + \beta k$.

An embedding φ of T is a (v, X', U) -embedding in \mathcal{Z} , if $\varphi(V(T) \setminus \{r\}) \subseteq \bigcup \mathcal{Z} \setminus U$, if $\varphi(r) = v$, and if each vertex at odd distance to the root r is mapped to a vertex that is typical to X' .

A vertex is \mathcal{Z} -typical, if it is typical to each cluster from \mathcal{Z} .

The set \mathcal{Z} is said to be (m, U) -large for X , if

$$\text{d\ddot{e}g}_{\mathcal{Z}}(X) > m + |U \cap \bigcup \mathcal{Z}| + \frac{\pi k}{100N} |\mathcal{Z}|.$$

Lemma 12. *Let $(T, r), X', X$ and U be as above with $|X'| \geq |X|/2$.*

A) *Suppose M_X is a matching in $\bar{H} - X$ so that $V(M_X)$ is $(|V(T)|, U)$ -large for X , so that $v \in X$ is $V(M_X)$ -typical, and so that U has Property (\diamond) in (C, D) with respect to X , for each $CD \in M_X$.*

Then, there is a (v, X', U) -embedding φ of T in $V(M_X)$ such that $U \cup \varphi(V(T))$ has Property (\diamond) with respect to X for every $CD \in M_X$.

B) *Let $L_X, N_{L_X} \subseteq V(\bar{H})$ be such that L_X is $(|V(T)|, U)$ -large for X , and N_{L_X} is $(|V(T)|, U)$ -large for each $Y \in L_X$. If $v \in X$ is L_X -typical, then there is a (v, X', U) -embedding φ of T in $L_X \cup N_{L_X}$.*

Proof. We map r to v and embed the trees from the forest $F := T - \{r\}$ inductively. In each step $j \geq 1$, we embed a tree T^j of the forest F . Denote by $V^{<j} \subseteq V(F)$ the set $\bigcup_{i < j} V(T^i)$ of vertices we have already embedded before step j . Let S be the set of vertices in $\bigcup V(\bar{H})$ that are not typical to X' . Set $U^{<j} := U \cup S \cup \varphi(V^{<j})$. In particular, $U^{<1} = U \cup S$.

For Part **A)**, we shall moreover use two properties of U during our embedding. Firstly, if $CD \in M_X$ satisfies $||C \cap U| - |D \cap U|| \leq \beta k$, then we require that in each step $j \geq 1$

(I) $U^{<j}$ has Property (\star) for $N(v) \cap (C \cup D)$.

This property holds for $j = 1$, as the condition of Property (\star) is void, and we shall check it for each later step.

Secondly, for the edges with $||C \cap U| - |D \cap U|| > \beta k$, observe that, as the sets $U^{<j}$ are growing, Property (\diamond) ensures that for all $j \geq 1$

(II) $\min_{Y \in \{C, D\}} \{\text{d\ddot{e}g}_Y(X)\} \leq \min_{Y \in \{C, D\}} \{|Y \cap U^{<j}|\} + 5\epsilon s + \beta k$.

So, assume now that we are in step $j \geq 1$, that is, $\varphi(x)$ has been defined for all $x \in V^{<j}$, and we are about to embed T^j .

Claim 13. *There is an edge CD , with $CD \in M_X$ for Part **A)** and with $C \in L$, for Part **B)**, such that*

$$\min\{|(N(v) \cap C) \setminus U^{<j}|, |D \setminus U^{<j}|\} \geq \frac{2}{p}(\epsilon s + \beta k).$$

Before proving Claim 13, we shall show how we complete our embedding of T^j under the assumption that the claim holds for the edge $e = CD$.

Set $N := N(v) \cap e$ and let $r^j := N(r) \cap V(T^j)$ be the root of T^j . Apply Lemma 11 to embed (T^j, r^j) in $e \setminus U^{<j}$, mapping r^j to $N(v)$. Lemma 11 together with (I) for j ensures (I) for $j + 1$. As our embedding avoids S , all vertices in $\varphi(T)$ are typical to X' . This terminates step j .

Say we terminate the embedding procedure in step ℓ . Then φ is a (v, X', U) -embedding. So, for Part B), we are done. For Part A), however, we still have to prove that $U^{<\ell} \setminus S$ has Property (\diamond) in each $CD \in M_X$. To this end, assume that $\|C \cap (U^{<\ell} \setminus S)\| - \|D \cap (U^{<\ell} \setminus S)\| > \beta k$. If $\|C \cap U\| - \|D \cap U\| \leq \beta k$, then by (I), $U^{<\ell}$ has Property (\star) in CD for $N(v) \cap (C \cup D)$. Hence,

$$\begin{aligned} \min_{Y=C,D} \{\text{d}\deg_Y(X)\} &\leq \min_{Y=C,D} \{\text{deg}_Y(v)\} + 2\varepsilon s \\ &\leq \min_{Y=C,D} \{|Y \cap (U^{<\ell})|\} + 4\varepsilon s + \beta k \\ &\leq \min_{Y=C,D} \{|Y \cap (U^{<\ell} \setminus S)|\} + 5\varepsilon s + \beta k. \end{aligned}$$

On the other hand, if $\|C \cap U\| - \|D \cap U\| > \beta k$, then (II) ensures that

$$\min_{Y \in \{C,D\}} \{\text{d}\deg_Y(X)\} \leq \min_{Y \in \{C,D\}} \{|Y \cap (U^{<\ell} \setminus S)|\} + 5\varepsilon s + \beta k.$$

This shows that $U^{<\ell} \setminus S$ has Property (\diamond) in each $CD \in M_X$ for Part A). It only remains to prove Claim 13.

Proof of Claim 13: First, suppose we are in Case A). Let us start by showing that there is an edge $e = CD \in M_X$ which satisfies

$$\text{d}\deg_e(X) - |e \cap U^{<j}| \geq \frac{8}{p}(\varepsilon s + \beta k) + 2\varepsilon s. \quad (24)$$

Indeed, suppose there is no such edge. Then, as $V(M_X)$ is $(V(T), U)$ -large, we have that

$$\begin{aligned} \frac{8}{p}(\varepsilon s + \beta k)|M_X| &> \sum_{e \in M_X} (\text{d}\deg_e(X) - |e \cap U^{<j}| - 2\varepsilon s) \\ &= \text{d}\deg_{M_X}(X) - |U \cap \bigcup M_X| - |U^{<j} \setminus U| - 2\varepsilon s|M_X| \\ &\geq \text{d}\deg_{M_X}(X) - |U \cap \bigcup M_X| - |V(T)| - |S \cap M_X| - 2\varepsilon s|M_X| \\ &\geq \frac{\pi k}{100N}|V(M_X)| - 4\varepsilon s|M_X| \\ &> \frac{\pi k}{100N}|M_X|, \end{aligned}$$

which, as $\beta k \leq \frac{\varepsilon}{M_0}n \leq \varepsilon s$, implies that $16\varepsilon/p > \pi q/100$, a contradiction. So, assume now that we have chosen an edge e for which (24) holds. Clearly, we can write $e = CD$ such that

$$\frac{4}{p}(\varepsilon s + \beta k) \leq \text{d}\deg_C(X) - 2\varepsilon s - |C \cap U^{<j}| \leq |N(v) \cap C \setminus U^{<j}|. \quad (25)$$

We claim that

$$|D \setminus U^{<j}| \geq \frac{2}{p}(2\varepsilon s + \beta k). \quad (26)$$

Indeed, suppose for contradiction (26) does not hold. Then (25) implies that

$$\begin{aligned}
|C \cap U^{<j}| &\leq s - \frac{4}{p}(\varepsilon s + \beta k) \\
&= |D \cap U^{<j}| + |D \setminus U^{<j}| - \frac{2}{p}(2\varepsilon s + \beta k) - \frac{2}{p}\beta k \\
&\leq |D \cap U^{<j}| - \frac{2}{p}\beta k.
\end{aligned}$$

Hence, either by (I), or by (II), it holds that

$$\min\{\text{d}\bar{\text{e}}\text{g}_C(X), \text{d}\bar{\text{e}}\text{g}_D(X)\} \leq |C \cap U^{<j}| + 5\varepsilon s + \beta k.$$

Thus, by (24),

$$\begin{aligned}
\frac{8}{p}(\varepsilon s + \beta k) + 2\varepsilon s &\leq \text{d}\bar{\text{e}}\text{g}_e(X) - |C \cap U^{<j}| - |D \cap U^{<j}| \\
&\leq \text{d}\bar{\text{e}}\text{g}_e(X) - \min\{\text{d}\bar{\text{e}}\text{g}_C(X), \text{d}\bar{\text{e}}\text{g}_D(X)\} + 5\varepsilon s + \beta k - |D \cap U^{<j}| \\
&\leq s + 5\varepsilon s + \beta k - |D \cap U^{<j}| \\
&< |D \setminus U^{<j}| + 5\varepsilon s + 2\beta k.
\end{aligned}$$

So, $|D \setminus U^{<j}| > (\frac{8}{p} - 3)(\varepsilon s + \beta k)$, a contradiction to our assumption that (26) does not hold. This proves (26), which together with (25) implies Claim 13 for Case **A**).

Now, assume that we are in Case **B**). First we show that if some $\mathcal{Z} \subseteq V(\bar{H})$ is $(|V(T)|, U)$ -large for some $Y \in V(\bar{H})$, then there is a $Z \in \mathcal{Z} \cap N(Y)$ such that

$$\text{d}\bar{\text{e}}\text{g}_Z(Y) - |Z \cap U^{<j}| \geq \frac{2}{p}(\varepsilon s + \beta k) + 2\varepsilon s.$$

Indeed, otherwise

$$\begin{aligned}
\frac{2}{p}(\varepsilon s + \beta k)|\mathcal{Z}| &> \sum_{Z \in \mathcal{Z}} (\text{d}\bar{\text{e}}\text{g}_Z(Y) - |Z \cap U^{<j}| - 2\varepsilon s) \\
&\geq \text{d}\bar{\text{e}}\text{g}_Z(Y) - |U^{<j} \cap \bigcup \mathcal{Z}| - 2\varepsilon s|\mathcal{Z}| \\
&> \left(\frac{\pi k}{100N} - 4\varepsilon s\right)|\mathcal{Z}| \\
&\geq \frac{\pi k}{200N}|\mathcal{Z}|,
\end{aligned}$$

a contradiction, as above.

So there is a $C \in L_X$ and a $D \in N_{L_X} \cap N(C)$ such that

$$|N(v) \cap C \setminus U^{<j}| \geq \text{d}\bar{\text{e}}\text{g}_C(X) - |C \cap U^{<j}| - 2\varepsilon s \geq \frac{2}{p}(\varepsilon s + \beta k)$$

and

$$|D \setminus U^{<j}| \geq \text{d}\bar{\text{e}}\text{g}_D(C) - |D \cap U^{<j}| \geq \frac{2}{p}(\varepsilon s + \beta k),$$

as desired for Claim 13. □

3.7 The embedding in Case 1

In this subsection, we shall complete the proof of Theorem 2 under the assumption that Case 1 of Section 3.2 holds. So, we assume that there are an edge $AB \in E(\bar{H})$ and a matching $M = M_A \cup M_B$ in $\bar{H} - \{A, B\}$ as in Section 3.5. These, together with the sets $SD = SD^A \cup SD^B$, \mathcal{T}_A and \mathcal{T}_B from Section 3.3, satisfy (17).

Our embedding φ will be defined in $|SD|$ steps. In each step $i \geq 1$, we choose a suitable vertex $r_i \in SD$ and embed it together with all trees from

$$\mathcal{T}_i := \{T \in \mathcal{T}_A \cup \mathcal{T}_B : sd(T) = r_i\}.$$

Set $V_0 := \emptyset$ and for $i \geq 1$, let

$$V_i := V_{i-1} \cup \{r_i\} \cup \bigcup_{T \in \mathcal{T}_i} V(T).$$

We start with $r_1 := R$, and in each step $i > 1$, we shall choose a vertex $r_i \in SD \setminus V_{i-1}$ that is adjacent to V_{i-1} . The seed r_i will be embedded in a vertex $v_i \in A \cup B$, while \mathcal{T}_i will be mapped to edges from M (or more precisely, to the corresponding subgraph of G'_p). Set $U_0 := \emptyset$, and once φ is defined on V_i , set $U_i := \varphi(V_i)$.

For each $i \geq 0$, the following conditions will hold.

- (i) $|(A \cup B) \cap U_i| \leq i$,
- (ii) if $x \in V_i \cap N(SD^A)$, resp. $x \in V_i \cap N(SD^B)$, then $\varphi(x)$ has at least $\frac{p}{4}s$ neighbours in A , resp. in B ,
- (iii) for $CD \in M_A$, the set U_i has Property (\diamond) in CD with respect to A .
- (iv) for $CD \in M_B$, the set U_i has Property (\diamond) in CD with respect to B .

Observe that properties (i)–(iv) trivially hold for $i = 0$.

So, suppose now that we are in some step $i \geq 1$ of our embedding process. Choose $r_i \in SD$ as detailed above. Let us assume that $r_i \in SD^A$, the case when $r_i \in SD^B$ is analogous.

We embed r_i in a vertex $v_i = \varphi(r_i) \in A$ that is typical with respect to B and typical w. r. t. all but at most $2\sqrt{\varepsilon}|M_A|$ clusters of M_A . Properties (i) and (ii) for $i - 1$ ensure that if x is the predecessor of r_i in T^* , then $\varphi(x)$ has at least $\frac{ps}{4} - i$ neighbours in $A \setminus U_{i-1}$. By (2) and (4), at most $2\sqrt{\varepsilon}s$ of these vertices do not have the required properties. Hence, there are at least $(\frac{p}{4} - 2\sqrt{\varepsilon})s - i \geq 1$ suitable vertices we may choose $v_i := \varphi(r_i)$ from.

Let $M_A^i \subseteq M_A$ be a maximal submatching such that v_i is typical w. r. t. each of the end-clusters of each edge of M_A^i . Then by (4) and (17) we obtain

$$\begin{aligned} \text{deg}_{M_A^i}(A) &\geq \text{deg}_{M_A}(A) - 4\sqrt{\varepsilon}|M_A|s \\ &> |V(F_A)| + \frac{\pi k}{40} - 4\sqrt{\varepsilon}|M_A|s \\ &> |V(F_A)| + \frac{\pi k}{80} \\ &> |V(\bigcup \mathcal{T}_i)| + |U_{i-1} \cap \bigcup V(M_A)| + \frac{\pi k}{80N}|V(M_A^i)|. \end{aligned}$$

Now we use Lemma 12 Part **A**) letting (T, r) be the tree induced by r_i and the trees from \mathcal{T}_i , and setting $M_X := M_A^i$, $U := U_{i-1}$, $v := v_i$, and $X = X' = A$. It is easy to see that (i), (ii), and (iv) hold for i , as they hold for $i-1$, and by our choice of $\varphi(V_i \setminus V_{i-1})$. Lemma 12 Part **A**) ensures (\diamond) for all edges $CD \in M_A^i$. As we did not embed anything in the edges outside M_A^i , (iii) for $i-1$ implies (iii) for i , for all $CD \in M_A$.

This completes the embedding of the tree T^* in $G'_p \subseteq G$ in Case 1.

3.8 The embedding in Case 2

We shall now complete the proof of Theorem 2 under the assumption that Case 2 of Section 3.2 holds. That is, there are an edge $AB \in E(\bar{H})$ and a matching $M = M_F \cup \bar{M}_B$ in $\bar{H} - \{A, B\}$ together with sets $\bar{S}D = \bar{S}D^A \cup \bar{S}D^B$, \mathcal{T}_F , $\bar{\mathcal{T}}_A$, \mathcal{T}_B^M and \mathcal{T}_B^L from Sections 3.3 and 3.4 satisfying (20) and (21) from Section 3.5. Our embedding will be defined in three phases. In the first phase, we shall embed all vertices from $\bar{S}D$ in $A \cup B$, embed F_F in edges of M_F , and embed F_B^M in edges of \bar{M}_B . In the second phase, we shall embed F_B^L in edges incident with $L' \cap N(B)$, and in the third phase, we shall embed \bar{F}_A in the remaining space inside edges from M .

Denote by A' the set of vertices in A that are typical to all but at most $2\sqrt{\varepsilon}|M|$ clusters of $V(M)$, and denote by B' the set of vertices in B that are typical to all but at most $\sqrt{\varepsilon}|L'|$ clusters of L' .

The first phase is done analogously as in Case 1, always considering A' and B' instead of A and B . In each step, Lemma 12 Part **A**) is used in the following setting.

The rooted tree (T, r) is the tree induced by r_i and the trees from

$$\mathcal{T}_i := \{T \in \mathcal{T}_F \cup \mathcal{T}_B^M : sd(T) = r_i\}.$$

We set either $(X', X) = (A', A)$ or $(X', X) = (B', B)$, and let $v = \varphi(r_i)$. The matching M_X is a maximal submatching either of M_F or of \bar{M}_B , so that $\varphi(r_i)$ is $V(M_X)$ -typical. Finally, the set U is the set of the vertices used before step i .

For the second phase, assume that $V(F_B^L) \neq \emptyset$ (otherwise we shall skip the second phase). We define the second phase of our embedding process in $|\bar{S}D^B|$ steps.

In each step $i \geq 1$, we embed the trees $\mathcal{T}^i := \{T \in \mathcal{T}_B^L : sd(T) = r_i\}$ in edges incident with L' . (Recall that $L' = L \setminus M$.) Suppose that we are at step i of this procedure, i. e. that we have already embedded the trees from $\mathcal{T}^1, \dots, \mathcal{T}^{i-1}$. Denote by U_{i-1} the set of vertices used so far for the embedding. Let L'_i be the set of those clusters of L' to which $\varphi(r_i)$ is typical. As $\varphi(r_i) \in B'$, (4) and (21) imply that

$$\text{d\ddot{e}g}_{L'_i}(B) \geq |V(\bigcup \mathcal{T}^i)| + |U_{i-1} \cap L'_i| + \frac{\pi k}{100N}|L'_i|.$$

Furthermore, by (9), for all $Y \in L'_i$ we have that

$$\text{d\ddot{e}g}(Y) \geq |V(\bigcup \mathcal{T}^i)| + |U_{i-1}| + \frac{\pi k}{100}.$$

Use Lemma 12 Part **B**) to embed \mathcal{T}_i , letting the rooted tree be the tree induced by r_i and the trees from \mathcal{T}^i , and setting $X := B$, $X' := B'$, $v := \varphi(r_i)$, $L_X := L'_i$, $N_{L_X} := N(L'_i)$, and $U := U_{i-1}$.

The third phase of our embedding process takes place in $|SD^A|$ steps, where in each step $i \geq 1$, we embed the trees from $\mathcal{T}^i := \{T \in \bar{\mathcal{T}}_A : sd(T) = r_i\}$. Suppose that we are at step i of this procedure, i. e. that we have already embedded the trees from $\mathcal{T}^1, \dots, \mathcal{T}^{i-1}$. Denote by \bar{U}_{i-1} the set of vertices used so far for the embedding. Let M_i be the maximal submatching of M such that $\varphi(r_i)$ is typical to all cluster of $V(M_i)$. As $\varphi(r_i) \in A'$, we have by (4) and (10) that

$$\text{deg}_{M_i}(A) \geq |V(\bigcup \mathcal{T}^i)| + |\bar{U}_i| + \frac{\pi k}{100}.$$

Observe that, as each edge $CD \in M$ meets $N(A)$ in at most one end-cluster, the set U_i trivially has Property (\diamond) in CD with respect to A . We use Lemma 12 Part **A**) to embed \mathcal{T}_i , letting (T, r) be the tree induced by r_i together with the trees from \mathcal{T}^i , and setting $X := A$, $X' := A'$, $v := \varphi(r_i)$, $M_X := M_i$, and $U := \bar{U}_{i-1}$.

This terminated our embedding of T^* , and thus the proof of Theorem 2.

4 Extensions and applications

In this last section, we explore applications and generalisations of Theorem 2. In Section 4.1 we show how our theorem implies an asymptotic upper bound on the Ramsey number of trees. We extend Theorem 2 so that it allows for embedding subgraphs other than trees in Section 4.2.

4.1 A bound on the Ramsey number of trees

Recall that $r(\mathcal{H}, \mathcal{H}')$ denotes the Ramsey number for the classes \mathcal{H} and \mathcal{H}' of graphs, and that \mathcal{T}_ℓ denotes the class of trees of order ℓ .

Based on ideas from [6] and using Theorem 2, we prove Proposition 3, which stated that $r(\mathcal{T}_{k+1}, \mathcal{T}_{m+1}) \leq k + m + o(k + m)$. The sharp bound $k + m$ has been conjectured in [6].

Proof of Proposition 3. Given $0 < \varepsilon < 1/4$, we apply Theorem 2 to $\eta = q = \varepsilon/4$ to obtain an $n_0 \in \mathbb{N}$. Now, let $n \geq n_0$, and let G be a graph on $n' = (1 + 2\varepsilon)n + 1$ vertices. Let k and m be such that $k + m = n$.

Clearly, either at least half of the vertices of G have degree at least $k + \varepsilon n$, or in the complement \bar{G} of G , at least half of the vertices have degree at least $m + \varepsilon n$. First, suppose that the former of these assertions is true. Then it is easy to calculate that

$$k + \varepsilon n \geq (1 + \eta)(k + qn').$$

Thus, we may apply Theorem 2, which yields that each tree in $\mathcal{T}_{k+qn'+1}$ is a subgraph of G . Hence, also each tree in \mathcal{T}_{k+1} is a subgraph of G .

Now, assume that the second assertion from above holds, that is, in the complement \bar{G} of G , there are at least $\frac{1}{2}(1 + \varepsilon)n$ vertices of degree at least $m + \frac{\varepsilon}{2}n$. We then find all trees from \mathcal{T}_{m+1} as subgraphs in \bar{G} . This is done analogously. We have thus shown that for every $\varepsilon > 0$ there is an n_0 so that for all k, m with $k + m \geq n_0$, we have that $R(\mathcal{T}_{k+1}, \mathcal{T}_{m+1}) \leq (1 + 2\varepsilon)(k + m) + 1$. This proves Proposition 3. \square

4.2 Graphs with few cycles

The question we pursue in this subsection is whether the condition of Theorem 2 allows for embedding other graphs on $k + 1$ vertices, apart from trees. For instance, may we add an edge to our tree T^* and still embed it in G ? In Theorem 4 we show that we may indeed add constantly many edges, as long as our graph stays bipartite.

Observe that the argument for the bound on Ramsey number from Subsection 4.1 would apply here as well. We thus get an upper bound of $k+m+o(k+m)$ for the Ramsey numbers of graphs Q_k, Q_m as in Theorem 4, although the sharp bound does not hold (cf. the example given in the introduction).

Our proof of Theorem 4 follows closely the lines of the proof of Theorem 2. We embed a rooted spanning tree (T^*, R) of Q , and choosing φ carefully, we ensure the adjacencies for the edges from $E(Q) \setminus E(T^*)$.

Proof of Theorem 4. Set $\pi := \min\{\eta, q\}$ and set

$$\varepsilon' := \frac{\varepsilon^{c+1}}{c+4}, \quad \text{and} \quad m_0 := \frac{500}{\pi^2 q},$$

where ε is the constant from the proof of Theorem 2. As in the proof of Theorem 2, the regularity lemma applied to ε' , and m_0 , yields natural numbers N_0 and M'_0 . Set $M_0 := \max\{M'_0, c\}$, define β and p accordingly, and set

$$n_0 := \max \left\{ N_0, \frac{9M_0}{\beta} \cdot \left(\frac{8}{p} \right)^{c+1} \right\}.$$

Now, let G be a graph on $n \geq n_0$ vertices which satisfies the condition of Theorem 4, let $k \geq qn$, and let Q be a connected bipartite graph of order $k + 1$ with at most $k + c$ edges, with a spanning tree T^* . Fix a root R in T^* . Denote by M^* the subgraph of Q induced by the edges in $E(Q) \setminus E(T^*)$ and let N^* be the set of predecessors of $V(M^*)$ in the tree order of T^* .

We decompose T^* as in Section 3.3, with the difference that we now add the vertices from $V(M^*) \cup N^*$ to the sets A' and B' (from the definition of SD), depending on the parity of their distance in T^* to R . In this way, and since Q is bipartite, we obtain, after the switching, two independent sets $\bar{S}D^A$ and SD^B so that

$$|\bar{S}D^A| + |SD^B| \leq \frac{8}{\beta} + 8c < \frac{9}{\beta},$$

which is constant in n .

The definition of our the embedding φ is similar as in the proof of Theorem 2, except for some extra precautions we take for vertices from $V(M^*) \cup N^*$. At step i , for each vertex $r \in \bar{S}D^A$, define

$$N_r^i := \bigcap_{\ell=1}^j N(\varphi(x_\ell)) \cap A,$$

where x_1, \dots, x_j are the already embedded neighbours of r in $\bar{S}D^A$. If none of the neighbours of r in $\bar{S}D^A$ has been embedded before step i , then set $N_r^i := A$. Analogously define N_r^i for $r \in SD^B$.

In each step i of our embedding process, we shall ensure the following.

(i) If $r \in V(M^*)$ is not yet embedded, then $|N_r^i| \geq \left(\frac{p}{4}\right)^j s$,

where $j = j(r, i)$ is the number of neighbours of r in \bar{SD}^A resp. SD^B that have already been embedded before step i .

Observe that in step $i = 0$, either $N_r^0 = A$ or $N_r^0 = B$, and therefore condition (i) is met.

Suppose that at step $i \geq 1$ of our embedding process, we are about to embed a vertex $r = r_i \in V(M^*) \cup N^*$. Assume that $r \in \bar{SD}^A$ (the case when $r \in SD^B$ is analogous). Denote by x_1, \dots, x_ℓ the neighbours of r in $V(M^*)$ that have not been embedded yet.

Now, embed r in a vertex from N_r^{i-1} that satisfies the at most 3 conditions of typicality from the proof of Theorem 2, except the typicality w. r. t. B , which we replace with typicality w. r. t. each $N_{x_j}^{i-1}$, for $1 \leq j \leq \ell$. This is possible, since our embedding scheme and the condition on the number of edges of Q ensure that r has at most $c + 1$ neighbours in Q that are already embedded. Thus, it follows from (i) for $i - 1$ and for r , from (2), and from the choice of n_0 that there are at least

$$\left(\left(\frac{p}{4}\right)^{c+1} - (c+3)\varepsilon' \right) s - |\bar{SD}| + 1 \geq \frac{1}{2} \left(\frac{p}{4}\right)^{c+1} s - \frac{9}{\beta} + 1 \geq 1$$

unused typical vertices we can choose $\varphi(r)$ from.

Finally, observe that since we chose $\varphi(r)$ typical w. r. t. each $N_{x_j}^{i-1}$, we have ensured property (i) for i and for every $r' \in V(M^*)$ that is not yet embedded. This completes the proof of Theorem 4. \square

Acknowledgment

The authors would like to thank Martin Loeb1 for helpful discussions, Miklós Simonovits for his valuable comments, and an anonymous referee for many useful remarks that helped to improve the presentation of the paper.

References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, On a conjecture of Loeb1, In *Proc. of the 7th International Conference on Graph Theory, Combinatorics, and Algorithms*, pages 1135-1146, Wiley, New York, 1995.
- [2] C. Bazgan, H. Li and M. Woźniak, On the Loeb1-Komlós-Sós conjecture, *J. Graph Theory*, 34:269-276, 2000.
- [3] S. Brandt and E. Dobson, The Erdős-Sós conjecture for graphs of girth 5, *Discr. Math.*, 150: 411-414,1996.
- [4] O. Cooley, Proof of the Loeb1-Komlós-Sós Conjecture for large, dense graphs. Preprint 2008
- [5] R. Diestel, *Graph Theory* (3rd edition). Springer-Verlag, 2005.
- [6] P. Erdős, Z. Füredi, M. Loeb1 and V. T. Sós, Discrepancy of trees. *Studia Sci. Math. Hungar.*, 30(1-2):47-57, 1995.

- [7] P. Haxell, T. Łuczak and P. Tingley, Ramsey numbers for trees of small maximum degree. *Combinatorica*, 22(2):287-320, 2002.
- [8] J. Hladký and D. Piguet, Loebel–Komlós–Sós conjecture: dense case. Preprint 2008.
- [9] J. Komlós and M. Simonovits, Szemerédi’s regularity lemma and its applications in graph theory. In *Paul Erdős is eighty, (Keszthely, 1993)*, volume 2 of *Bolyai Soc. Math. Stud.*, pages 295-352, Budapest, 1996. János Bolyai Math. Soc.
- [10] D. Piguet and M. Stein, The Loebel–Komlós–Sós conjecture for trees of diameter 5 and other special cases. *Electr. J. of Comb.*, 15:R106, 2008.
- [11] J.-F. Saclé and M. Woźniak, A note on the Erdős–Sós conjecture for graphs without C_4 . *J. Combin. Theory (Series B)*, 70(2):229-234, 1997.
- [12] E. Szemerédi, Regular partitions of graphs. *Colloques Internationaux C.N.R.S. 260 – Problèmes Combinatoires et Théorie des Graphes, Orsay*, pages 399-401, 1976.
- [13] M. Woźniak, On the Erdős–Sós conjecture. *J. Graph Theory*, 21(2):229-234, 1996.
- [14] Y. Zhao, Proof of the $(n/2 - n/2 - n/2)$ conjecture for large n . Preprint.