

The Approximate Loebel–Komlós–Sós Conjecture IV: Embedding techniques and the proof of the main result

Jan Hladký^{*†} János Komlós[‡] Diana Piguet^{§¶}
Miklós Simonovits^{||} Maya Stein^{**†} Endre Szemerédi^{‡‡}

Abstract

This is the last paper of a series of four papers in which we prove the following relaxation of the Loebel–Komlós–Sós Conjecture: For every $\alpha > 0$ there exists a number k_0 such that for every $k > k_0$ every n -vertex graph G with at least $(\frac{1}{2} + \alpha)n$ vertices of degree at least $(1 + \alpha)k$ contains each tree T of order k as a subgraph.

In the first two papers of this series, we decomposed the host graph G , and found a suitable combinatorial structure inside the decomposition. In the third paper, we refined this structure, and proved that any graph satisfying the conditions of the above approximate version of the Loebel–Komlós–Sós Conjecture contains one of ten specific configurations. In this paper we embed the tree T in each of the ten configurations.

Mathematics Subject Classification: 05C35 (primary), 05C05 (secondary).

Keywords: extremal graph theory; Loebel–Komlós–Sós Conjecture; tree; regularity lemma; sparse graph; graph decomposition.

^{*} *Corresponding author.* Centre for Discrete Mathematics and its Applications (DIMAP) and Mathematics Institute, University of Warwick, UK. Email: honzahladky@gmail.com

[†]Supported by an EPSRC postdoctoral fellowship.

[‡]Department of Mathematics, Rutgers University, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA

[§]European Centre of Excellence NTIS, Faculty of Applied Sciences, University of West Bohemia.

[¶]Supported by the Marie Curie fellowship FIST, DFG grant TA 309/2-1, Czech Ministry of Education project 1M0545, EPSRC award EP/D063191/1, and EPSRC Additional Sponsorship EP/J501414/1. The work leading to this invention has received funding from the European Research Council under the European Unions Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 259385, and was supported by the European Regional Development Fund (ERDF), project NTIS - New Technologies for Information Society, European Centre of Excellence, CZ.1.05/1.1.00/02.0090.

^{||}Rényi Institute, Budapest, Hungary.

^{**}Centro de Modelamiento Matemático, Universidad de Chile, Santiago, Chile.

^{††}Supported by Fondecyt Iniciación grant 11090141 and Fondecyt Regular grant 1140766.

^{‡‡}Department of Computer Science, Rutgers University, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA

Contents

1	Introduction	1
2	Notation and preliminaries	1
2.1	General notation	1
2.2	Regular pairs	2
2.3	LKS graphs	2
3	Trees	3
4	Necessary facts and notation from [HKP^a, HKP^b, HKP^c]	7
4.1	Sparse decomposition	7
4.2	Shadows	9
5	Configurations	9
5.1	Common settings	9
5.2	The ten configurations	13
6	Embedding trees	17
6.1	Overview of the embedding procedures	17
6.1.1	Embedding overview for Configurations ($\diamond 2$)–($\diamond 5$)	17
6.1.2	Embedding overview for Configurations ($\diamond 6$)–($\diamond 7$)	18
6.1.3	Embedding overview for Configuration ($\diamond 8$)	20
6.1.4	Embedding overview for Configuration ($\diamond 9$)	21
6.1.5	Embedding overview for Configuration ($\diamond 10$)	22
6.2	The role of random splitting	23
6.3	Stochastic process Duplicate(ℓ)	23
6.4	Embedding small trees	24
6.5	Main embedding lemmas	30
6.5.1	Embedding in Configuration ($\diamond 1$)	30
6.5.2	Embedding in Configurations ($\diamond 2$)–($\diamond 5$)	30
6.5.3	Embedding in Configurations ($\diamond 6$)–($\diamond 10$)	40
7	Proof of Theorem 1.2	60
8	Theorem 1.2 algorithmically	63
9	Acknowledgments	63
	Symbol index	65
	General index	66
	Bibliography	67

1 Introduction

This is the fourth and last of a series of four papers [HKP⁺a, HKP⁺b, HKP⁺c, HKP⁺d] in which we provide an approximate solution of the Loeb–Komlós–Sós Conjecture. The conjecture reads as follows.

Conjecture 1.1 (Loeb–Komlós–Sós Conjecture 1995 [EFLS95]). *Suppose that G is an n -vertex graph with at least $n/2$ vertices of degree more than $k - 2$. Then G contains each tree of order k .*

We discuss the history and state of the art in detail in the first paper [HKP⁺a] of our series. Our main result, which we prove in the present paper, is the approximate solution of the Loeb–Komlós–Sós Conjecture, and reads as follows.

Theorem 1.2 (Main result). *For every $\alpha > 0$ there exists k_0 such that for any $k > k_0$ we have the following. Each n -vertex graph G with at least $(\frac{1}{2} + \alpha)n$ vertices of degree at least $(1 + \alpha)k$ contains each tree T of order k .*

In the first paper [HKP⁺a] we exposed the decomposition techniques (*the sparse decomposition*), and in the second paper [HKP⁺b], we found a rough combinatorial structure in the host graph G . In [HKP⁺c], we refined this structure, and obtained one of ten possible *configurations*, which appear in any graph satisfying the hypotheses of Theorem 1.2. These configurations will be reintroduced in Section 5. All the configurations are build up from basic elements which are inherited from the sparse decomposition.

In the present paper, we will embed the tree T in the thus prepared host graph G . Let us give a short outline of this procedure. First, we cut the tree into smaller subtrees, connected by few vertices. This will be done in Section 3.

We then develop techniques to embed the smaller subtrees in different building blocks of the configurations. Then, for each of the configurations, we show how to combine the embedding techniques for smaller trees to embed the whole tree T . All of this will be done in Section 6. For the reader not interested in the details, we mention that Section 6.1 contains a 5-page overview of the embedding procedures, with all the relevant ideas.

Finally, in Section 7, we prove Theorem 1.2, with the help of the main results from the earlier papers [HKP⁺a, HKP⁺b, HKP⁺c].

2 Notation and preliminaries

2.1 General notation

The set $\{1, 2, \dots, n\}$ of the first n positive integers is denoted by $[n]$. We frequently employ indexing by many indices. We write superscript indices in parentheses (such as $a^{(3)}$), as opposed to notation of powers (such as a^3). We use sometimes subscript to refer to parameters appearing in a fact/lemma/theorem. For example $\alpha_{T1.2}$ refers to the parameter α from Theorem 1.2. We omit rounding symbols when this does not affect the correctness of the arguments.

We use lower case Greek letters to denote small positive constants. The exception is the letter ϕ which is reserved for embedding of a tree T in a graph G , $\phi : V(T) \rightarrow V(G)$. The capital Greek letters are used for large constants.

We write $V(G)$ and $E(G)$ for the vertex set and edge set of a graph G , respectively. Further, $v(G) = |V(G)|$ is the order of G , and $e(G) = |E(G)|$ is its number of edges. If $X, Y \subseteq V(G)$ are

two, not necessarily disjoint, sets of vertices we write $e(X)$ for the number of edges induced by X , and $e(X, Y)$ for the number of ordered pairs $(x, y) \in X \times Y$ such that $xy \in E(G)$. In particular, note that $2e(X) = e(X, X)$.

For a graph G , a vertex $v \in V(G)$ and a set $U \subseteq V(G)$, we write $\deg(v)$ and $\deg(v, U)$ for the degree of v , and for the number of neighbours of v in U , respectively. We write $\deg^{\min}(G)$ for the minimum degree of G , $\deg^{\min}(U) := \min\{\deg(u) : u \in U\}$, and $\deg^{\min}(V_1, V_2) = \min\{\deg(u, V_2) : u \in V_1\}$ for two sets $V_1, V_2 \subseteq V(G)$. Similar notation is used for the maximum degree, denoted by $\deg^{\max}(G)$. The neighbourhood of a vertex v is denoted by $N(v)$. We set $N(U) := \bigcup_{u \in U} N(u)$. The symbol $-$ is used for two graph operations: if $U \subseteq V(G)$ is a vertex set then $G - U$ is the subgraph of G induced by the set $V(G) \setminus U$. If $H \subseteq G$ is a subgraph of G then the graph $G - H$ is defined on the vertex set $V(G)$ and corresponds to deletion of edges of H from G .

2.2 Regular pairs

In this section we introduce the notion of regular pairs which is central for Szemerédi's regularity lemma. We also list some simple properties of regular pairs that will be useful in our embedding process.

Given a graph H and a pair (U, W) of disjoint sets $U, W \subseteq V(H)$ the *density of the pair* (U, W) is defined as

$$d(U, W) := \frac{e(U, W)}{|U||W|}.$$

Similarly, for a bipartite graph G with colour classes U, W we talk about its *bipartite density* $d(G) = \frac{e(G)}{|U||W|}$. For a given $\varepsilon > 0$, a pair (U, W) of disjoint sets $U, W \subseteq V(H)$ is called an ε -regular pair if $|d(U, W) - d(U', W')| < \varepsilon$ for every $U' \subseteq U$, $W' \subseteq W$ with $|U'| \geq \varepsilon|U|$, $|W'| \geq \varepsilon|W|$. If the pair (U, W) is not ε -regular, then we call it ε -irregular. A stronger notion than regularity is that of super-regularity which we recall now. A pair (A, B) is (ε, γ) -super-regular if it is ε -regular, and we have $\deg^{\min}(A, B) \geq \gamma|B|$, and $\deg^{\min}(B, A) \geq \gamma|A|$. Note that then (A, B) has bipartite density at least γ .

The following well-known property of regular pairs will be useful.

Fact 2.1. *Suppose that (U, W) is an ε -regular pair of density d . Let $U' \subseteq U, W' \subseteq W$ be sets of vertices with $|U'| \geq \alpha|U|$, $|W'| \geq \alpha|W|$, where $\alpha > \varepsilon$. Then the pair (U', W') is a $2\varepsilon/\alpha$ -regular pair of density at least $d - \varepsilon$.*

2.3 LKS graphs

Write $\mathbf{LKS}(n, k, \alpha)$ for the class of all n -vertex graphs with at least $(\frac{1}{2} + \alpha)n$ vertices of degrees at least $(1 + \alpha)k$. With this notation Conjecture 1.1 states that every graph in $\mathbf{LKS}(n, k, 0)$ contains every tree from $\mathbf{trees}(k + 1)$.

Define $\mathbf{LKSmin}(n, k, \eta)$ as the set of all graphs $G \in \mathbf{LKS}(n, k, \eta)$ that are edge-minimal with respect to the membership in $\mathbf{LKS}(n, k, \eta)$.

Definition 2.2. *Let $\mathbf{LKSsmall}(n, k, \eta)$ be the class of those graphs $G \in \mathbf{LKS}(n, k, \eta)$ for which we have the following three properties:*

1. *All the neighbours of every vertex $v \in V(G)$ with $\deg(v) > \lceil (1 + 2\eta)k \rceil$ have degrees at most $\lceil (1 + 2\eta)k \rceil$.*

-
2. All the neighbours of every vertex of $\mathbb{S}_{\eta,k}(G)$ have degree exactly $\lceil(1 + \eta)k\rceil$.
 3. We have $e(G) \leq kn$.

3 Trees

In this section we will show how to partition any given tree into small subtrees, connected by only a few vertices; this is what we call an ℓ -fine partition. This notion is essential for our proof of Theorem 1.2, as we can embed these small subtrees one at a time.

Similar but simpler tree-cutting procedures were used earlier for the dense case of the Loeb–Komlós–Sós Conjecture [AKS95, HP, PS12, Zha11]. There, the small trees were embedded in regular pairs of a regularity lemma decomposition of the host graph G . Since here, we use the sparse decomposition instead, we had to take more care when cutting the tree. (In particular, features (h), (i), (j) of Definition 3.3 are needed for the more complex setting here.)

If T is a tree and $r \in V(T)$, then the pair (T, r) is a *rooted tree* with root r . We write $V_{\text{odd}}(T, r) \subseteq V(T)$ for the set of vertices of T of odd distance from r , and analogously define $V_{\text{even}}(T, r)$. Note that $r \in V_{\text{even}}(T, r)$. The distance between two vertices v_1 and v_2 in a tree is denoted by $\text{dist}(v_1, v_2)$.

We start with a simple well-known fact about the number of leaves in a tree. For completeness we include a proof.

Fact 3.1. *Let T be a tree with ℓ vertices of degree at least three. Then T has at least $\ell + 2$ leaves.*

Proof. Let D_1 be the set of leaves, D_2 the set of vertices of degree two and D_3 be the set of vertices of degree of at least three. Then

$$2(|D_1| + |D_2| + |D_3|) - 2 = 2v(T) - 2 = 2e(T) = \sum_{v \in V(T)} \deg(v) \geq |D_1| + 2|D_2| + 3|D_3| ,$$

and the statement follows. □

Let T be a tree rooted at r , inducing the partial order \preceq on $V(T)$ (with r as the minimal element). If $a \preceq b$ and $ab \in E(T)$ then we say b is a *child of a* and a is the *parent of b* . $\text{Ch}(a)$ denotes the set of children of a , and the parent of a vertex $b \neq r$ is denoted $\text{Par}(b)$. For a set $U \subseteq V(T)$ write $\text{Par}(U) := \bigcup_{u \in U \setminus \{r\}} \text{Par}(u) \setminus U$ and $\text{Ch}(U) := \bigcup_{u \in U} \text{Ch}(u) \setminus U$.

The next simple fact has already appeared in [Zha11, HP] (and most likely in some more classic texts as well). Nevertheless, for completeness we give a proof here.

Fact 3.2. *Let T be a tree with color-classes X and Y , and $v(T) \geq 2$. Then the set X contains at least $|X| - |Y| + 1$ leaves of T .*

Proof. Root T at an arbitrary vertex $r \in Y$. Let I be the set of internal vertices of T that belong to X . Each $v \in I$ has at least one immediate successor in the tree order induced by r . These successors are distinct for distinct $v \in I$ and all lie in $Y \setminus \{r\}$. Thus $|I| \leq |Y| - 1$. The claim follows. □

We say that a tree $T' \subseteq T$ is *induced* by a vertex $x \in V(T)$ if $V(T')$ is the up-closure of x in $V(T)$, i.e., $V(T') = \{v \in V(T) : x \preceq v\}$. We then write $T' = T(r, \uparrow x)$, or $T' = T(\uparrow x)$, if the root is obvious from the context and call T' an *end subtree*. Subtrees of T that are not end subtrees are called *internal subtrees*.

Let T be a tree rooted at r and let $T' \subseteq T$ be a subtree with $r \notin V(T')$. The *seed* of T' is the \preceq -maximal vertex $x \in V(T) \setminus V(T')$ such that $x \preceq v$ for all $v \in V(T')$. We write $\text{Seed}(T') = x$. A *fruit* in a rooted tree (T, r) is any vertex $u \in V(T)$ whose distance from r is even and at least four.

We can now state the most important definition of this section.

Definition 3.3 (*ℓ -fine partition*). *Let $T \in \mathbf{trees}(k)$ be a tree rooted at r . An ℓ -fine partition of T is a quadruple $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$, where $W_A, W_B \subseteq V(T)$ and $\mathcal{S}_A, \mathcal{S}_B$ are families of subtrees of T such that*

- (a) *the three sets W_A, W_B and $\{V(T^*)\}_{T^* \in \mathcal{S}_A \cup \mathcal{S}_B}$ partition $V(T)$,*
- (b) *$r \in W_A \cup W_B$,*
- (c) *$\max\{|W_A|, |W_B|\} \leq 336k/\ell$,*
- (d) *for $w_1, w_2 \in W_A \cup W_B$ the distance $\text{dist}(w_1, w_2)$ is odd if and only if one of them lies in W_A and the other one in W_B ,*
- (e) *$v(T^*) \leq \ell$ for every tree $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$,*
- (f) *$V(T^*) \cap N(W_B) = \emptyset$ for every $T^* \in \mathcal{S}_A$ and $V(T^*) \cap N(W_A) = \emptyset$ for every $T^* \in \mathcal{S}_B$,*
- (g) *each tree of $\mathcal{S}_A \cup \mathcal{S}_B$ has its seed in $W_A \cup W_B$,*
- (h) *$|V(T^*) \cap N(W_A \cup W_B)| \leq 2$ for each $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$,*
- (i) *if $V(T^*) \cap N(W_A \cup W_B)$ contains two distinct vertices y_1, y_2 for some $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$, then $\text{dist}(y_1, y_2) \geq 4$,*
- (j) *if $T_1, T_2 \in \mathcal{S}_A \cup \mathcal{S}_B$ are two internal subtrees of T such that $v_1 \in T_1$ precedes $v_2 \in T_2$ then $\text{dist}_T(v_1, v_2) > 2$,*
- (k) *\mathcal{S}_B does not contain any internal tree of T , and*
- (l) *$\sum_{\substack{T^* \in \mathcal{S}_A \\ \text{end tree of } T}} v(T^*) \geq \sum_{T^* \in \mathcal{S}_B} v(T^*)$.*

Remark 3.4. *Suppose that $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ is an ℓ -fine partition of a tree (T, r) , and suppose that $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$ is such that $|V(T^*) \cap N(W_A \cup W_B)| = 2$. Let us root T^* at the neighbour r_1 of its seed, and let r_2 be the other vertex of $V(T^*) \cap N(W_A \cup W_B)$. Then (d), (f), and (i) imply that r_2 is a fruit in (T^*, r_1) .*

The following is the main lemma of this section.

Lemma 3.5. *Let $T \in \mathbf{trees}(k)$ be a tree rooted at r and let $\ell \in \mathbb{N}$ with $\ell \leq k$. Then T has an ℓ -fine partition.*

Proof. First we shall use an inductive construction to get candidates for W_A , W_B , \mathcal{S}_A and \mathcal{S}_B , which we shall modify later on, so that they satisfy all the conditions required by Definition 3.3.

Set $T_0 := T$. Now, inductively for $i \geq 1$ choose a \preceq -maximal vertex $x_i \in V(T_{i-1})$ with the property that $v(T_{i-1}(\uparrow x_i)) > \ell$. We set $T_i := T_{i-1} - (V(T_{i-1}(\uparrow x_i)) \setminus \{x_i\})$. If, say at step $i = i_{\text{end}}$, no such x_i exists, then $v(T_{i-1}) \leq \ell$. In that case, set $x_i := r$, set $W_1 := \{x_i\}_{i=1}^{i_{\text{end}}}$ and terminate. The fact that $v(T_{i-1} - V(T_i)) \geq \ell$ for each $i < i_{\text{end}}$ implies that

$$|W_1| - 1 = i_{\text{end}} - 1 \leq k/\ell. \quad (3.1)$$

Let \mathcal{C} be the set of all components of the forest $T - W_1$. Observe that by the choice of the x_i each $T^* \in \mathcal{C}$ has order at most ℓ .

Let A and B be the colour classes of T such that $r \in A$. Now, choosing W_A as $W_1 \cap A$ and W_B as $W_1 \cap B$ and dividing \mathcal{C} adequately into sets \mathcal{S}_A and \mathcal{S}_B would yield a quadruple that satisfies conditions (a), (b), (c), (d), (e) and (g). In order to find also the remaining properties satisfied, we shall refine our tree partition by adding more vertices to W_1 , thus making the trees in $\mathcal{S}_A \cup \mathcal{S}_B$ smaller. In doing so, we have to be careful not to end up violating (c). We shall enlarge the set of cut vertices in several steps, accomplishing sequentially, in this order, also properties (h), (j), (f), (i), and in the last step at the same time (k) and (l). It will be easy to check that in each of the steps none of the previously established properties is lost, so we will not explicitly check them, except for (c).

For condition (h), first define T' as the subtree of T that contains all vertices of W_1 and all vertices that lie on paths in T which have both endvertices in W_1 . Now, if a subtree $T^* \in \mathcal{C}$ does not already satisfy (h) for W_1 , then $V(T^*) \cap V(T')$ must contain some vertices of degree at least three. We will add the set $Y(T^*)$ of all these vertices to W_1 . Formally, let Y be the union of the sets $Y(T^*)$ over all $T^* \in \mathcal{C}$, and set $W_2 := W_1 \cup Y$. Then the components of $T - W_2$ satisfy (h).

Let us bound the size of the set W_2 . For each $T^* \in \mathcal{C}$, note that by Fact 3.1 for $T^* \cap T'$, we know that $|Y(T^*)|$ is at most the number of leaves of $T^* \cap T'$ (minus two). On the other hand, each leaf of $T^* \cap T'$ has a child in W_1 (in T). As these children are distinct for different trees $T^* \in \mathcal{C}$, we find that $|Y| \leq |W_1|$ and thus

$$|W_2| \leq 2|W_1|. \quad (3.2)$$

Next, for condition (j), observe that by setting $W_3 := W_2 \cup \text{Par}_T(W_2)$ the components of $T - W_3$ fulfill (j). We have

$$|W_3| \leq 2|W_2| \stackrel{(3.2)}{\leq} 4|W_1|. \quad (3.3)$$

In order to ensure condition (f), let R^* be the set of the roots (\preceq -minimal vertices) of those components T^* of $T - W_3$ which contain neighbours of both colour classes of T . Setting $W_4 := W_3 \cup R^*$ we see that (f) is satisfied for W_4 . Furthermore, as for each vertex in R^* there is a distinct member of W_3 above it in the order on T , we obtain

$$|W_4| \leq 2|W_3| \stackrel{(3.3)}{\leq} 8|W_1|. \quad (3.4)$$

Next, we shall aim for a stronger version of property (i), namely,

(i') if $V(T^*) \cap N_T(W_A \cup W_B) = \{y_1, y_2\}$ with $y_1 \neq y_2$ for some $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$, then $\text{dist}(y_1, y_2) \geq 6$.

The reason for requiring this strengthening is that later we might introduce additional cut vertices which would “shorten T^* by two”.

Consider a component T^* of $T - W_4$ which is an internal tree of T . If T^* contains two distinct neighbours y_1, y_2 of W_4 such that $\text{dist}_{T^*}(y_1, y_2) < 6$, then we call T^* *short*. Observe that there are at most $|W_4|$ short trees, because each of these trees has a unique vertex from W_4 above it. Let $Z(T^*) \subseteq V(T^*)$ be the vertices on the path from y_1 to y_2 . Then $|Z(T^*)| \leq 6$. Letting Z be the union of the sets $Z(T^*)$ over all short trees in $T - W_4$, and set $W_5 := W_4 \cup Z$, we obtain

$$|W_5| \leq |W_4| + 6|W_4| \stackrel{(3.4)}{\leq} 56|W_4| \stackrel{(3.1)}{\leq} 112k/\ell. \quad (3.5)$$

We still need to ensure (k) and (l). To this end, consider the set \mathcal{C}' of all components of $T - W_5$. Set $\mathcal{C}'_A := \{T^* \in \mathcal{C}' : \text{Seed}(T^*) \in A\}$ and set $\mathcal{C}'_B := \mathcal{C}' \setminus \mathcal{C}'_A$. We assume that

$$\sum_{T^* \in \mathcal{C}'_A : T^* \text{ end tree of } T} v(T^*) \geq \sum_{T^* \in \mathcal{C}'_B : T^* \text{ end tree of } T} v(T^*), \quad (3.6)$$

as otherwise we can simply swap A and B . Now, for each $T^* \in \mathcal{C}'_B$ that is not a end subtree of T , set $X(T^*) := V(T^*) \cap N_T(W_5)$. Let X be the union of all such sets $X(T^*)$. Observe that

$$|X| \leq 2|W_5 \cap B| \leq 2|W_5|. \quad (3.7)$$

For $W := W_5 \cup X$, all internal trees of $T - W$ have their seeds in A . This will guarantee (k), and, together with (3.6), also (l).

Finally, set $W_A := W \cap A$ and $W_B := W \cap B$, and let \mathcal{S}_A and \mathcal{S}_B be the sets of those components of $T - W$ that have their seeds in W_A and W_B , respectively. By construction, $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ has all the properties of an ℓ -fine partition. In particular, for (c), we find with (3.5) and (3.7) that $|W| \leq |W_5| + 2|W_5 \cap B| \leq 336k/\ell$. \square

For an ℓ -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a rooted tree (T, r) , the trees $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$ are called *shrubs*. An *end shrub* is a shrub that is an end subtree. An *internal shrub* is a shrub that is an internal subtree. A *knag* is a component of the forest $T[W_A \cup W_B]$.

Remark 3.6. (i) In our proof of Theorem 1.2, we shall apply Lemma 3.5 to a tree $T_{\mathbb{T}1.2} \in \mathbf{trees}(k)$. The number $\ell_{\mathbb{L}3.5}$ will be linear in k , and thus (c) of Definition 3.3 tells us that the size of the sets W_A and W_B is bounded by an absolute constant.

(ii) Each internal tree in \mathcal{S}_A of an ℓ -fine partition has a unique vertex from W_A above it. Thus with $\ell_{\mathbb{L}3.5}$ as above also the number of internal trees in \mathcal{S}_A is bounded by an absolute constant. This need not be the case for the number of end trees. For instance, if $(T_{\mathbb{T}1.2}, r)$ is a star with $k - 1$ leaves and rooted at its centre r then $W_A = \{r\}$ while the $k - 1$ leaves of $T_{\mathbb{T}1.2}$ form the end shrubs in \mathcal{S}_A .

For an ℓ -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a rooted tree (T, r) , the trees $T^* \in \mathcal{S}_A \cup \mathcal{S}_B$ are called *shrubs*. An *end shrub* is a shrub which is an end subtree. An *internal shrub* is a shrub which is an internal subtree. A *knag* is a component of the forest $T[W_A \cup W_B]$. Suppose that $T^* \in \mathcal{S}_A$ is an internal shrub, and r^* its \preceq_r -minimal vertex. Then $T^* - r^*$ contains a unique component with a vertex from $N_T(W_A)$. We call this component *principal subshrub*, and the other components *peripheral subshrubs*.

Definition 3.7 (ordered skeleton). We say that the sequence (X_0, X_1, \dots, X_m) is an ordered skeleton of the ℓ -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a rooted tree (T, r) if

- X_0 is a knag and contains r , and all other X_i are either knags or shrubs,
- $V(\bigcup_{i \leq m} X_i) = V(T)$, and
- for each $i = 1, \dots, m$, the subgraph formed by $X_0 \cup X_1 \cup \dots \cup X_i$ is connected in T .

Directly from Definition 3.3 we get:

Lemma 3.8. Any ℓ -fine partition of any rooted tree has an ordered skeleton.

4 Necessary facts and notation from [HKP^a, HKP^b, HKP^c]

4.1 Sparse decomposition

We first recall the notion of dense spots and related concepts from the earlier papers.

Definition 4.1 ((m, γ) -dense spot, (m, γ) -nowhere-dense). An (m, γ) -dense spot in a graph G is a non-empty bipartite subgraph $D = (U, W; F)$ of G with $d(D) > \gamma$ and $\deg^{\min}(D) > m$. We call G (m, γ) -nowhere-dense if it does not contain any (m, γ) -dense spot.

Definition 4.2 ((m, γ) -dense cover). A (m, γ) -dense cover of a graph G is a family \mathcal{D} of edge-disjoint (m, γ) -dense spots such that $E(G) = \bigcup_{D \in \mathcal{D}} E(D)$.

The proofs of the following facts can be found in [HKP^b].

Fact 4.3. Let $(U, W; F)$ be a $(\gamma k, \gamma)$ -dense spot in a graph G of maximum degree at most Ωk . Then $\max\{|U|, |W|\} \leq \frac{\Omega}{\gamma} k$.

Fact 4.4. Let H be a graph of maximum degree at most Ωk , let $v \in V(H)$, and let \mathcal{D} be a family of edge-disjoint $(\gamma k, \gamma)$ -dense spots. Then less than $\frac{\Omega}{\gamma}$ dense spots from \mathcal{D} contain v .

In the following definition, note that a subset of a $(\Lambda, \varepsilon, \gamma, k)$ -avoiding set is also $(\Lambda, \varepsilon, \gamma, k)$ -avoiding.

Definition 4.5 ($(\Lambda, \varepsilon, \gamma, k)$ -avoiding set). Suppose that G is a graph and \mathcal{D} is a family of dense spots in G . A set $\mathfrak{A} \subseteq \bigcup_{D \in \mathcal{D}} V(D)$ is $(\Lambda, \varepsilon, \gamma, k)$ -avoiding with respect to \mathcal{D} if for every $\bar{U} \subseteq V(G)$ with $|\bar{U}| \leq \Lambda k$ the following holds that for all but at most εk vertices $v \in \mathfrak{A}$. There is a dense spot $D \in \mathcal{D}$ with $|\bar{U} \cap V(D)| \leq \gamma^2 k$ that contains v .

Through the two definitions below, we expose the most important tool in the proof of our main result (Theorem 1.2): the *sparse decomposition*. It generalises the notion of equitable partition from Szemerédi's regularity lemma.

Definition 4.6 ($(k, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -bounded decomposition). Let $\mathcal{V} = \{V_1, V_2, \dots, V_s\}$ be a partition of the vertex set of a graph G . We say that $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ is a $(k, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -bounded decomposition of G with respect to \mathcal{V} if the following properties are satisfied:

1. The elements of \mathbf{V} are disjoint subsets of $V(G)$.

4.1 Sparse decomposition

2. G_{reg} is a subgraph of $G - G_{\text{exp}}$ on the vertex set $\bigcup \mathbf{V}$. For each edge $xy \in E(G_{\text{reg}})$ there are distinct $C_x \ni x$ and $C_y \ni y$ from \mathbf{V} , and $G[C_x, C_y] = G_{\text{reg}}[C_x, C_y]$. Furthermore, $G[C_x, C_y]$ forms an ε -regular pair of density at least γ^2 .
3. We have $\nu k \leq |C| = |C'| \leq \varepsilon k$ for all $C, C' \in \mathbf{V}$.
4. \mathcal{D} is a family of edge-disjoint $(\gamma k, \gamma)$ -dense spots in $G - G_{\text{exp}}$. For each $D = (U, W; F) \in \mathcal{D}$ all the edges of $G[U, W]$ are covered by \mathcal{D} (but not necessarily by D).
5. If G_{reg} contains at least one edge between $C_1, C_2 \in \mathbf{V}$ then there exists a dense spot $D = (U, W; F) \in \mathcal{D}$ such that $C_1 \subseteq U$ and $C_2 \subseteq W$.
6. For all $C \in \mathbf{V}$ there is $V \in \mathcal{V}$ so that either $C \subseteq V \cap V(G_{\text{exp}})$ or $C \subseteq V \setminus V(G_{\text{exp}})$. For all $C \in \mathbf{V}$ and $D = (U, W; F) \in \mathcal{D}$ we have $C \cap U \in \{\emptyset, C\}$.
7. G_{exp} is a $(\gamma k, \gamma)$ -nowhere-dense subgraph of G with $\deg^{\min}(G_{\text{exp}}) > \rho k$.
8. \mathfrak{A} is a $(\Lambda, \varepsilon, \gamma, k)$ -avoiding subset of $V(G) \setminus \bigcup \mathbf{V}$ with respect to dense spots \mathcal{D} .

We say that the bounded decomposition $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ respects the avoiding threshold b if for each $C \in \mathbf{V}$ we either have $\deg^{\max}_G(C, \mathfrak{A}) \leq b$, or $\deg^{\min}_G(C, \mathfrak{A}) > b$.

The members of \mathbf{V} are called *clusters*. Define the *cluster graph* \mathbf{G}_{reg} as the graph on the vertex set \mathbf{V} that has an edge $C_1 C_2$ for each pair (C_1, C_2) which has density at least γ^2 in the graph G_{reg} .

Definition 4.7 ($(k, \Omega^{**}, \Omega^*, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -sparse decomposition). Let $\mathcal{V} = \{V_1, V_2, \dots, V_s\}$ be a partition of the vertex set of a graph G . We say that $\nabla = (\Psi, \mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ is a $(k, \Omega^{**}, \Omega^*, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -sparse decomposition of G with respect to V_1, V_2, \dots, V_s if the following holds.

1. $\Psi \subseteq V(G)$, $\deg^{\min}_G(\Psi) \geq \Omega^{**} k$, $\deg^{\max}_H(V(G) \setminus \Psi) \leq \Omega^* k$, where H is spanned by the edges of $\bigcup \mathcal{D}$, G_{exp} , and edges incident with Ψ ,
2. $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ is a $(k, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -bounded decomposition of $G - \Psi$ with respect to $V_1 \setminus \Psi, V_2 \setminus \Psi, \dots, V_s \setminus \Psi$.

If the parameters do not matter, we call ∇ simply a *sparse decomposition*, and similarly we speak about a *bounded decomposition*.

Fact 4.8 ([HKP⁺a, Fact 3.10]). Let $\nabla = (\Psi, \mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ be a $(k, \Omega^{**}, \Omega^*, \Lambda, \gamma, \varepsilon, \nu, \rho)$ -sparse decomposition of a graph G . Let $x \in V(G) \setminus \Psi$. Assume that $\mathbf{V} \neq \emptyset$, and let \mathbf{c} be the size of each of the members of \mathbf{V} . Then there are less than

$$\frac{2(\Omega^*)^2 k}{\gamma^2 \mathbf{c}} \leq \frac{2(\Omega^*)^2}{\gamma^2 \nu}$$

clusters $C \in \mathbf{V}$ with $\deg_{G_{\mathcal{D}}}(x, C) > 0$.

Definition 4.9 (captured edges). In the situation of Definition 4.7, we refer to the edges in $E(G_{\text{reg}}) \cup E(G_{\text{exp}}) \cup E_G(\Psi, V(G)) \cup E_G(\mathfrak{A}, \mathfrak{A} \cup \bigcup \mathbf{V})$ as captured by the sparse decomposition. We write G_{∇} for the subgraph of G on the same vertex set which consists of the captured edges. Likewise, the captured edges of a bounded decomposition $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$ of a graph G are those in $E(G_{\text{reg}}) \cup E(G_{\text{exp}}) \cup E_G(\mathfrak{A}, \mathfrak{A} \cup \bigcup \mathbf{V})$.

The last definition we need is the notion of a semiregular matching.

Definition 4.10 ((ε, d, ℓ) -semiregular matching). *A collection \mathcal{N} of pairs (A, B) with $A, B \subseteq V(H)$ is called an (ε, d, ℓ) -semiregular matching of a graph H if*

- (i) $|A| = |B| \geq \ell$ for each $(A, B) \in \mathcal{N}$,
- (ii) (A, B) induces in H an ε -regular pair of density at least d , for each $(A, B) \in \mathcal{N}$, and
- (iii) all involved sets A and B are pairwise disjoint.

Sometimes, when the parameters do not matter we simply write semiregular matching.

Fact 4.11 ([HKP⁺b, Fact 4.3]). *Suppose that \mathcal{M} is an (ε, d, ℓ) -semiregular matching in a graph H . Then $|C| \leq \frac{\deg^{\max}(H)}{d}$ for each $C \in \mathcal{V}(\mathcal{M})$.*

4.2 Shadows

We recall the notion of a shadow given in [HKP⁺c]. Given a graph H , a set $U \subseteq V(H)$, and a number ℓ we define inductively

$$\begin{aligned} \text{shadow}_H^{(0)}(U, \ell) &:= U, \text{ and} \\ \text{shadow}_H^{(i)}(U, \ell) &:= \{v \in V(H) : \deg_H(v, \text{shadow}_H^{(i-1)}(U, \ell)) > \ell\} \text{ for } i \geq 1. \end{aligned}$$

We abbreviate $\text{shadow}_H^{(1)}(U, \ell)$ as $\text{shadow}_H(U, \ell)$. Further, the graph H is omitted from the subscript if it is clear from the context. Note that the shadow of a set U might intersect U .

The proofs of the following facts can be found in [HKP⁺c].

Fact 4.12. *Suppose H is a graph with $\deg^{\max}(H) \leq \Omega k$. Then for each $\alpha > 0, i \in \{0, 1, \dots\}$, and each set $U \subseteq V(H)$, we have*

$$|\text{shadow}^{(i)}(U, \alpha k)| \leq \left(\frac{\Omega}{\alpha}\right)^i |U|.$$

Fact 4.13. *Let $\alpha, \gamma, Q > 0$ be three numbers such that $Q \geq 1$ and $16Q \leq \frac{\alpha}{\gamma}$. Suppose that H is a $(\gamma k, \gamma)$ -nowhere-dense graph, and let $U \subseteq V(H)$ with $|U| \leq Qk$. Then we have*

$$|\text{shadow}(U, \alpha k)| \leq \frac{16Q^2\gamma}{\alpha} k.$$

5 Configurations

5.1 Common settings

We repeat some common settings that already appeared in [HKP⁺c].

Setting 5.1. *We assume that the constants $\Lambda, \Omega^*, \Omega^{**}, k_0$ and $\hat{\alpha}, \gamma, \varepsilon, \varepsilon', \eta, \pi, \rho, \tau, d$ satisfy*

$$\eta \gg \frac{1}{\Omega^*} \gg \frac{1}{\Omega^{**}} \gg \rho \gg \gamma \gg d \geq \frac{1}{\Lambda} \geq \varepsilon \geq \pi \geq \hat{\alpha} \geq \varepsilon' \geq \nu \gg \tau \gg \frac{1}{k_0} > 0, \quad (5.1)$$

5.1 Common settings

and that $k \geq k_0$. Here, by writing $c > a_1 \gg a_2 \gg \dots \gg a_\ell > 0$ we mean that there exist non-decreasing functions $f_i : (0, c)^i \rightarrow (0, c)$ ($i = 1, \dots, \ell - 1$) such that for each $i \in [\ell - 1]$ we have $a_{i+1} < f_i(a_1, \dots, a_i)$.¹

Suppose that $G \in \mathbf{LKSSsmall}(n, k, \eta)$ is given together with its $(k, \Omega^{**}, \Omega^*, \Lambda, \gamma, \varepsilon', \nu, \rho)$ -sparse decomposition

$$\nabla = (\Psi, \mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A}),$$

with respect to the partition $\{\mathbb{S}_{\eta, k}(G), \mathbb{L}_{\eta, k}(G)\}$, and with respect to the avoiding threshold $\frac{\rho k}{100\Omega^*}$. We write

$$V_{\rightsquigarrow \mathfrak{A}} := \text{shadow}_{G_{\nabla} - \Psi}(\mathfrak{A}, \frac{\rho k}{100\Omega^*}) \quad \text{and} \quad \mathbf{V}_{\rightsquigarrow \mathfrak{A}} := \{C \in \mathbf{V} : C \subseteq V_{\rightsquigarrow \mathfrak{A}}\}. \quad (5.2)$$

The graph \mathbf{G}_{reg} is the corresponding cluster graph. Let \mathfrak{c} be the size of an arbitrary cluster in \mathbf{V} .² Let G_{∇} be the spanning subgraph of G formed by the edges captured by ∇ . There are two $(\varepsilon, d, \pi\mathfrak{c})$ -semiregular matchings \mathcal{M}_A and \mathcal{M}_B in $G_{\mathcal{D}}$, with the following properties (we abbreviate $\mathbb{X}\mathbb{A} := \mathbb{X}\mathbb{A}(\eta, \nabla, \mathcal{M}_A, \mathcal{M}_B)$, $\mathbb{X}\mathbb{B} := \mathbb{X}\mathbb{B}(\eta, \nabla, \mathcal{M}_A, \mathcal{M}_B)$, and $\mathbb{X}\mathbb{C} := \mathbb{X}\mathbb{C}(\eta, \nabla, \mathcal{M}_A, \mathcal{M}_B)$):

1. $V(\mathcal{M}_A) \cap V(\mathcal{M}_B) = \emptyset$,
2. $V_1(\mathcal{M}_B) \subseteq S^0$, where

$$S^0 := \mathbb{S}_{\eta, k}(G) \setminus (V(G_{\text{exp}}) \cup \mathfrak{A}), \quad (5.3)$$

3. for each $(X, Y) \in \mathcal{M}_A \cup \mathcal{M}_B$, there is a dense spot $(U, W; F) \in \mathcal{D}$ with $X \subseteq U$ and $Y \subseteq W$, and further, either $X \subseteq \mathbb{S}_{\eta, k}(G)$ or $X \subseteq \mathbb{L}_{\eta, k}(G)$, and $Y \subseteq \mathbb{S}_{\eta, k}(G)$ or $Y \subseteq \mathbb{L}_{\eta, k}(G)$,
4. for each $X_1 \in \mathcal{V}_1(\mathcal{M}_A \cup \mathcal{M}_B)$ there exists a cluster $C_1 \in \mathbf{V}$ such that $X_1 \subseteq C_1$, and for each $X_2 \in \mathcal{V}_2(\mathcal{M}_A \cup \mathcal{M}_B)$ there exists $C_2 \in \mathbf{V} \cup \{\mathbb{L}_{\eta, k}(G) \cap \mathfrak{A}\}$ such that $X_2 \subseteq C_2$,
5. each pair of the semiregular matching $\mathcal{M}_{\text{good}} := \{(X_1, X_2) \in \mathcal{M}_A : X_1 \cup X_2 \subseteq \mathbb{X}\mathbb{A}\}$ corresponds to an edge in \mathbf{G}_{reg} ,
6. $e_{G_{\nabla}}(\mathbb{X}\mathbb{A}, S^0 \setminus V(\mathcal{M}_A)) \leq \gamma kn$,
7. $e_{G_{\text{reg}}}(V(G) \setminus V(\mathcal{M}_A \cup \mathcal{M}_B)) \leq \gamma^2 kn$,
8. for the semiregular matching $\mathcal{N}_{\mathfrak{A}} := \{(X, Y) \in \mathcal{M}_A \cup \mathcal{M}_B : (X \cup Y) \cap \mathfrak{A} \neq \emptyset\}$ we have $e_{G_{\text{reg}}}(V(G) \setminus V(\mathcal{M}_A \cup \mathcal{M}_B), V(\mathcal{N}_{\mathfrak{A}})) \leq \gamma^2 kn$,
9. $|E(G) \setminus E(G_{\nabla})| \leq 2\rho kn$,
10. $|E(G_{\mathcal{D}}) \setminus (E(G_{\text{reg}}) \cup E_G[\mathfrak{A}, \mathfrak{A} \cup \bigcup \mathbf{V}])| \leq \frac{5}{4}\gamma kn$.

¹The precise relation between the parameters will be determined in [HKP⁺d].

²The number \mathfrak{c} is not defined when $\mathbf{V} = \emptyset$. However in that case \mathfrak{c} is never actually used.

5.1 Common settings

We write

$$V_+ := V(G) \setminus (S^0 \setminus V(\mathcal{M}_A \cup \mathcal{M}_B)) \quad (5.4)$$

$$= \mathbb{L}_{\eta,k}(G) \cup V(G_{\text{exp}}) \cup \mathfrak{A} \cup V(\mathcal{M}_A \cup \mathcal{M}_B), \quad (5.5)$$

$$L_{\#} := \mathbb{L}_{\eta,k}(G) \setminus \mathbb{L}_{\frac{9}{10}\eta,k}(G_{\nabla}), \text{ and} \quad (5.6)$$

$$V_{\text{good}} := V_+ \setminus (\Psi \cup L_{\#}), \quad (5.7)$$

$$\mathbb{Y}\mathbb{A} := \text{shadow}_{G_{\nabla}} \left(V_+ \setminus L_{\#}, \left(1 + \frac{\eta}{10}\right)k \right) \setminus \text{shadow}_{G-G_{\nabla}} \left(V(G), \frac{\eta}{100}k \right), \quad (5.8)$$

$$\mathbb{Y}\mathbb{B} := \text{shadow}_{G_{\nabla}} \left(V_+ \setminus L_{\#}, \left(1 + \frac{\eta}{10}\right)\frac{k}{2} \right) \setminus \text{shadow}_{G-G_{\nabla}} \left(V(G), \frac{\eta}{100}k \right), \quad (5.9)$$

$$V_{\neq\Psi} := (\mathbb{X}\mathbb{A} \cup \mathbb{X}\mathbb{B}) \cap \text{shadow}_G \left(\Psi, \frac{\eta}{100}k \right), \quad (5.10)$$

$$\mathbb{P}_{\mathfrak{A}} := \text{shadow}_{G_{\text{reg}}} (V(\mathcal{M}_{\mathfrak{A}}), \gamma k) \setminus V(\mathcal{M}_A \cup \mathcal{M}_B),$$

$$\mathbb{P}_1 := \text{shadow}_{G_{\text{reg}}} (V(G) \setminus V(\mathcal{M}_A \cup \mathcal{M}_B), \gamma k) \setminus V(\mathcal{M}_A \cup \mathcal{M}_B),$$

$$\mathbb{P} := (\mathbb{X}\mathbb{A} \setminus \mathbb{Y}\mathbb{A}) \cup ((\mathbb{X}\mathbb{A} \cup \mathbb{X}\mathbb{B}) \setminus \mathbb{Y}\mathbb{B}) \cup V_{\neq\Psi} \cup L_{\#} \cup \mathbb{P}_1$$

$$\cup \text{shadow}_{G_{\mathcal{D}} \cup G_{\nabla}} (V_{\neq\Psi} \cup L_{\#} \cup \mathbb{P}_{\mathfrak{A}} \cup \mathbb{P}_1, \frac{\eta^2 k}{10^5}),$$

$$\mathbb{P}_2 := \mathbb{X}\mathbb{A} \cap \text{shadow}_{G_{\nabla}} (S^0 \setminus V(\mathcal{M}_A), \sqrt{\gamma}k),$$

$$\mathbb{P}_3 := \mathbb{X}\mathbb{A} \cap \text{shadow}_{G_{\nabla}} (\mathbb{X}\mathbb{A}, \eta^3 k / 10^3),$$

$$\mathcal{F} := \{C \in \mathcal{V}(\mathcal{M}_A) : C \subseteq \mathbb{X}\mathbb{A}\} \cup \mathcal{V}_1(\mathcal{M}_B). \quad (5.11)$$

Lemma 5.2. *For each $p \in \mathbb{N}$ and $a > 0$ there exists $k_0 > 0$ such that for each $k > k_0$ we have the following.*

Suppose G is a graph of order $n \geq k_0$ and $\deg^{\max}(G) \leq \Omega^* k$ with its $(k, \Lambda, \gamma, \varepsilon, k^{-0.05}, \rho)$ -bounded decomposition $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$. As usual, we write G_{∇} for the subgraph captured by $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$, and $G_{\mathcal{D}}$ for the spanning subgraph of G consisting of the edges in \mathcal{D} . Let \mathcal{M} be an $(\varepsilon, d, k^{0.95})$ -semiregular matching in G , and $\mathfrak{U}_1, \dots, \mathfrak{U}_p$ be subsets of $V(G)$. Suppose that $\Omega^* \geq 1$ and $\Omega^*/\gamma < k^{0.1}$.

Suppose that $\mathfrak{q}_1, \dots, \mathfrak{q}_p \in \{0\} \cup [a, 1]$ are reals with $\sum \mathfrak{q}_i \leq 1$. Then there exists a partition $\mathfrak{Q}_1 \cup \dots \cup \mathfrak{Q}_p = V(G)$, and sets $\bar{V} \subseteq V(G)$, $\bar{\mathcal{V}} \subseteq \mathcal{V}(\mathcal{M})$, $\bar{\mathbf{V}} \subseteq \mathbf{V}$ with the following properties.

$$(1) |\bar{V}| \leq \exp(-k^{0.1})n, |\bigcup \bar{\mathcal{V}}| \leq \exp(-k^{0.1})n, |\bigcup \bar{\mathbf{V}}| < \exp(-k^{0.1})n.$$

$$(2) \text{ For each } i \in [p] \text{ and each } C \in \mathbf{V} \setminus \bar{\mathbf{V}} \text{ we have } |C \cap \mathfrak{Q}_i| \geq \mathfrak{q}_i |\mathfrak{Q}_i| - k^{0.9}.$$

$$(3) \text{ For each } i \in [p] \text{ and each } C \in \mathcal{V}(\mathcal{M}) \setminus \bar{\mathcal{V}} \text{ we have } |C \cap \mathfrak{Q}_i| \geq \mathfrak{q}_i |\mathfrak{Q}_i| - k^{0.9}.$$

$$(4) \text{ For each } i \in [p], D = (U, W; F) \in \mathcal{D} \text{ and } \deg^{\min}_D(U \setminus \bar{V}, W \cap \mathfrak{Q}_i) \geq \mathfrak{q}_i \gamma k - k^{0.9}.$$

$$(5) \text{ For each } i, j \in [p] \text{ we have } |\mathfrak{Q}_i \cap \mathfrak{U}_j| \geq \mathfrak{q}_i |\mathfrak{U}_j| - n^{0.9}.$$

$$(6) \text{ For each } i \in [p] \text{ each } J \subseteq [p] \text{ and each } v \in V(G) \setminus \bar{V} \text{ we have}$$

$$\deg_H(v, \mathfrak{Q}_i \cap \mathfrak{U}_J) \geq \mathfrak{q}_i \deg_H(v, \mathfrak{U}_J) - 2^{-p} k^{0.9},$$

for each of the graphs $H \in \{G, G_{\nabla}, G_{\text{exp}}, G_{\mathcal{D}}, G_{\nabla} \cup G_{\mathcal{D}}\}$, where $\mathfrak{U}_J := (\bigcap_{j \in J} \mathfrak{U}_j) \setminus (\bigcup_{j \in [p] \setminus J} \mathfrak{U}_j)$.

5.1 Common settings

(7) For each $i, i', j, j' \in [p]$ ($j \neq j'$), we have

$$\begin{aligned} e_H(\mathfrak{Q}_i \cap \mathfrak{U}_j, \mathfrak{Q}_{i'} \cap \mathfrak{U}_{j'}) &\geq \mathfrak{q}_i \mathfrak{q}_{i'} e_H(\mathfrak{U}_j, \mathfrak{U}_{j'}) - k^{0.6} n^{0.6}, \\ e_H(\mathfrak{Q}_i \cap \mathfrak{U}_j, \mathfrak{Q}_{i'} \cap \mathfrak{U}_j) &\geq \mathfrak{q}_i \mathfrak{q}_{i'} e(H[\mathfrak{U}_j]) - k^{0.6} n^{0.6} \quad \text{if } i \neq i', \text{ and} \\ e(H[\mathfrak{Q}_i \cap \mathfrak{U}_j]) &\geq \mathfrak{q}_i^2 e(H[\mathfrak{U}_j]) - k^{0.6} n^{0.6}. \end{aligned}$$

for each of the graphs $H \in \{G, G_\nabla, G_{\text{exp}}, G_{\mathcal{D}}, G_\nabla \cup G_{\mathcal{D}}\}$.

(8) For each $i \in [p]$ if $\mathfrak{q}_i = 0$ then $\mathfrak{Q}_i = \emptyset$.

Definition 5.3 (Proportional splitting). Let $\mathfrak{p}_0, \mathfrak{p}_1, \mathfrak{p}_2 > 0$ be three positive reals with $\sum_i \mathfrak{p}_i \leq 1$. Under Setting 5.1, suppose that $(\mathfrak{P}_0, \mathfrak{P}_1, \mathfrak{P}_2)$ is a partition of $V(G) \setminus \Psi$ which satisfies assertions of Lemma 5.2 with parameter $p_{L5.2} := 10$ for graph $G_{L5.2}^* := (G_\nabla - \Psi) \cup G_{\mathcal{D}}$ (here, by the union, we mean union of the edges), bounded decomposition $(\mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$, matching $\mathcal{M}_{L5.2} := \mathcal{M}_A \cup \mathcal{M}_B$, sets $\mathfrak{U}_1 := V_{\text{good}}, \mathfrak{U}_2 := \mathbb{X}\mathbb{A} \setminus (\Psi \cup \mathbb{P}), \mathfrak{U}_3 := \mathbb{X}\mathbb{B} \setminus \mathbb{P}, \mathfrak{U}_4 := V(G_{\text{exp}}), \mathfrak{U}_5 := \mathfrak{A}, \mathfrak{U}_6 := V_{\rightsquigarrow \mathfrak{A}}, \mathfrak{U}_7 := \mathbb{P}_{\mathfrak{A}}, \mathfrak{U}_8 := \mathbb{L}_{\eta, k}(G), \mathfrak{U}_9 := L_{\#}, \mathfrak{U}_{10} := V_{\neq \Psi}$ and reals $\mathfrak{q}_1 := \mathfrak{p}_0, \mathfrak{q}_2 := \mathfrak{p}_1, \mathfrak{q}_3 := \mathfrak{p}_2, \mathfrak{q}_4 := \dots \mathfrak{q}_{10} = 0$. Note that by Lemma 5.2(8) we have that $(\mathfrak{P}_0, \mathfrak{P}_1, \mathfrak{P}_2)$ is a partition of $V(G) \setminus \Psi$. We call $(\mathfrak{P}_0, \mathfrak{P}_1, \mathfrak{P}_2)$ proportional $(\mathfrak{p}_0 : \mathfrak{p}_1 : \mathfrak{p}_2)$ splitting.

We refer to properties of the proportional $(\mathfrak{p}_0 : \mathfrak{p}_1 : \mathfrak{p}_2)$ splitting $(\mathfrak{P}_0, \mathfrak{P}_1, \mathfrak{P}_2)$ using the numbering of Lemma 5.2; for example, ‘‘Definition 5.3(5)’’ tells us among other things that $|(\mathbb{X}\mathbb{A} \setminus \mathbb{P}) \cap \mathfrak{P}_0| \geq \mathfrak{p}_0 |\mathbb{X}\mathbb{A} \setminus (\mathbb{P} \cup \Psi)| - n^{0.9}$.

Setting 5.4. Under Setting 5.1, suppose that we are given a proportional $(\mathfrak{p}_0 : \mathfrak{p}_1 : \mathfrak{p}_2)$ splitting $(\mathfrak{P}_0, \mathfrak{P}_1, \mathfrak{P}_2)$ of $V(G) \setminus \Psi$. We assume that

$$\mathfrak{p}_0, \mathfrak{p}_1, \mathfrak{p}_2 \geq \frac{\eta}{100}. \quad (5.12)$$

Let $\bar{V}, \bar{\mathcal{V}}, \bar{\mathbf{V}}$ be the exceptional sets as in Definition 5.3(1).

We write

$$\mathbb{F} := \text{shadow}_{G_{\mathcal{D}}} \left(\bigcup \bar{\mathcal{V}} \cup \bigcup \bar{\mathcal{V}}^* \cup \bigcup \bar{\mathbf{V}}, \frac{\eta^2 k}{10^{10}} \right), \quad (5.13)$$

where $\bar{\mathcal{V}}^*$ are the partners of $\bar{\mathcal{V}}$ in $\mathcal{M}_A \cup \mathcal{M}_B$.

We have

$$|\mathbb{F}| \leq \varepsilon n. \quad (5.14)$$

For an arbitrary set $U \subseteq V(G)$ and for $i \in \{0, 1, 2\}$ we write $U^{\uparrow i}$ for the set $U \cap \mathfrak{P}_i$.

For each $(X, Y) \in \mathcal{M}_A \cup \mathcal{M}_B$ such that $X, Y \notin \bar{\mathcal{V}}$ we write $(X, Y)^{\uparrow i}$ for an arbitrary fixed pair $(X' \subseteq X, Y' \subseteq Y)$ with the property that $|X'| = |Y'| = \min\{|X^{\uparrow i}|, |Y^{\uparrow i}|\}$. We extend this notion of restriction to an arbitrary semiregular matching $\mathcal{N} \subseteq \mathcal{M}_A \cup \mathcal{M}_B$ as follows. We set

$$\mathcal{N}^{\uparrow i} := \{(X, Y)^{\uparrow i} : (X, Y) \in \mathcal{N} \text{ with } X, Y \notin \bar{\mathcal{V}}\}.$$

In [HKP⁺c] it was shown that the above setting yields the following property.

Lemma 5.5 ([HKP⁺c, Lemma 3.8]). Assume Setting 5.4. Then for each $i \in \{0, 1, 2\}$, and for each $\mathcal{N} \subseteq \mathcal{M}_A \cup \mathcal{M}_B$ we have that $\mathcal{N}^{\uparrow i}$ is a $(\frac{400\varepsilon}{\eta}, \frac{d}{2}, \frac{\eta\pi}{200}\mathfrak{c})$ -semiregular matching satisfying

$$|V(\mathcal{N}^{\uparrow i})| \geq \mathfrak{p}_i |V(\mathcal{N})| - 2k^{-0.05}n. \quad (5.15)$$

Moreover for all $v \notin \mathbb{F}$ and for all $i = 0, 1, 2$ we have $\deg_{G_{\mathcal{D}}}(v, V(\mathcal{N})^{\uparrow i} \setminus V(\mathcal{N}^{\uparrow i})) \leq \frac{\eta^2 k}{10^5}$.

5.2 The ten configurations

We start by giving definitions of Configuration $(\diamond 1)$. This is a very easy configuration in which a modification of the greedy tree-embedding strategy works.

Definition 5.6 (Configuration $(\diamond 1)$). *We say that a graph G is in Configuration $(\diamond 1)$ if there exists a non-empty bipartite graph $H \subseteq G$ with $\deg^{\min}_G(V(H)) \geq k$ and $\deg^{\min}(H) \geq k/2$.*

We now introduce the configurations $(\diamond 2)$ – $(\diamond 5)$ which make use of the set Ψ . These configurations build on Preconfiguration (\clubsuit) .

Definition 5.7 (Preconfiguration (\clubsuit)). *Suppose that we are in Setting 5.1. We say that the graph G is in Preconfiguration $(\clubsuit)(\Omega^*)$ if the following conditions are met. G contains non-empty sets $L'' \subseteq L' \subseteq \mathbb{L}_{\frac{9}{10}\eta, k}(G_{\nabla}) \setminus \Psi$, and a non-empty set $\Psi' \subseteq \Psi$ such that*

$$\deg^{\max}_{G_{\nabla}}(L', \Psi \setminus \Psi') < \frac{\eta k}{100}, \quad (5.16)$$

$$\deg^{\min}_{G_{\nabla}}(\Psi', L'') \geq \Omega^* k, \text{ and} \quad (5.17)$$

$$\deg^{\max}_{G_{\nabla}}(L'', \mathbb{L}_{\frac{9}{10}\eta, k}(G_{\nabla}) \setminus (\Psi \cup L')) \leq \frac{\eta k}{100}. \quad (5.18)$$

Definition 5.8 (Configuration $(\diamond 2)$). *Suppose that we are in Setting 5.1. We say that the graph G is in Configuration $(\diamond 2)(\Omega^*, \tilde{\Omega}, \beta)$ if the following conditions are met.*

The triple L'', L', Ψ' witnesses preconfiguration $(\clubsuit)(\Omega^)$ in G . There exist a non-empty set $\Psi'' \subseteq \Psi'$, a set $V_1 \subseteq V(G_{\text{exp}}) \cap \mathbb{YB} \cap L''$, and a set $V_2 \subseteq V(G_{\text{exp}})$ with the following properties.*

$$\deg^{\min}_{G_{\nabla}}(\Psi'', V_1) \geq \tilde{\Omega} k$$

$$\deg^{\min}_{G_{\nabla}}(V_1, \Psi'') \geq \beta k,$$

$$\deg^{\min}_{G_{\text{exp}}}(V_1, V_2) \geq \beta k,$$

$$\deg^{\min}_{G_{\text{exp}}}(V_2, V_1) \geq \beta k.$$

Definition 5.9 (Configuration $(\diamond 3)$). *Suppose that we are in Setting 5.1. We say that the graph G is in Configuration $(\diamond 3)(\Omega^*, \tilde{\Omega}, \zeta, \delta)$ if the following conditions are met.*

The triple L'', L', Ψ' witnesses preconfiguration $(\clubsuit)(\Omega^)$ in G . There exist a non-empty set $\Psi'' \subseteq \Psi'$, a set $V_1 \subseteq \mathfrak{A} \cap \mathbb{YB} \cap L''$, and a set $V_2 \subseteq V(G) \setminus \Psi$ such that the following properties are satisfied.*

$$\deg^{\min}_{G_{\nabla}}(\Psi'', V_1) \geq \tilde{\Omega} k,$$

$$\deg^{\min}_{G_{\nabla}}(V_1, \Psi'') \geq \delta k,$$

$$\deg^{\max}_{G_{\mathcal{D}}}(V_1, V(G) \setminus (V_2 \cup \Psi)) \leq \zeta k, \quad (5.19)$$

$$\deg^{\min}_{G_{\mathcal{D}}}(V_2, V_1) \geq \delta k. \quad (5.20)$$

Definition 5.10 (Configuration $(\diamond 4)$). *Suppose that we are in Setting 5.1. We say that the graph G is in Configuration $(\diamond 4)(\Omega^*, \tilde{\Omega}, \zeta, \delta)$ if the following conditions are met.*

5.2 The ten configurations

The triple L'', L', Ψ' witnesses preconfiguration $(\clubsuit)(\Omega^*)$ in G . There exists a non-empty set $\Psi'' \subseteq \Psi'$, sets $V_1 \subseteq \mathbb{YB} \cap L''$, $\mathfrak{A}' \subseteq \mathfrak{A}$, and $V_2 \subseteq V(G) \setminus \Psi$ with the following properties

$$\begin{aligned} \deg_{G_{\nabla}}^{\min}(\Psi'', V_1) &\geq \tilde{\Omega}k, \\ \deg_{G_{\nabla}}^{\min}(V_1, \Psi'') &\geq \delta k, \\ \deg_{G_{\nabla} \cup G_{\mathcal{D}}}^{\min}(V_1, \mathfrak{A}') &\geq \delta k, \end{aligned} \tag{5.21}$$

$$\deg_{G_{\nabla} \cup G_{\mathcal{D}}}^{\min}(\mathfrak{A}', V_1) \geq \delta k, \tag{5.22}$$

$$\deg_{G_{\nabla} \cup G_{\mathcal{D}}}^{\min}(V_2, \mathfrak{A}') \geq \delta k, \tag{5.23}$$

$$\deg_{G_{\nabla} \cup G_{\mathcal{D}}}^{\max}(\mathfrak{A}', V(G) \setminus (\Psi \cup V_2)) \leq \zeta k. \tag{5.24}$$

Definition 5.11 (Configuration $(\diamond 5)$). Suppose that we are in Setting 5.1. We say that the graph G is in Configuration $(\diamond 5)(\Omega^*, \tilde{\Omega}, \delta, \zeta, \tilde{\pi})$ if the following conditions are met.

The triple L'', L', Ψ' witnesses preconfiguration $(\clubsuit)(\Omega^*)$ in G . There exists a non-empty set $\Psi'' \subseteq \Psi'$, and a set $V_1 \subseteq (\mathbb{YB} \cap L'' \cap \mathbf{V}) \setminus V(G_{\text{exp}})$ such that the following conditions are fulfilled.

$$\deg_{G_{\nabla}}^{\min}(\Psi'', V_1) \geq \tilde{\Omega}k, \tag{5.25}$$

$$\deg_{G_{\nabla}}^{\min}(V_1, \Psi'') \geq \delta k, \tag{5.26}$$

$$\deg_{G_{\text{reg}}}^{\min}(V_1) \geq \zeta k. \tag{5.27}$$

Further, we have

$$C \cap V_1 = \emptyset \text{ or } |C \cap V_1| \geq \tilde{\pi}|C| \tag{5.28}$$

for every $C \in \mathbf{V}$.

It remains to introduce configurations $(\diamond 6)$ – $(\diamond 10)$. In these configurations the set Ψ is not utilized. All these configurations make use of Setting 5.4, i.e., the set $V(G) \setminus \Psi$ is partitioned into three sets $\mathfrak{P}_0, \mathfrak{P}_1$ and \mathfrak{P}_2 . The purpose of $\mathfrak{P}_0, \mathfrak{P}_1$ and \mathfrak{P}_2 is to make possible to embed the knags, the internal shrubs, and the end shrubs of $T_{T1.2}$, respectively. Thus the parameters $\mathfrak{p}_0, \mathfrak{p}_1$ and \mathfrak{p}_2 are chosen proportionally to the sizes of these respective parts of $T_{T1.2}$.

We first introduce four preconfigurations $(\heartsuit 1)$, $(\heartsuit 2)$, (\mathbf{exp}) and (\mathbf{reg}) which are building bricks for configurations $(\diamond 6)$ – $(\diamond 9)$. The preconfigurations $(\heartsuit 1)$ and $(\heartsuit 2)$ will be used for embedding end shrubs of a fine partition of the tree $T_{T1.2}$, and preconfigurations (\mathbf{exp}) and (\mathbf{reg}) will be used for embedding its knags.

An \mathcal{M} -cover of a semiregular matching \mathcal{M} is a family $\mathcal{F} \subseteq \mathcal{V}(\mathcal{M})$ with the property that at least one of the elements S_1 and S_2 is a member of \mathcal{F} , for each $(S_1, S_2) \in \mathcal{M}$.

Definition 5.12 (Preconfiguration $(\heartsuit 1)$). Suppose that we are in Setting 5.1 and Setting 5.4. We say that the graph G is in Preconfiguration $(\heartsuit 1)(\gamma', h)$ of $V(G)$ if there are two non-empty sets $V_0, V_1 \subseteq \mathfrak{P}_0 \setminus \left(\mathbb{F} \cup \text{shadow}_{G_{\mathcal{D}}}(V_{\mathcal{A}\Psi}, \frac{\eta^2 k}{10^5}) \right)$ with the following properties.

$$\deg_{G_{\nabla}}^{\min}(V_0, V_{\text{good}}^{\uparrow 2}) \geq h/2, \text{ and} \tag{5.29}$$

$$\deg_{G_{\nabla}}^{\min}(V_1, V_{\text{good}}^{\uparrow 2}) \geq h. \tag{5.30}$$

Further, there is an $(\mathcal{M}_A \cup \mathcal{M}_B)$ -cover \mathcal{F} such that

$$\deg_{G_{\nabla}}^{\max}(V_1, \bigcup \mathcal{F}) \leq \gamma'k. \tag{5.31}$$

Definition 5.13 (Preconfiguration (♡2)). Suppose that we are in Setting 5.1 and Setting 5.4. We say that the graph G is in Preconfiguration (♡2)(h) of $V(G)$ if there are two non-empty sets $V_0, V_1 \subseteq \mathfrak{P}_0 \setminus \left(\mathbb{F} \cup \text{shadow}_{G_{\mathcal{D}}}(V_{\mathcal{A}\Psi}, \frac{\eta^2 k}{10^5}) \right)$ with the following properties.

$$\deg^{\min}_{G_{\nabla}}(V_0 \cup V_1, V_{\text{good}}^2) \geq h. \quad (5.32)$$

Definition 5.14 (Preconfiguration (exp)). Suppose that we are in Setting 5.1 and Setting 5.4. We say that the graph G is in Preconfiguration (exp)(β) if there are two non-empty sets $V_0, V_1 \subseteq \mathfrak{P}_0$ with the following properties.

$$\deg^{\min}_{G_{\text{exp}}}(V_0, V_1) \geq \beta k, \quad (5.33)$$

$$\deg^{\min}_{G_{\text{exp}}}(V_1, V_0) \geq \beta k. \quad (5.34)$$

Definition 5.15 (Preconfiguration (reg)). Suppose that we are in Setting 5.1 and Setting 5.4. We say that the graph G is in Preconfiguration (reg)($\tilde{\varepsilon}, d', \mu$) if there are two non-empty sets $V_0, V_1 \subseteq \mathfrak{P}_0$ and a non-empty family of vertex-disjoint ($\tilde{\varepsilon}, d'$)-super-regular pairs $\{(Q_0^{(j)}, Q_1^{(j)})\}_{j \in \mathcal{Y}}$ (with respect to the edge set $E(G)$) with $V_0 := \bigcup Q_0^{(j)}$ and $V_1 := \bigcup Q_1^{(j)}$ such that

$$\min \left\{ |Q_0^{(j)}|, |Q_1^{(j)}| \right\} \geq \mu k. \quad (5.35)$$

Definition 5.16 (Configuration (◇6)). Suppose that we are in Settings 5.1 and 5.4. We say that the graph G is in Configuration (◇6)($\delta, \tilde{\varepsilon}, d', \mu, \gamma', h_2$) if the following conditions are met.

The vertex sets V_0, V_1 witness Preconfiguration (reg)($\tilde{\varepsilon}, d', \mu$) or Preconfiguration (exp)(δ) and either Preconfiguration (♡1)(γ', h_2) or Preconfiguration (♡2)(h_2). There exist non-empty sets $V_2, V_3 \subseteq \mathfrak{P}_1$ such that

$$\deg^{\min}_G(V_1, V_2) \geq \delta k, \quad (5.36)$$

$$\deg^{\min}_G(V_2, V_1) \geq \delta k, \quad (5.37)$$

$$\deg^{\min}_{G_{\text{exp}}}(V_2, V_3) \geq \delta k, \text{ and} \quad (5.38)$$

$$\deg^{\min}_{G_{\text{exp}}}(V_3, V_2) \geq \delta k. \quad (5.39)$$

Definition 5.17 (Configuration (◇7)). Suppose that we are in Settings 5.1 and 5.4. We say that the graph G is in Configuration (◇7)($\delta, \rho', \tilde{\varepsilon}, d', \mu, \gamma', h_2$) if the following conditions are met.

The sets V_0, V_1 witness Preconfiguration (reg)($\tilde{\varepsilon}, d', \mu$) and either Preconfiguration (♡1)(γ', h_2) or Preconfiguration (♡2)(h_2). There exist non-empty sets $V_2 \subseteq \mathfrak{A}^1 \setminus \bar{V}$ and $V_3 \subseteq \mathfrak{P}_1$ such that

$$\deg^{\min}_G(V_1, V_2) \geq \delta k, \quad (5.40)$$

$$\deg^{\min}_G(V_2, V_1) \geq \delta k, \quad (5.41)$$

$$\deg^{\max}_{G_{\mathcal{D}}}(V_2, \mathfrak{P}_1 \setminus V_3) < \rho' k \text{ and} \quad (5.42)$$

$$\deg^{\min}_{G_{\mathcal{D}}}(V_3, V_2) \geq \delta k. \quad (5.43)$$

Definition 5.18 (Configuration (◇8)). Suppose that we are in Settings 5.1 and 5.4. We say that the graph G is in Configuration (◇8)($\delta, \rho', \varepsilon_1, \varepsilon_2, d_1, d_2, \mu_1, \mu_2, h_1, h_2$) if the following conditions are met.

5.2 The ten configurations

The vertex sets V_0, V_1 witness Preconfiguration $(\mathbf{reg})(\varepsilon_2, d_2, \mu_2)$ and Preconfiguration $(\heartsuit 2)(h_2)$. There exist non-empty sets $V_2 \subseteq \mathfrak{P}_0$, $V_3, V_4 \subseteq \mathfrak{P}_1$, $V_3 \subseteq \mathfrak{A} \setminus \bar{V}$, and an $(\varepsilon_1, d_1, \mu_1 k)$ -semiregular matching \mathcal{N} absorbed by $(\mathcal{M}_A \cup \mathcal{M}_B) \setminus \mathcal{N}_{\mathfrak{A}}$, $V(\mathcal{N}) \subseteq \mathfrak{P}_1 \setminus V_3$ such that

$$\deg_G^{\min}(V_1, V_2) \geq \delta k, \quad (5.44)$$

$$\deg_G^{\min}(V_2, V_1) \geq \delta k, \quad (5.45)$$

$$\deg_{G_{\nabla}}^{\min}(V_2, V_3) \geq \delta k, \quad (5.46)$$

$$\deg_{G_{\nabla}}^{\min}(V_3, V_2) \geq \delta k, \quad (5.47)$$

$$\deg_{G_{\mathcal{D}}}^{\max}(V_3, \mathfrak{P}_1 \setminus V_4) < \rho' k, \quad (5.48)$$

$$\deg_{G_{\mathcal{D}}}^{\min}(V_4, V_3) \geq \delta k, \text{ and} \quad (5.49)$$

$$\deg_{G_{\mathcal{D}}}(v, V_3) + \deg_{G_{\mathbf{reg}}}(v, V(\mathcal{N})) \geq h_1 \text{ for each } v \in V_2. \quad (5.50)$$

Definition 5.19 (Configuration $(\diamond 9)$). Suppose that we are in Settings 5.1, and 5.4. We say that the graph G is in Configuration $(\diamond 9)(\delta, \gamma', h_1, h_2, \varepsilon_1, d_1, \mu_1, \varepsilon_2, d_2, \mu_2)$ if the following conditions are met.

The sets V_0, V_1 together with the $(\mathcal{M}_A \cup \mathcal{M}_B)$ -cover \mathcal{F}' witness Preconfiguration $(\heartsuit 1)(\gamma', h_2)$. There exists an $(\varepsilon_1, d_1, \mu_1 k)$ -semiregular matching \mathcal{N} absorbed by $\mathcal{M}_A \cup \mathcal{M}_B$, $V(\mathcal{N}) \subseteq \mathfrak{P}_1$. Further, there is a family $\{(Q_0^{(j)}, Q_1^{(j)})\}_{j \in \mathcal{Y}}$ as in Preconfiguration $(\mathbf{reg})(\varepsilon_2, d_2, \mu_2)$. There is a set $V_2 \subseteq V(\mathcal{N}) \setminus \bigcup \mathcal{F}' \subseteq \bigcup \mathbf{V}$ with the following properties:

$$\deg_{G_{\mathcal{D}}}^{\min}(V_1, V_2) \geq h_1, \quad (5.51)$$

$$\deg_{G_{\mathcal{D}}}^{\min}(V_2, V_1) \geq \delta k. \quad (5.52)$$

Our last configuration, Configuration $(\diamond 10)$, will lead to an embedding very similar to the one in the dense case (as treated in [PS12]; this will be explained in detail in [HKP⁺d]). In order to be able to formalize the configuration we need a preliminary definition. We shall generalize the standard concept of a regularity graph (in the context of regular partitions and Szemerédi's regularity lemma) to graphs with clusters whose sizes are only bounded from below.

Definition 5.20 ($(\varepsilon, d, \ell_1, \ell_2)$ -regularized graph). Let G be a graph, and let \mathcal{V} be an ℓ_1 -ensemble that partitions $V(G)$. Suppose that $G[X]$ is empty for each $X \in \mathcal{V}$ and suppose $G[X, Y]$ is ε -regular and of density either 0 or at least d for each $X, Y \in \mathcal{V}$. Further suppose that for all $X \in \mathcal{V}$ it holds that $|\bigcup N_G(X)| \leq \ell_2$. Then we say that (G, \mathcal{V}) is an $(\varepsilon, d, \ell_1, \ell_2)$ -regularized graph.

A semiregular matching \mathcal{M} of G is consistent with (G, \mathcal{V}) if $\mathcal{V}(\mathcal{M}) \subseteq \mathcal{V}$.

Definition 5.21 (Configuration $(\diamond 10)(\tilde{\varepsilon}, d', \ell_1, \ell_2, \eta')$). Assume Setting 5.1. The graph G contains an $(\tilde{\varepsilon}, d', \ell_1, \ell_2)$ -regularized graph (\tilde{G}, \mathcal{V}) and there is a $(\tilde{\varepsilon}, d', \ell_1)$ -semiregular matching \mathcal{M} consistent with (\tilde{G}, \mathcal{V}) . There are a family $\mathcal{L}^* \subseteq \mathcal{V}$ and distinct clusters $A, B \in \mathcal{V}$ with

(a) $E(\tilde{G}[A, B]) \neq \emptyset$,

(b) $\deg_{\tilde{G}}(v, V(\mathcal{M}) \cup \bigcup \mathcal{L}^*) \geq (1 + \eta')k$ for all but at most $\tilde{\varepsilon}|A|$ vertices $v \in A$ and for all but at most $\tilde{\varepsilon}|B|$ vertices $v \in B$, and

(c) for each $X \in \mathcal{L}^*$ we have $\deg_{\tilde{G}}(v) \geq (1 + \eta')k$ for all but at most $\tilde{\varepsilon}|X|$ vertices $v \in X$.

6 Embedding trees

In this section we provide an embedding of a tree $T_{T1.2} \in \mathbf{trees}(k)$ in the setting of the configurations introduced in Subsection 5.2. In Section 6.1 we first give a fairly detailed overview of the embedding techniques used. In Section 6.3 we introduce a class of stochastic processes which will be used for some embeddings. Section 6.4 contains a number of lemmas about embedding small trees, and use them for embedding knags and shrubs of a given fine partition of $T_{T1.2}$. Embedding the entire tree $T_{T1.2}$ is then handled in the final Section 6.5. There we have to distinguish between particular configurations. The configurations are grouped into three categories (Section 6.5.1, Section 6.5.2, and Section 6.5.3) corresponding to the similarities between the configurations.

6.1 Overview of the embedding procedures

Recall that we are working under Setting 5.1. Given a host graph $G_{T1.2}$ with one of the Configurations $(\diamond 2)$ – $(\diamond 10)$, we have to embed in it a given tree $T = T_{T1.2} \in \mathbf{trees}(k)$, which comes with its (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$. The τk -fine partition of T will make it possible to combine embeddings of smaller parts of T into one embedding of the whole tree. This means that we will first develop tools for embedding singular shrubs and knags of the (τk) -fine partition into various basic building bricks of the configurations: the avoiding set \mathfrak{A} , the expander G_{exp} , regular pairs, and vertices of huge degree Ψ . Second, we will combine these basic techniques to embed the entire tree T . Here, the order in which different parts of T are embedded is important. Also, it will be crucial at some points to reserve places for parts of the tree which will be embedded only later.

In the following subsections, we draft our embedding techniques. We group them into five categories comprising of related configurations³: Configurations $(\diamond 2)$ – $(\diamond 5)$, Configurations $(\diamond 6)$ – $(\diamond 7)$, Configuration $(\diamond 8)$, Configuration $(\diamond 9)$, and Configuration $(\diamond 10)$, treated in Sections 6.1.1, 6.1.2, 6.1.3, 6.1.4, 6.1.5, respectively.

6.1.1 Embedding overview for Configurations $(\diamond 2)$ – $(\diamond 5)$

In each of the Configurations $(\diamond 2)$ – $(\diamond 5)$ we have sets Ψ', Ψ'', L'', L' and V_1 . Further, we have some additional sets (V_2 and/or \mathfrak{A}') depending on the particular configuration.

A common embedding scheme for Configurations $(\diamond 2)$ – $(\diamond 5)$ is illustrated in Figure 6.1. There are two stages of the embedding procedure: the knags, the shrubs \mathcal{S}_A and some parts of the shrubs \mathcal{S}_B are embedded in Stage 1, and then in Stage 2 the remainders of \mathcal{S}_B are embedded. Recall that \mathcal{S}_A contains both internal and end shrubs while \mathcal{S}_B contains exclusively end shrubs. We note that here the shrubs \mathcal{S}_B are further subdivided and some parts of them are embedded in the Stage 1 and some in Stage 2.

- In Stage 1, the knags of T are embedded in Ψ'' and V_1 so that W_A is mapped to Ψ'' and W_B is mapped to V_1 .
- In Stage 1, the internal and end shrubs of \mathcal{S}_A are embedded using the sets V_1, V_2 and \mathfrak{A}' which are specific to the particular Configurations $(\diamond 2)$ – $(\diamond 5)$. The vertices of \mathcal{S}_A neighbouring W_A are always embedded in V_1 . Parts of the shrubs \mathcal{S}_B are embedded while the ancestors of the unembedded remainders are embedded on vertices which have large degrees in Ψ' .

³Configuration $(\diamond 1)$ is trivial (see Section 6.5.1) and needs no draft.




Figure 6.1: An overview of embedding of a tree $T \in \mathbf{trees}(k)$ given with its fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ using Configurations $(\diamond 2)$ – $(\diamond 5)$. The knags are embedded between Ψ'' and V_1 , all the shrubs \mathcal{S}_A are embedded into sets specific to particular configurations so that the vertices neighbouring W_A are embedded in V_1 . Parts of the shrubs \mathcal{S}_B are embedded directly (using various embedding techniques), while the rest is “suspended”, i.e., the ancestors of the unembedded remainders are embedded on vertices which have large degrees in Ψ' . The embedding of \mathcal{S}_B is then finalized in the last stage.

- In Stage 2, the embedding of \mathcal{S}_B is finalized. The remainders of \mathcal{S}_B are embedded starting with embedding their roots in Ψ' .

A hierarchy of the embedding lemmas used to resolve Configurations $(\diamond 2)$ – $(\diamond 5)$ is given in Table 6.1.

6.1.2 Embedding overview for Configurations $(\diamond 6)$ – $(\diamond 7)$

Suppose Setting 5.1 and 5.4 (see Remark 6.1 below for a comment on the constants $\mathfrak{p}_0, \mathfrak{p}_1, \mathfrak{p}_2$). Recall that we have in each of these configurations sets $V_0 \cup V_1 \subseteq \mathfrak{P}_0$, sets $V_2 \cup V_3 \subseteq \mathfrak{P}_1$ and $V_{\text{good}}^{\text{I2}}$.

A common embedding scheme for Configurations $(\diamond 6)$ – $(\diamond 7)$ is illustrated in Figure 6.2. The embedding has three parts.

- The knags of T are embedded between V_0 and V_1 so that W_A is mapped to V_1 and W_B is mapped to V_0 using either the Preconfiguration **(exp)** or **(reg)**. Thus $W_A \cup W_B$ are mapped to $\subseteq \mathfrak{P}_0$.
- The internal shrubs of T are embedded in $V_2 \cup V_3$, always putting neighbours of W_A into V_2 . Note that the internal shrubs are therefore embedded in \mathfrak{P}_1 , and thus there is no interference with embedding the knags. We need to understand why a mere degree of δk (from V_1 to

6.1 Overview of the embedding procedures

Main embedding lemma: Lemma 6.19		
↑	↑	↑
Shrubs \mathcal{S}_A ($\diamond 2$): Lemma 6.5 ($\diamond 3$): Lemma 6.14 ($\diamond 4$): Lemma 6.15 ($\diamond 5$): regularity	Shrubs \mathcal{S}_B (Stage 1): Lemma 6.18	Shrubs \mathcal{S}_B (Stage 2): Lemma 6.17

Table 6.1: Embedding lemmas employed for Configurations ($\diamond 2$)–($\diamond 5$).




Figure 6.2: An overview of embedding a fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a tree $T \in \mathbf{trees}(k)$ using Configurations ($\diamond 6$)–($\diamond 7$). The knags are embedded between V_0 and V_1 , the internal shrubs are embedded in $V_2 \cup V_3$, and the end shrubs are embedded using $V_{\text{good}}^{|2}$.

V_2 , ensured by (5.36) and (5.40), with $\delta \ll 1$) is sufficient for embedding internal shrubs of potentially big total order, that is, how to ensure that already embedded internal trees do not cause a blockage later. Here the expansion⁴ ruling between the V_2 and V_3 comes into play. This property (together with other properties of Preconfigurations **(exp)** and **(reg)**) will allow that, once finished embedding an internal tree, the follow-up snag can be embedded in a place (in V_1) which sees very little of the previously embedded internal shrubs.

This is the only part of the embedding process which makes use of the specifics of Configurations ($\diamond 6$) and ($\diamond 7$). For this reason we will be able to follow the same embedding scheme as presented here also for Configuration ($\diamond 8$), the only difference being the embedding of the internal shrubs (see Section 6.1.3).

- The end shrubs are embedded in the yet unoccupied part of G . For this we use the properties of Preconfigurations ($\heartsuit 1$) or ($\heartsuit 2$). The end shrubs are embedded using (but not entirely into) the designated vertex set $V_{\text{good}}^{|2}$.

⁴This expansion is given by the presence of G_{exp} in Configurations ($\diamond 6$) (cf. (5.38)–(5.39)), and by the presence of the avoiding set \mathfrak{A} in Configurations ($\diamond 7$) ($V_2 \subseteq \mathfrak{A}^{|1} \setminus \bar{V}$).




Table 6.2: Embedding lemmas employed for Configurations ($\diamond 6$)–($\diamond 8$) when embedding a tree $T \in \mathbf{trees}(k)$ with a given fine partition.

The above embedding scheme is divided in two main steps: first the knags and the internal trees are embedded (see Lemma 6.20), and this partial embedding is then extended to end shrubs (see Lemmas 6.22 and 6.23). A more detailed hierarchy of the embedding lemmas which are used is given in Table 6.2.

Remark 6.1. *In Configuration ($\diamond 6$), the number \mathfrak{p}_1 will be approximately the proportion of the total order of the internal shrubs of a given fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of T while \mathfrak{p}_2 will be approximately the proportion of the total order of the end shrubs. The number \mathfrak{p}_0 is just a small constant.*

These numbers – scaled up by k – determine the parameter $h_1 \approx \mathfrak{p}_1 k$ (in Configurations ($\diamond 8$) and ($\diamond 9$)) and $h_2 \approx \mathfrak{p}_2 k$ (in Configurations ($\diamond 6$)–($\diamond 9$)). The properties of these configurations will then allow to embed all the internal shrubs and end shrubs. Note that the parameter h_1 does not appear in Configurations ($\diamond 6$) and ($\diamond 7$). This suggests that the total order of the internal shrubs is not at all important in Configurations ($\diamond 6$)–($\diamond 7$). Indeed, we would succeed even embedding a tree with internal shrubs of total order say $100k$.⁵

In view of this it might be tempting to think that the end shrubs in \mathcal{S}_A could also be embedded using the same technique as the internal shrubs into the sets $V_2 \cup V_3$ provided by these configurations (cf. Figure 6.2). This is however not the case. Indeed, the minimum degree conditions (5.36), (5.40), and (5.44) allow embedding only a small number of shrubs from a single cut-vertex $x \in W_A$ while there may be many end shrubs attached to x ; cf. Remark 3.6(ii).

6.1.3 Embedding overview for Configuration ($\diamond 8$)

Suppose Setting 5.1 and 5.4. We are working with sets $V_0, V_1, V_{\text{good}}^{\uparrow 2}, V_2, V_3$ and V_4 and with semiregular matching \mathcal{N} coming from the configuration.

The embedding scheme follows Table 6.2, and is illustrated in Figure 6.3. Embedding of the knags and of the external shrubs is done in the same way as in Configurations ($\diamond 6$)–($\diamond 7$). We only describe here the way the internal shrubs are embedded. Their roots are embedded in V_2 . From that point we proceed embedding subshrub by subshrub. Some of the subshrubs get embedded between V_3 and V_4 . This pair of sets has the same expansion property as the pair V_2, V_3 in Configuration ($\diamond 7$).

⁵Configuration ($\diamond 8$) has this property only in part. We would succeed even embedding a tree with principal subshrubs of total order say $100k$ provided that the total order of peripheral subshrubs is somewhat smaller than h_1 .




Figure 6.3: An overview of embedding a fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a tree $T \in \mathbf{trees}(k)$ using Configuration $(\diamond 8)$. The knags are embedded between V_0 and V_1 . The roots of the internal shrubs are embedded in V_2 . Some of the subshrubs of the internal shrubs are embedded in $V_3 \cup V_4$ and some in \mathcal{N} ; principal subshrubs are always embedded in $V_3 \cup V_4$. The end shrubs are embedded in using $V_{\text{good}}^{|2}$.

In particular, it allows to avoid the shadow of the already occupied set so that the follow-up snag can be embedded in location almost isolated from the previous images, similarly as described in Section 6.1.2. For this reason we make sure that principal subshrubs get embedded here. The degree condition from V_2 to V_3 is too weak to ensure that all remaining subshrubs are embedded between V_3 and V_4 . Therefore we might have to embed some subshrubs in \mathcal{N} . Condition (5.50) — where h_1 is approximately the order of the internal shrubs, as in Remark 6.1 — indicates that it should be possible to accommodate all the subshrubs. For technical reasons, the order in which different types of subshrubs are embedded is very important.

6.1.4 Embedding overview for Configuration $(\diamond 9)$

The embedding process in Configuration $(\diamond 9)$ follows the same scheme as in Configurations $(\diamond 6)$ – $(\diamond 8)$, but the embedding of the internal shrubs follows the regularity method. Assuming the simplest situation $\mathcal{F} = \mathcal{V}_2(\mathcal{N})$ and $V_2 = V_1(\mathcal{N})$, we would have $\deg^{\min}_{G_{\text{reg}}}(V_1, V_1(\mathcal{N})) \geq h_1$ (cf. (5.51)). See Figure 6.4 for an illustration. Similarly as above, the knags are embedded between V_0 and V_1 . The internal shrubs are accommodated using the regularity method in \mathcal{N} , and the end shrubs are embedded in $V_{\text{good}}^{|2}$ using Preconfiguration $(\heartsuit 1)$. The embedding lemma for this configuration is given in Lemma 6.25.




Figure 6.4: An overview of embedding a fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of a tree $T \in \mathbf{trees}(k)$ using Configuration $(\diamond 9)$. The knags are embedded between V_0 and V_1 , the internal shrubs using the regularity method in \mathcal{N} and the end shrubs are embedded using $V_{\text{good}}^{|2}$.

6.1.5 Embedding overview for Configuration $(\diamond 10)$

Configuration $(\diamond 10)$ is very closely related to the structure obtained by Piguet and Stein [PS12] in their solution of the dense approximate case of Conjecture 1.1.

Theorem 6.2 (Piguet–Stein [PS12]). *For any $q > 0$ and $\alpha > 0$ there exists a number n_0 such that for any $n > n_0$ and $k > qn$ the following holds. For each n -vertex graph G with at least $n/2$ vertices of degree at least $(1 + \alpha)k$ we have $\mathbf{trees}(k + 1) \subseteq G$.*

Let us describe their proof first. Piguet and Stein prove that when $k > qn$ (for some fixed $q > 0$ and k sufficiently large) the cluster graph⁶ \mathbf{G}_{reg} of a graph $G \in \mathbf{LKS}(n, k, \eta)$ contains the following structure (cf. [PS12, Lemma 8]). There is a set of clusters $\mathbf{L} \subseteq \mathbf{V}$ such that each cluster in \mathbf{L} contains only vertices of captured degrees at least $(1 + \frac{\eta}{2})k$. There is a matching $M \subseteq \mathbf{G}_{\text{reg}}$, and an edge AB , with $A, B \in \mathbf{L}$. One of the following conditions is satisfied

- (H1) M covers $N_{\mathbf{G}_{\text{reg}}}(\{A, B\})$, or
- (H2) M covers $N_{\mathbf{G}_{\text{reg}}}(A)$, and the vertices in B have captured degrees at least $(1 + \frac{\eta}{2})\frac{k}{2}$ into $\bigcup(\mathbf{L} \cup V(M))$. Further, each edge in M has at most one endvertex in $N_{\mathbf{G}_{\text{reg}}}(A)$.

Piguet and Stein use structures (H1) and (H2) to embed any given tree $T \in \mathbf{trees}(k)$ into G using the regularity method; see Sections 3.6 and 3.7 in [PS12], respectively. Actually, a slight relaxation of (H1) and (H2) would be sufficient for the embedding to work, as can be easily seen from their proof: Again, there is a set of clusters $\mathbf{L} \subseteq \mathbf{V}$ such that each cluster in \mathbf{L} contains only vertices of captured degrees at least $(1 + \frac{\eta}{2})k$, there is a matching $M \subseteq \mathbf{G}_{\text{reg}}$, and an edge AB , $A, B \in \mathbf{L}$. One of the following conditions is satisfied

- (H1') the vertices in $A \cup B$ have captured degrees at least $(1 + \frac{\eta}{2})k$ into the vertices of $\bigcup(\mathbf{L} \cup V(M))$,
or

⁶ordinary, in the sense of the classic regularity lemma

(H2') the vertices in A have captured degrees at least $(1 + \frac{\eta}{2})k$ into the vertices of $\bigcup V(M)$, and the vertices in B have captured degrees at least $(1 + \frac{\eta}{2})\frac{k}{2}$ into $\bigcup(\mathbf{L} \cup V(M))$. Further, each edge in M has at most one endvertex in $\mathbf{N}_{\mathbf{G}_{\text{reg}}}(A)$.

It can be seen that Configuration $(\diamond 10)$ is a direct counterpart to **(H1')**.⁷ (The counterpart of **(H2')** is contained in Configuration $(\diamond 9)$ and the similarity is somewhat weaker.)

The embedding lemma for Configuration $(\diamond 10)$ is stated in Lemma 6.26.

6.2 The role of random splitting

The random splitting as introduced in Setting 5.4 is used in Configurations $(\diamond 6)$ – $(\diamond 9)$; the set \mathfrak{P}_0 will host the cut-vertices $W_A \cup W_B$, the set \mathfrak{P}_1 will host the internal shrubs, and the set \mathfrak{P}_2 will (essentially) host the end shrubs of a (τk) -fine partition of $T_{T1.2}$.

The need for introducing the random splitting is dictated by Configurations $(\diamond 6)$ – $(\diamond 9)$. To see this, let us try to follow the embedding plan from, for example, Section 6.1.2 without the random splitting, i.e., dropping the conditions $\subseteq \mathfrak{P}_0, \subseteq \mathfrak{P}_1, \subseteq \mathfrak{P}_2$ from Definitions 5.12–5.17. Then the sets V_2 and V_3 in Figure 6.2, which will host the internal shrubs, may interfere with V_0 and V_1 primarily designated for W_A and W_B . In particular, the conditions on degrees between V_0 and V_1 given by (5.33)–(5.34) in Definition 5.14, or given by the super-regularity in Definition 5.15 (in which $\beta_{D5.14} > 0$, or $d'_{D5.15}\mu_{D5.15} > 0$ are tiny) need not be sufficient for embedding greedily all the cut-vertices and all the internal shrubs of $T_{T1.2}$. It should be noted that this problem occurs even in Preconfiguration **(exp)**, i.e., the expanding property does not add enough strength to the minimum degree conditions.⁸ Restricting V_0 and V_1 to host only the cut-vertices (only $O(1/\tau) = o(k)$ of them in total, cf. Definition 3.3(c)), resolves the problem.

The above justifies the distinction between the space \mathfrak{P}_0 for embedding the cut-vertices and the space $\mathfrak{P}_1 \cup \mathfrak{P}_2$ for embedding the shrubs. There are some other approaches which do not need to further split $\mathfrak{P}_1 \cup \mathfrak{P}_2$ but doing so seems to be the most convenient.

6.3 Stochastic process Duplicate(ℓ)

Let us introduce a class of stochastic processes, which we call Duplicate(ℓ) ($\ell \in \mathbb{N}$). These are discrete processes $(X_1, Y_1), (X_2, Y_2), \dots, (X_q, Y_q) \in \{0, 1\}^2$ (where $q \in \mathbb{N}$ is arbitrary) satisfying the following.

- For each $i \in [q]$, we have either
 - (a) $X_i = Y_i = 0$ (deterministically), or
 - (b) $X_i = Y_i = 1$ (deterministically), or
 - (c) exactly one of X_i and Y_i is one, and in that case $\mathbf{P}[X_i = 1] = \frac{1}{2}$.
- If the distribution of (X_i, Y_i) is according to (c), then the random choice is made independently of the values (X_j, Y_j) ($j < i$).
- We have $\sum_{i=1}^q (X_i + Y_i) \leq \ell$.

⁷Observe that some parts of \mathbf{G}_{reg} are irrelevant in the embedding process of [PS12]. The objects \mathbf{G}_{reg} , \mathbf{L} , and M in the structural result of [PS12] correspond to (\tilde{G}, \mathcal{V}) , \mathcal{L}^* , and \mathcal{M} in Configuration $(\diamond 10)$.

⁸See [HKP⁺a, Section 3.6] for details.

6.4 Embedding small trees

Needless to say that this definition is not deep and its purpose is only to adopt the language we shall be using later. The following lemma asserts that the first and second component of a process $\text{Duplicate}(\ell)$ are typically balanced.

Lemma 6.3. *Suppose that $(X_1, Y_1), (X_2, Y_2), \dots, (X_q, Y_q)$ is a process in $\text{Duplicate}(\ell)$. Then for any $a > 0$ we have*

$$\mathbf{P} \left[\sum_{i=1}^q X_i - \sum_{i=1}^q Y_i \geq a \right] \leq \exp \left(-\frac{a^2}{2\ell} \right).$$

Proof. We shall be using the following version of the Chernoff bound for sums of independent random variables Z_i , with distribution $\mathbf{P}[Z_i = 1] = \mathbf{P}[Z_i = -1] = \frac{1}{2}$.

$$\mathbf{P} \left[\sum_{i=1}^n Z_i \geq a \right] \leq \exp \left(-\frac{a^2}{2n} \right). \quad (6.1)$$

Let $J \subseteq [q]$ be the set of all indices i with $X_i + Y_i = 1$. By the definition of $\text{Duplicate}(\ell)$, we have $|J| \leq \ell$. By (6.1) we have

$$\mathbf{P} \left[\sum_J (X_i - Y_i) \geq a \right] \leq \exp \left(-\frac{a^2}{2|J|} \right) \leq \exp \left(-\frac{a^2}{2\ell} \right).$$

□

We shall use the stochastic process Duplicate to guarantee that certain fixed vertex sets do not get overfilled during our tree embedding procedure. Duplicate is used in Lemmas 6.12 and 6.13 through Lemma 6.11.

6.4 Embedding small trees

When embedding the tree $T_{T1.2}$ in our proof of Theorem 1.2 it will be important to control where different bits of $T_{T1.2}$ go. This motivates the following notation. Let $X_1, \dots, X_\ell \subseteq V(T)$ be arbitrary vertex sets of a tree T , and let $V_1, \dots, V_\ell \subseteq V(G)$ be arbitrary vertex sets of a graph G . Then an embedding $\phi : V(T) \rightarrow V(G)$ of T in G is an $(X_1 \hookrightarrow V_1, \dots, X_\ell \hookrightarrow V_\ell)$ -embedding if $\phi(X_i) \subseteq V_i$ for each $i \in [\ell]$.

We provide several sufficient conditions for embedding a small tree with additional constraints.

The first lemma deals with embedding using an avoiding set.

Lemma 6.4. *Let $\Lambda, k \in \mathbb{N}$ and let $\varepsilon, \gamma \in (0, \frac{1}{2})$ with $\gamma^2 > \varepsilon$. Suppose \mathfrak{A} is a $(\Lambda, \varepsilon, \gamma, k)$ -avoiding set with respect to a set \mathcal{D} of $(\gamma k, \gamma)$ -dense spots in a graph H . Suppose that $(T_1, r_1), \dots, (T_\ell, r_\ell)$ are rooted trees with $|\bigcup_i T_i| \leq \gamma k/2$. Let $U \subseteq V(H)$ with $|U| \leq \Lambda k$, and let $U^* \subseteq \mathfrak{A}$ with $|U^*| \geq \varepsilon k + \ell$. Then there are mutually disjoint $(r_i \hookrightarrow U^*, V(T_i) \setminus \{r_i\} \hookrightarrow V(H) \setminus U)$ -embeddings of the trees (T_i, r_i) in H .*

Proof. Since \mathfrak{A} is $(\Lambda, \varepsilon, \gamma, k)$ -avoiding, there exists a set $Y \subseteq \mathfrak{A}$ with $|Y| \leq \varepsilon k$, such that each vertex v in $\mathfrak{A} \setminus Y$ has degree at least γk into some $(\gamma k, \gamma)$ -dense spot $D \in \mathcal{D}$ with $|U \cap V(D)| \leq \gamma^2 k$. In particular, $U^* \setminus Y$ is large enough so that we can embed all vertices r_i there. We extend this embedding successively to an embedding of $\bigcup_i T_i$, in each step finding a suitable image in $V(D) \setminus U$ for one neighbour of an already embedded vertex $v \in \bigcup_i V(T_i)$. This is possible since the image of v has degree at least $\gamma k - |U \cap V(D)| > \gamma k/2 \geq \sum_i v(T_i)$ into $V(D) \setminus U$. □

The next lemma deals with embedding a tree into a nowhere-dense graph, a primal example of which is the graph G_{exp} .

Lemma 6.5. *Let $k \in \mathbb{N}$, let $Q \geq 1$ and let $\gamma, \zeta \in (0, 1)$ be such that $128Q\gamma \leq \zeta^2$. Let H be a $(\gamma k, \gamma)$ -nowhere-dense graph. Let $(T_1, r_1), \dots, (T_\ell, r_\ell)$ be rooted trees of total order less than $\zeta k/4$. Let $V_1, V_2, U, U^* \subseteq V(H)$ be four sets with $U^* \subseteq V_1$, $|U| < Qk$, $|U^*| > \frac{32Q^2\gamma}{\zeta}k + \ell$, and $\deg^{\min}_H(V_j, V_{3-j}) \geq \zeta k$ for $j = 1, 2$. Then there are mutually disjoint $(r_i \hookrightarrow U^*, V_{\text{even}}(T_i) \hookrightarrow V_1 \setminus U, V_{\text{odd}}(T_i) \hookrightarrow V_2 \setminus U)$ -embeddings of the trees (T_i, r_i) in H .*

Proof. Set $B := \text{shadow}_H(U, \zeta k/2)$. By Fact 4.13, we have $|B| \leq \frac{32Q^2\gamma}{\zeta}k \leq \frac{\zeta}{4}k$. In particular, $U^* \setminus B$ is large enough to accommodate the images $\phi(r_i)$ of all vertices r_i .

Successively, extend ϕ , in each step mapping a neighbour u of some already embedded vertex $v \in \bigcup_i V(T_i)$ to a yet unused neighbour of $\phi(v)$ in $V_j \setminus (B \cup U)$, where j is either 1 or 2, depending on the parity of $\text{dist}_T(r, v)$. This is possible as $\phi(v)$, lying outside B , has at least $\zeta k/2$ neighbours in $V_i \setminus U$. Thus $\phi(v)$ has at least $\zeta k/4$ neighbours in $V_i \setminus (U \cup B)$, which is more than $\sum_i v(T_i)$. \square

The next three standard lemmas deal with embedding trees in a regular or a super-regular pair. We omit their proofs.

Lemma 6.6. *Let $\varepsilon > 0$ and $\beta > 2\varepsilon$. Let (C, D) be an ε -regular pair in a graph H , with $|C| = |D| =: \ell$, and with density $d(C, D) \geq 3\beta$. Suppose that there are sets $X \subseteq C$, $Y \subseteq D$, and $X^* \subseteq X$ satisfying $\min\{|X|, |Y|\} \geq 4\frac{\varepsilon}{\beta}\ell$ and $|X^*| > \frac{\beta}{2}\ell$. Let (T, r) be a rooted tree of order $v(T) \leq \varepsilon\ell$. Then there exists an $(r \hookrightarrow X^*, V_{\text{even}}(T) \hookrightarrow X, V_{\text{odd}}(T) \hookrightarrow Y)$ -embedding of T in H .*

Lemma 6.7. *Let $\beta, \varepsilon > 0$ and $\ell \in \mathbb{N}$ be such that $\beta > 2\varepsilon$. Let (C, D) be an ε -regular pair with $|C| = |D| = \ell$ of density $d(C, D) \geq 3\beta$ in a graph H . Let $(T_1, r_1), (T_2, r_2), \dots, (T_s, r_s)$ be rooted trees with $v(T_i) \leq \varepsilon\ell$ for all $i \in [s]$. Let $U \subseteq V(H)$ fulfill $|C \cap U| = |D \cap U|$, and let $X^* \subseteq (C \cup D) \setminus U$ be such that*

$$|X^*| \geq \sum_{i=1}^s v(T_i) + 50\beta\ell. \quad (6.2)$$

Then there are mutually disjoint $(r_i \hookrightarrow X^, V(T_i) \hookrightarrow (C \cup D) \setminus U)$ -embeddings of the trees (T_i, r_i) in H .*

Lemma 6.8. *Let $d > 10\varepsilon > 0$. Suppose that (A, B) forms an (ε, d) -super-regular pair with $|A|, |B| \geq \ell$. Let $U_A \subseteq A$, $U_B \subseteq B$ be such that $|U_A| \leq |A|/2$ and $|U_B| \leq d|B|/4$. Let (T, r) be a rooted tree of order at most $d\ell/4$, and let $v \in A \setminus U_A$ be arbitrary. Then there exists an $(r \hookrightarrow v, V_{\text{even}}(T, r) \hookrightarrow A \setminus U_A, V_{\text{odd}}(T, r) \hookrightarrow B \setminus U_B)$ -embedding of T .*

Suppose that we have a rooted tree (T, r) to be embedded, and its root was already on a vertex $\phi(r)$. Suppose that r has degree $\ell_X + \ell_Y$ in a regular pair (X, Y) , where $\ell_X := \deg(\phi(r), X)$, $\ell_Y := \deg(\phi(r), Y)$, with $\ell_X \geq \ell_Y$, say. The hope is that we can embed T in (X, Y) as long as $v(T)$ is a bit smaller than $\ell_X + \ell_Y$. For this, the greedy strategy does not work (see Figure 6.5) and we need to be somewhat more careful. We split the embedding process into two stages. In the first stage we choose a subset of the components of $T - r$ of total order approximately $2 \min(\ell_X, \ell_Y) = 2\ell_Y$. When embedding these, we choose orientations of each component in such a way that the image is approximately balanced with respect to X and Y . In the second stage we embed the remaining components so that their roots are embedded in X . We refer to the first stage as embedding in an *balanced way*, and as embedding in an *unbalanced way* to the second stage.

The next lemma says that each regular pair can be filled-up in a balanced way by trees.




Figure 6.5: An example of a rooted tree (T, r) , depicted on the left. The forest $T - r$ has three components (I), (II), (III) of total order 12. Say the vertex r is embedded so that for the regular pair (X, Y) we have $\deg(\phi(r), X) = 8$, $\deg(\phi(r), Y) = 4$ (neighbourhoods of $\phi(r)$ hatched). While the greedy strategy does not work (middle), splitting the process into a balanced and an unbalanced stage (right) does — here the components (I) and (II) are embedded in the balanced stage and the component (III) in the unbalanced stage.

Lemma 6.9. *Let G be a graph, $v \in V(G)$ be a vertex, \mathcal{M} be an $(\varepsilon, d, \nu k)$ -semiregular matching in G , and $\{f_{CD}\}_{(C,D) \in \mathcal{M}}$ a family of integers between $-\tau k$ and τk . Suppose (T, r) is a rooted tree,*

$$v(T) \leq \left(1 - \frac{4(\varepsilon + \frac{\tau}{\nu})}{d - 2\varepsilon}\right) |V(\mathcal{M})|,$$

with the property that each component of $T - r$ has order at most τk . If $V(\mathcal{M}) \subseteq N_G(v)$ then there exists an $(r \hookrightarrow v, V(T - r) \hookrightarrow V(\mathcal{M}))$ -embedding ϕ of T such that for each $(C, D) \in \mathcal{M}$ we have $|C \cap \phi(T)| + f_{CD} = |D \cap \phi(T)| \pm \tau k$.

The proof of Lemma 6.9 is standard, and is given for example in [HP, Lemma 5.11].

Lemma 6.9 suggests the following definitions. The *discrepancy* of a set X with respect to a pair of sets (C, D) is the number $|C \cap X| - |D \cap X|$. X is *s-balanced* with respect to a semiregular matching \mathcal{M} if the discrepancy of X with respect to each $(C, D) \in \mathcal{M}$ is at most s in absolute value.

Lemma 6.10. *Let G be a graph, $v \in V(G)$ be a vertex, \mathcal{M} be an $(\varepsilon, d, \nu k)$ -semiregular matching in G with an \mathcal{M} -cover \mathcal{F} , and $U \subseteq V(G)$. Suppose (T, r) is a rooted tree with*

$$v(T) + |U| \leq \deg_G\left(v, V(\mathcal{M}) \setminus \bigcup \mathcal{F}\right) - \frac{4(\varepsilon + \frac{\tau}{\nu})}{d - 2\varepsilon} |V(\mathcal{M})|,$$

such that each component of $T - r$ has order at most τk . Then there exists an $(r \hookrightarrow v, V(T - r) \hookrightarrow V(\mathcal{M}) \setminus U)$ -embedding ϕ of T .

The proof of Lemma 6.10 is again standard and we again omit it.

The following lemma uses a probabilistic technique to embed a shrub while reserving a set of vertices in the host graph for later use. We wish the reserved set to use about as much space inside certain given sets P_i as the image of our shrub does. (In later applications the sets P_i correspond to neighbourhoods of vertices which are still ‘active’.)

Lemma 6.11 will find an immediate application in all the remaining lemmas of this subsection. However it is really necessary only for Lemmas 6.12–6.13, which deal with embedding shrubs in the presence of one of the Configurations $(\diamond 6)$ – $(\diamond 8)$. For Lemmas 6.14 and 6.15, which are for Configurations $(\diamond 3)$ and $(\diamond 4)$ a simpler auxiliary lemma (without reservations) would suffice.

Lemma 6.11. *Let H be a graph, let $X^*, X_1, X_2, P_1, P_2, \dots, P_L \subseteq V(H)$, and let $(T_1, r_1), \dots, (T_\ell, r_\ell)$ be rooted trees, such that $L \leq k$, $|P_j| \leq k$ for each $j \in [L]$, and $|X^*| \geq 2\ell$. Suppose that $\deg^{\min}(X_1 \cup X^*, X_2) \geq 2 \sum v(T_i)$ and $\deg^{\min}(X_2, X_1) \geq 2 \sum v(T_i)$.*

Then there exist pairwise disjoint $(r_i \hookrightarrow X^, V_{\text{even}}(T_i, r_i) \setminus \{r_i\} \hookrightarrow X_1, V_{\text{odd}}(T_i, r_i) \hookrightarrow X_2)$ -embeddings ϕ_i of T_i in G and a set $C \subseteq (X_1 \cup X_2) \setminus \bigcup \phi_i(T_i)$ of size $\sum v(T_i)$ such that for each $j \in [L]$ we have*

$$|P_j \cap \bigcup \phi_i(T_i)| \leq |P_j \cap C| + k^{3/4}. \quad (6.3)$$

Proof. Let $m := \sum v(T_i)$.

We construct pairwise disjoint random $(r_i \hookrightarrow X^*, V_{\text{even}}(T_i, r_i) \setminus \{r_i\} \hookrightarrow X_1, V_{\text{odd}}(T_i, r_i) \hookrightarrow X_2)$ -embeddings ϕ_i and a set $C \subseteq V(H) \setminus \bigcup \phi_i(T_i)$ which satisfies (6.3) with positive probability. Then the statement follows.

Enumerate the vertices of $\bigcup T_i$ as $\bigcup V(T_i) = \{v_1, \dots, v_m\}$ such that $v_i = r_i$ for $i = 1, \dots, \ell$, and such that for each $j > \ell$ we have that the parent of v_j lies in the set $\{v_1, \dots, v_{j-1}\}$. Pick pairwise disjoint sets $A_1, \dots, A_\ell \subseteq X^*$ of size two. Uniformly at random denote one element of A_j as x_j and the other as y_j .

Now, successively for $i = \ell + 1, \dots, m$, we shall define vertices x_i and y_i . Let r denote the root of the tree in which v_i lies, and let $v_s = \text{Par}(v_i)$. We shall choose $x_i, y_i \in X_{j_i}$ where $j_i = \text{dist}(r, v_i) \bmod 2 + 1$. In step i , proceed as follows. Since $x_s \in X_{j_s}$ (or since $x_s \in X^*$), we have

$$\deg(x_s, X_{j_i} \setminus \bigcup_{h < i} \{x_h, y_h\}) \geq 2.$$

Hence, we may take an arbitrary subset $A_i \subseteq (N(x_s) \cap X_{j_i}) \setminus \bigcup_{h < i} \{x_h, y_h\}$ of size exactly two. As above, randomly label its elements as x_i and y_i independently of all other choices.

The choices of the maps $(v_j \mapsto x_j)_{j=1}^m$ determine ϕ_1, \dots, ϕ_ℓ . Then $C := \{y_1, \dots, y_m\}$ has size exactly m and avoids $\bigcup \phi_i(T_i)$.

For each $j \in [L]$ we set up a stochastic process $\mathfrak{S}^{(j)} = \left((X_i^{(j)}, Y_i^{(j)}) \right)_{i=1}^m$, defined by $X_i^{(j)} = \mathbf{1}_{\{x_i \in P_j\}}$ and $Y_i^{(j)} = \mathbf{1}_{\{y_i \in P_j\}}$. Note that $\mathfrak{S}^{(j)} \in \text{Duplicate}(|P_j|) \subseteq \text{Duplicate}(k)$. Thus, for a fixed $j \in [L]$, by Lemma 6.3, the probability that $|P_j \cap (\bigcup \phi_i(T_i))| > |P_j \cap C| + k^{3/4}$ is at most $\exp(-\sqrt{k}/2)$. Using the union bound over all $j \in [L]$ we get that Property 6.5 holds with probability at least

$$1 - L \cdot \exp\left(-\frac{\sqrt{k}}{2}\right) > 0.$$

This finishes the proof. \square

We now get to the first application of Lemma 6.11.

Lemma 6.12. *Assume we are in Setting 5.1. Suppose that the sets V_2, V_3 are such that for $j = 2, 3$ we have*

$$\deg^{\min}_H(V_j, V_{5-j}) \geq \delta k, \quad (6.4)$$

where $\delta > 300/k$, and H is a $(\gamma k, \gamma)$ -nowhere dense graph. Suppose that $U, U^, P_1, P_2, \dots, P_L \subseteq V(G)$, and $L \leq k$, are such that $|U| \leq \frac{\delta}{24\sqrt{\gamma}}k$, $U^* \subseteq V_2$, $|U^*| \geq \frac{\delta}{4}k$, and $|P_j| \leq k$ for each $j \in [L]$. Let (T, r) be a rooted tree of order at most $\delta k/8$.*

6.4 Embedding small trees

Then there exists a $(r \hookrightarrow U^*, V_{\text{even}}(T, r) \setminus \{r\} \hookrightarrow V_2 \setminus U, V_{\text{odd}}(T, r) \hookrightarrow V_3 \setminus U)$ -embedding ϕ of T in G and a set $C \subseteq (V_2 \cup V_3) \setminus (U \cup \phi(T))$ of size $v(T)$ such that for each $j \in [L]$ we have

$$|P_j \cap \phi(T)| \leq |P_j \cap C| + k^{3/4}. \quad (6.5)$$

Proof. Set $B := \text{shadow}_{G_{\text{exp}}}(U, \delta k/4)$. By Fact 4.13, we have that $|B| \leq 64 \frac{\gamma}{\delta} (\frac{\delta}{24\sqrt{\gamma}})^2 k \leq \frac{\delta}{4} k - 2$. In particular, $X^* := U^* \setminus B$ has size at least 2. Set $X_1 := V_2 \setminus (U \cup B)$ and set $X_2 := V_3 \setminus (U \cup B)$. Using (6.4), we find that

$$\deg_{G_{\text{exp}}}^{\min}(X_j, X_{3-j}) \geq \delta k - \deg_{G_{\text{exp}}}^{\max}(X_j, U) - |B| \geq \delta k - \frac{\delta}{4} k - \frac{\delta}{4} k \geq 2v(T)$$

for $j = 1, 2$. We may thus apply Lemma 6.11 to obtain the desired embedding ϕ and the set C . \square

Lemma 6.13. *Assume Setting 5.1 and Setting 5.4. Suppose that we are given sets $Y_1, Y_2 \subseteq \mathfrak{P}_1 \setminus \bar{V}$ with $Y_1 \subseteq \mathfrak{A}$, and*

$$(i) \deg_{G_{\mathcal{D}}}^{\max}(Y_1, \mathfrak{P}_1 \setminus Y_2) \leq \frac{\eta\gamma}{400}, \text{ and}$$

$$(ii) \deg_{G_{\mathcal{D}}}^{\min}(Y_2, Y_1) \geq \delta k.$$

Suppose that $U, U^*, P_1, P_2, \dots, P_L \subseteq V(G)$ are sets such that $|U| \leq \frac{\Lambda\delta}{2\Omega^*} k$, $U^* \subseteq Y_1$, with $|U^*| \geq \frac{\delta}{4} k$, $|P_j| \leq k$ for each $j \in [L]$, and $L \leq k$. Suppose $(T_1, r_1), \dots, (T_\ell, r_\ell)$ are rooted trees of total order at most $\delta k/1000$. Suppose further that $\delta < \eta\gamma/100$, $\varepsilon' < \delta/1000$, and $k > 1000/\delta$.

Then there exist pairwise disjoint $(r_i \hookrightarrow U^*, V_{\text{even}}(T_i, r_i) \hookrightarrow Y_1 \setminus U, V_{\text{odd}}(T_i, r_i) \hookrightarrow Y_2 \setminus U)$ -embeddings ϕ_i of T_i in G and a set $C \subseteq V(G - \bigcup \phi_i(T_i))$ of size $\sum v(T_i)$ such that for each $j \in [L]$ we have that

$$|P_j \cap \bigcup \phi_i(T_i)| \leq |P_j \cap C| + k^{3/4}. \quad (6.6)$$

Proof. Set $U' := \text{shadow}_{G_{\mathcal{D}}}(U, \delta k/2) \cup U$. By Fact 4.12, we have $|U'| \leq \Lambda k$. As Y_1 is a $(\Lambda, \varepsilon', \gamma, k)$ -avoiding set, by Definition 4.5 there exists a set $B \subseteq Y_1$, $|B| \leq \varepsilon' k$ such that for all $v \in Y_1 \setminus B$ there exists a dense spot $D_v \in \mathcal{D}$ with $v \in V(D_v)$ and $|V(D_v) \cap U'| \leq \gamma^2 k$. As Y_1 is disjoint from \bar{V} , by Definition 5.3(4) and by (5.12), we have that $\deg_{D_v}(v, V(D_v)^{\uparrow 1}) \geq \frac{\eta\gamma}{200} k$. By (i), we have that $\deg_{G_{\mathcal{D}}}(v, V(D_v)^{\uparrow 1} \setminus Y_2) < \frac{\eta\gamma}{400} k$, and hence,

$$\deg_{G_{\mathcal{D}}}(v, (V(D_v)^{\uparrow 1} \cap Y_2) \setminus U') \geq \frac{\eta\gamma k}{400} - \gamma^2 k \geq \frac{\eta\gamma k}{800}.$$

Thus,

$$\deg_{G_{\mathcal{D}}}^{\min}(Y_1 \setminus B, Y_2 \setminus (U' \cup B)) \geq \frac{\eta\gamma k}{800} - \varepsilon' k \geq 2 \sum v(T_i). \quad (6.7)$$

Further, by the definition of U' and by (ii), we have

$$\deg_{G_{\mathcal{D}}}^{\min}(Y_2 \setminus U', Y_1 \setminus (U \cup B)) \geq \frac{\delta k}{2} - \varepsilon' k \geq 2 \sum v(T_i). \quad (6.8)$$

Set $X^* := U^* \setminus B$, and note that $|X^*| \geq \delta k/4 - \varepsilon' k \geq 2\ell$. Set $X_1 := Y_1 \setminus (U \cup B)$ and $X_2 := Y_2 \setminus (U' \cup B)$. Inequalities (6.7) and (6.8) guarantee that we may apply Lemma 6.11 to obtain the desired embeddings ϕ_i . \square

6.4 Embedding small trees

Lemma 6.14. *Assume Setting 5.1. Suppose that the sets $L', L'', \Psi', \Psi'', V_1, V_2$ witness Configuration $(\diamond 3)(0, 0, \gamma/4, \delta)$. Suppose that $U, U^* \subseteq V(G)$ are sets such that $|U| \leq k$, $U^* \subseteq V_1$, $|U^*| \geq \frac{\delta}{4}k$. Suppose (T, r) is a rooted tree of order at most $\delta k/1000$. Suppose further that $\delta \leq \gamma/100$, $\varepsilon' < \delta/1000$, and $4\Omega^*/\delta \leq \Lambda$.*

Then there is an $(r \hookrightarrow U^, V_{\text{even}}(T, r) \setminus \{r\} \hookrightarrow V_1 \setminus U, V_{\text{odd}}(T, r) \hookrightarrow V_2 \setminus U)$ -embedding of T in G .*

Proof. The proof of this lemma is very similar to the one of Lemma 6.13 (in fact, even easier). Set $U' := \text{shadow}_{G_{\mathcal{D}}}(U, \delta k/2) \cup U$ and note that $|U'| \leq \Lambda k$ by Fact 4.12. As V_1 is $(\Lambda, \varepsilon', \gamma, k)$ -avoiding, by Definition 4.5 there is a set $B \subseteq V_1$, $|B| \leq \varepsilon' k$ such that for all $v \in V_1 \setminus B$ there exists a dense spot $D_v \in \mathcal{D}$ with $\deg_{D_v}(v, V(D_v) \setminus U') \geq \gamma k/2$. By (5.19), we know that $\deg_{G_{\mathcal{D}}}(v, V(D_v) \setminus V_2) \leq \gamma k/4$, and hence, $\deg_{G_{\mathcal{D}}}(v, (V(D_v) \cap V_2) \setminus U') \geq \gamma k/4$. Thus,

$$\deg^{\min}_{G_{\mathcal{D}}}(V_1 \setminus B, V_2 \setminus U') \geq \frac{\gamma k}{4} \geq 2v(T). \quad (6.9)$$

Further, by the definition of U' and by (5.20), we have

$$\deg^{\min}_{G_{\mathcal{D}}}(V_2 \setminus U', V_1 \setminus U) \geq \frac{\delta k}{2} \geq 2(T). \quad (6.10)$$

Set $X^* := U^* \setminus B$, and note that $|X^*| \geq \delta k/4 - \varepsilon' k \geq 2$. Set $X_1 := V_1 \setminus (U \cup B)$ and $X_2 := V_2 \setminus (U' \cup B)$. Inequalities (6.9) and (6.10) guarantee that we may apply Lemma 6.11 (with empty sets P_i) to obtain the desired embedding ϕ . \square

Lemma 6.15. *Assume Setting 5.1. Suppose that the sets $L', L'', \Psi', \Psi'', V_1, \mathfrak{A}', V_2$ witness Configuration $(\diamond 4)(0, 0, \gamma/4, \delta)$. Suppose that $U \subseteq V(G)$, $U^* \subseteq V_1$ are sets such that $|U| \leq k$ and $|U^*| \geq \frac{\delta}{4}k$. Suppose (T, r) is a rooted tree of order at most $\delta k/20$ with a fruit r' . Suppose further that $4\varepsilon' \leq \delta \leq \gamma/100$, and $\Lambda \geq 300(\frac{\Omega^*}{\delta})^3$.*

Then there exists an $(r \hookrightarrow U^, r' \hookrightarrow V_1 \setminus U, V(T) \setminus \{r, r'\} \hookrightarrow (\mathfrak{A}' \cup V_2) \setminus U)$ -embedding of T in G .*

Proof. Set

$$U' := \tilde{U} \cup \text{shadow}_{G_{\nabla-\Psi}}(U, \delta k/4) \cup \text{shadow}_{G_{\nabla-\Psi}}^{(2)}(\tilde{U}, \delta k/4)$$

and let

$$U'' := \tilde{U} \cup \text{shadow}_{G_{\mathcal{D}}}(U', \delta k/2).$$

We use Fact 4.12 to see that $|U'| \leq \frac{\delta}{4\Omega^*}\Lambda k$ and $|U''| \leq \Lambda k$. We then use Definition 4.5 and (5.24) to find a set $B \subseteq \mathfrak{A}'$ of size at most $\varepsilon' k$ such that

$$\deg^{\min}_{G_{\mathcal{D}}}(\mathfrak{A}' \setminus B, V_2 \setminus U'') \geq 2v(T). \quad (6.11)$$

Using (6.11), and employing (5.21) and (5.23), we see that we may apply Lemma 6.11 with $X_{L6.11}^* := U^*$, $X_{1,L6.11} := \mathfrak{A}' \setminus (B \cup U')$ and $X_{2,L6.11} := V_2 \setminus U''$ (and with empty sets P_i) in order to embed the tree $T - T(r, \uparrow r')$ rooted at r . Then embed $T(r, \uparrow r')$, by applying Lemma 6.11 a second time, using (5.21) and (5.22). \square

6.5 Main embedding lemmas

For this section, we need to introduce the notion of a ghost. Given a semiregular matching \mathcal{N} , we call an involution $\mathfrak{d} : V(\mathcal{N}) \rightarrow V(\mathcal{N})$ with the property that $\mathfrak{d}(S) = T$ for each $(S, T) \in \mathcal{N}$ a *matching involution*.

Assume Setting 5.1 and fix a matching involution \mathfrak{b} for $\mathcal{M}_A \cup \mathcal{M}_B$. For any set $U \subseteq V(G)$ we then define by

$$\text{ghost}(U) := U \cup \mathfrak{b}(U \cap V(\mathcal{M}_A \cup \mathcal{M}_B)) .$$

Clearly, we have that $|\text{ghost}(U)| \leq 2|U|$, and $|\text{ghost}(U) \cap S| = |\text{ghost}(U) \cap T|$ for each $(S, T) \in \mathcal{M}_A \cup \mathcal{M}_B$.

The notion of ghost extends to other semiregular matchings. If \mathcal{N} is a semiregular matching and \mathfrak{d} a matching involution for \mathcal{N} then we write $\text{ghost}_{\mathfrak{d}}(U) := U \cup \mathfrak{d}(U \cap V(\mathcal{N}))$.

6.5.1 Embedding in Configuration ($\diamond 1$)

This subsection contains an easy observation that $\mathbf{trees}(k) \subseteq G$ in case G contains Configuration ($\diamond 1$).

Lemma 6.16. *Let G be a graph, and let $A, B \subseteq V(G)$ be such that $\deg^{\min}(G[A, B]) \geq k/2$, and $\deg^{\min}(A) \geq k$. Then $\mathbf{trees}(k) \subseteq G$.*

Proof. Let $T \in \mathbf{trees}(k)$ have colour classes X and Y , with $|X| \geq k/2 \geq |Y|$. By Fact 3.2, for the set W of those leaves of T that lie in X , we have $|X \setminus W| \leq k/2$. We embed $T - W$ greedily in G , mapping Y to A and $X \setminus W$ to B . We then embed W using the fact that $\deg^{\min}(A) \geq k$. \square

6.5.2 Embedding in Configurations ($\diamond 2$)–($\diamond 5$)

In this section we show how to embed $T_{T_{1,2}}$ in the presence of configurations ($\diamond 2$)–($\diamond 5$). As outlined in Section 6.1.1 our main embedding lemma, Lemma 6.19, builds on Lemma 6.18 which handles Stage 1 of the embedding, and Lemma 6.17 which handles Stage 2.

Lemma 6.17. *Assume we are in Setting 5.1. Suppose L'', L' and Ψ' witness Preconfiguration (\clubsuit)($\frac{10^5 \Omega^*}{\eta}$). Let (T, r) be a rooted tree of order at most $\gamma^2 \nu k / 6$. Let $U \subseteq V(G)$ with $|U| + v(T) \leq k$, and let $v \in \Psi' \setminus U$. Then there exists an $(r \leftrightarrow v, V(T) \leftrightarrow V(G) \setminus U)$ -embedding of (T, r) .*

Proof. We proceed by induction on the order of T . The base $v(T) \leq 2$ obviously holds. Let us assume Lemma 6.17 is true for all trees T' with $v(T') < v(T)$.

Let $U_1 := \text{shadow}_{G_{\nabla}}(U - \Psi, \eta k / 200)$, and $U_2 := \bigcup \{C \in \mathbf{V} : |C \cap U| \geq \frac{1}{2}|C|\}$. We have $|U_1| \leq \frac{200\Omega^*}{\eta}k$ by Fact 4.12, and $|U_2| \leq 2|U|$. Set

$$\begin{aligned} L_{\mathfrak{A}} &:= L'' \cap \text{shadow}_{G_{\nabla}}(\mathfrak{A}, \frac{\eta k}{50}), \\ L_{\Psi} &:= L'' \cap \text{shadow}_{G_{\nabla}}\left(\Psi, |U \cap \Psi| + \frac{\eta k}{50}\right), \text{ and} \\ L_{\mathbf{V}} &:= L'' \cap \text{shadow}_{G_{\text{reg}}}\left(V(G_{\text{reg}}), (1 + \frac{\eta}{50})k - |U \cap \Psi|\right) . \end{aligned}$$

6.5 Main embedding lemmas

Observe that $L_{\mathbf{V}} \subseteq \bigcup \mathbf{V}$ and that since $L'' \subseteq \mathbb{L}_{\frac{9}{10}\eta, k}(G_{\nabla}) \setminus \Psi$, we have

$$L'' \subseteq V(G_{\text{exp}}) \cup \mathfrak{A} \cup L_{\Psi} \cup L_{\mathfrak{A}} \cup L_{\mathbf{V}}.$$

As by (5.18), we have $\deg_G(v, L'') \geq \frac{10^5 \Omega^* k}{\eta} > 5(|U \cup U_1 \cup U_2| + v(T) + \eta k)$, one of the following five cases must occur.

Case I: $\deg_G(v, V(G_{\text{exp}}) \setminus U) > v(T) + \eta k$. Lemma 6.5 gives an embedding of the forest $T-r$ (whose components are rooted at neighbours of r). The input sets/parameters of Lemma 6.5 are $Q_{\text{L6.5}} := 1$, $\zeta_{\text{L6.5}} := 12\sqrt{\gamma}$, $U_{\text{L6.5}}^* := (N_G(v) \cap V(G_{\text{exp}})) \setminus U$, $U_{\text{L6.5}} := U$, $V_1 = V_2 := V(G_{\text{exp}})$.

Case II: $\deg_G(v, \mathfrak{A} \setminus U) > v(T) + \eta k$. Lemma 6.4 gives an embedding of the forest $T-r$ (whose components are rooted at neighbours of r). The input sets/parameters of Lemma 6.4 are $U_{\text{L6.4}}^* := (N_G(v) \cap \mathfrak{A}) \setminus U$, $U_{\text{L6.4}} := U$ and $\varepsilon_{\text{L6.4}} := \varepsilon' \leq \eta$. Here, and below, we tacitly implicitly assume parameters of the same name to be the same, i.e. $\gamma_{\text{L6.4}} := \gamma$.

Case III: $\deg_G(v, L_{\mathfrak{A}} \setminus (U \cup U_1)) > v(T) + \eta k$. We only outline the strategy. Embed the children of r in $L_{\mathfrak{A}} \setminus (U \cup U_1)$ using a map $\phi : \text{Ch}_T(r) \rightarrow L_{\mathfrak{A}} \setminus (U \cup U_1)$. By definition of $L_{\mathfrak{A}}$, and U_1 , we have $\deg_{G_{\nabla}}(\phi(w), \mathfrak{A} \setminus U) > \frac{\eta k}{100}$ for each $w \in \text{Ch}_T(r)$. Now, for every $w \in \text{Ch}_T(r)$ we can proceed as in Case II to extend this embedding to the rooted tree $(T(r, \uparrow w), w)$. That is, Case III is ‘‘Case II with an extra step in the beginning’’.

Case IV: $\deg_G(v, L_{\Psi} \setminus U) > v(T) + \eta k$. We embed the children $\text{Ch}_T(r)$ of r in distinct vertices of $L_{\Psi} \setminus U$. This is possible by the assumption of Case IV.

Now, (5.16) implies that $\deg_{G_{\nabla}}^{\min}(L_{\Psi}, \Psi') \geq |U \cap \Psi| + \frac{\eta k}{100}$. Consequently, $\deg_{G_{\nabla}}^{\min}(L_{\Psi}, \Psi' \setminus U) \geq \frac{\eta k}{100}$. Therefore, for each $w \in \text{Ch}_T(r)$ embedded in $L_{\Psi} \setminus U$ we can find an embedding of $\text{Ch}_T(w)$ in $\Psi' \setminus U$ such that the images of grandchildren of r are disjoint. We fix such an embedding. We can now apply induction. More specifically, for each grandchild u of r we embed the rooted tree $(T(r, \uparrow u), u)$ using Lemma 6.17 (employing induction) using the updated set U , to which the images of the newly embedded vertices were added.

Case V: $\deg_G(v, L_{\mathbf{V}} \setminus (U \cup U_1 \cup U_2)) \geq v(T)$. Let u_1, \dots, u_{ℓ} be the children of r . Let us consider arbitrary distinct neighbours $x_1, \dots, x_{\ell} \in \overline{L_{\mathbf{V}}} \setminus (U \cup U_1 \cup U_2)$ of v . Let $T_i := T(r, \uparrow u_i)$. We sequentially embed the rooted trees (T_i, u_i) , $i = 1, \dots, \ell$, writing ϕ for the embedding. In step i , consider the set $W_i := \left(U \cup \bigcup_{j < i} \phi(T_j) \right) \setminus \Psi$. Let $D_i \in \mathbf{V}$ be the cluster containing x_i . By definition of $L_{\mathbf{V}}$ and of U_1 ,

$$\deg_{G_{\text{reg}}}(x_i, V(G_{\text{reg}}) \setminus W_i) \geq \frac{\eta k}{50} - \frac{\eta k}{200} \geq \frac{\eta k}{100}.$$

Fact 4.8 yields a cluster $C_i \in \mathbf{V}$ such that

$$\deg_{G_{\text{reg}}}(x_i, C_i \setminus W_i) \geq \frac{\eta}{100} \cdot \frac{\gamma \mathfrak{c}}{2(\Omega^*)^2} > \frac{\gamma^2 \mathfrak{c}}{2} + v(T) > \frac{12\varepsilon' \mathfrak{c}}{\gamma^2} + v(T).$$

In particular there is at least one edge from $E(G_{\text{reg}})$ between C_i and D_i , and therefore, (C_i, D_i) forms an ε' -regular pair of density at least γ^2 in G_{reg} . Map u_i to x_i and let F_1, \dots, F_m be the components of the forest $T_i - u_i$. We now sequentially embed the trees F_j in the pair (D_i, C_i) using Lemma 6.6, with $X_{\text{L6.6}} := C_i \setminus (W_i \cup \bigcup_{q < j} \phi(F_q))$, $X_{\text{L6.6}}^* := N_{G_{\text{reg}}}(x_i, X_{\text{L6.6}})$, $Y_{\text{L6.6}} := D_i \setminus (W_i \cup \{x_i\} \cup \bigcup_{q < j} \phi(F_q))$, $\varepsilon_{\text{L6.6}} := \varepsilon'$, and $\beta_{\text{L6.6}} := \gamma^2/3$. \square

We are now ready for the lemma that will handle Stage 1 in configurations $(\diamond 2)$ – $(\diamond 5)$.

Lemma 6.18. *Assume we are in Setting 5.1, with L'', L', Ψ' witnessing $(\clubsuit)(\Omega^*)$ in G . Let $U \subseteq V(G) \setminus \Psi$ and let (T, r) be a rooted tree with $v(T) \leq k/2$ and $|U| + v(T) \leq k$. Suppose that each component of $T - r$ has order at most τk . Let $x \in (L'' \cap \mathbb{YB}) \setminus \bigcup_{i=0}^2 \text{shadow}_{G_{\nabla}}^{(i)}(\text{ghost}(U), \eta k/1000)$.*

Then there is a subtree T' of T with $r \in V(T')$ which has an $(r \hookrightarrow x, V(T') \setminus \{r\} \hookrightarrow V(G) \setminus \Psi)$ -embedding ϕ . Further, the components of $T - T'$ can be partitioned into two (possibly empty) sets $\mathcal{C}_1, \mathcal{C}_2$, such that the following two assertions hold.

(a) *If $\mathcal{C}_1 \neq \emptyset$, then $\deg_{G_{\nabla}}^{\min}(\phi(\text{Par}(V(\bigcup \mathcal{C}_1))), \Psi') > k + \frac{\eta k}{100} - v(T')$,*

(b) *$\text{Par}(V(\bigcup \mathcal{C}_2)) \subseteq \{r\}$, and $\deg_{G_{\nabla}}(x, \Psi') > \frac{k}{2} + \frac{\eta k}{100} - v(T' \cup \bigcup \mathcal{C}_1)$.*

Proof. Let \mathcal{C} be the set of all components of $T - r$. We start by defining \mathcal{C}_2 . Then, we have to distribute $T - \bigcup \mathcal{C}_2$ between T' and \mathcal{C}_1 . First, we find a set $\mathcal{C}_M \subseteq \mathcal{C} \setminus \mathcal{C}_2$ which fits into the matching $\mathcal{M}_A \cup \mathcal{M}_B$ (and thus will form part of T'). Then, we consider the remaining components of $\mathcal{C} \setminus \mathcal{C}_2$: some of these will be embedded entirely, of others we only embed the root, and leave the rest for \mathcal{C}_1 . Everything embedded will become a part of T' .

Throughout the proof we write shadow for $\text{shadow}_{G_{\nabla}}$.

Set $\overline{V_{\text{good}}} := V_{\text{good}} \setminus \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000})$, and choose $\tilde{\mathcal{C}} \subseteq \mathcal{C}$ such that

$$\deg_{G_{\nabla}}(x, \overline{V_{\text{good}}}) - \frac{\eta k}{30} < \sum_{S \in \tilde{\mathcal{C}}} v(S) \leq \max \left\{ 0, \deg_{G_{\nabla}}(x, \overline{V_{\text{good}}}) - \frac{\eta k}{40} \right\}. \quad (6.12)$$

Set $\mathcal{C}_2 := \mathcal{C} \setminus \tilde{\mathcal{C}}$. Note that this choice clearly satisfies the first part of (b). Let us now verify the second part of (b). For this, we calculate

$$\begin{aligned} \deg_{G_{\nabla}}(x, \Psi') &\geq \deg_{G_{\nabla}}(x, V_+ \setminus L_{\#}) - \deg_{G_{\nabla}}(x, \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000})) \\ &\quad - \deg_{G_{\nabla}}(x, V_+ \setminus (L_{\#} \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}) \cup \Psi)) \\ &\quad - \deg_{G_{\nabla}}(x, \Psi \setminus \Psi') \\ &\stackrel{\text{(by (5.9), } x \notin \text{shadow}^{(2)}(\text{ghost}(U), \frac{\eta k}{1000}), \text{(6.12), (5.16))}}{\geq} \left(\frac{k}{2} + \frac{\eta k}{20} \right) - \frac{\eta k}{1000} - \left(\sum_{S \in \tilde{\mathcal{C}}} v(S) + \frac{\eta k}{30} \right) - \frac{\eta k}{100} \\ &> \frac{k}{2} - \sum_{S \in \tilde{\mathcal{C}}} v(S) + \frac{\eta k}{20} \\ &\geq \frac{k}{2} - v(T' \cup \bigcup \mathcal{C}_1) + \frac{\eta k}{100}, \end{aligned}$$

as desired for (b).

Now, set

$$\mathcal{M} := \{(X_1, X_2) \in \mathcal{M}_A \cup \mathcal{M}_B : \deg_{G_{\mathcal{D}}}(x, (X_1 \cup X_2) \setminus \mathfrak{A}) > 0\}. \quad (6.13)$$

Claim 6.18.1. We have $|V(\mathcal{M})| \leq \frac{4(\Omega^*)^2}{\gamma^2} k$.

6.5 Main embedding lemmas

Proof of Claim 6.18.1. Indeed, let $(X_1, X_2) \in \mathcal{M}$, i.e. $(X_1, X_2) \in \mathcal{M}_A \cup \mathcal{M}_B$ with $\deg_{G_{\mathcal{D}}}(x, (X_1 \cup X_2) \setminus \mathfrak{A}) > 0$. Then, using Property 4 of Setting 5.1, we see that there exists a cluster $C_{(X_1, X_2)} \in \mathbf{V}$ such that $\deg_{G_{\mathcal{D}}}(x, C_{(X_1, X_2)}) > 0$, and either $X_1 \subseteq C_{(X_1, X_2)}$ or $X_2 \subseteq C_{(X_1, X_2)}$. In particular, there exists a dense spot $(A_{(X_1, X_2)}, B_{(X_1, X_2)}; F_{(X_1, X_2)}) \in \mathcal{D}$ such that $x \in A_{(X_1, X_2)}$, and $X_1 \subseteq B_{(X_1, X_2)}$ or $X_2 \subseteq B_{(X_1, X_2)}$. By Fact 4.4, there are at most $\frac{\Omega^*}{\gamma}$ such dense spots, let Z denote the union of all vertices contained in these spots. Fact 4.3 implies that $|Z| \leq \frac{2(\Omega^*)^2}{\gamma^2}k$. Thus $|V(\mathcal{M})| \leq 2|V(\mathcal{M}) \cap Z| \leq 2|Z| \leq \frac{4(\Omega^*)^2}{\gamma^2}k$. \square

First we shall embed as many components from $\tilde{\mathcal{C}}$ as possible in \mathcal{M} . To this end, consider an inclusion-maximal subset \mathcal{C}_M of $\tilde{\mathcal{C}}$ with

$$\sum_{S \in \mathcal{C}_M} v(S) \leq \deg_{G_{\nabla}}(x, V(\mathcal{M})) - \frac{\eta k}{1000}. \quad (6.14)$$

We aim to utilize the degree of x to $V(\mathcal{M})$ to embed \mathcal{C}_M in $V(\mathcal{M})$ using the regularity method.

Remark 6.18.2. This remark (which may as well be skipped at a first reading) is aimed at those readers that are wondering about a seeming inconsistency of the defining formulas (6.13) for \mathcal{M} , and (6.14) for \mathcal{C}_M . That is, (6.13) involves the degree in $G_{\mathcal{D}}$ and excludes the set \mathfrak{A} , while (6.14) involves the degree in G_{∇} . The setting in (6.13) was chosen so that it allows us to control the size of \mathcal{M} in Claim 6.18.1, crucially relying on Property 4 of Setting 5.1. Such a control is necessary to make the regularity method work. Indeed, in each regular pair there may be a small number of atypical vertices⁹, and we must avoid these vertices when embedding the components by the regularity method. Thus without the control on $|\mathcal{M}|$ it might happen that the degree of x is unusable because x sees very small numbers of atypical vertices in an enormous number of sets corresponding to \mathcal{M} -vertices. On the other hand, the edges x sends to \mathfrak{A} can be utilized by other techniques in later stages. Once we have defined \mathcal{M} we want to use the full degree to $V(\mathcal{M})$ to ensure we can embed the shrubs as balanced as possible into the \mathcal{M} -edges. This is necessary as otherwise part of the degree of x might be unusable for embedding, e.g. because it might go to \mathcal{M} -vertices whose partners are already full.

For each $(C, D) \in \mathcal{M}$ we choose $\mathcal{C}_{CD} \subseteq \mathcal{C}_M$ maximal such that

$$\sum_{S \in \mathcal{C}_{CD}} v(S) \leq \deg_{G_{\nabla}}(x, (C \cup D) \setminus \text{ghost}(U)) - \left(\frac{\gamma}{\Omega^*}\right)^3 |C|, \quad (6.15)$$

and further, we require \mathcal{C}_{CD} to be disjoint from families $\mathcal{C}_{C'D'}$ defined in previous steps. We claim that $\{\mathcal{C}_{CD}\}_{(C,D) \in \mathcal{M}}$ forms a partition of \mathcal{C}_M , i.e., all the elements of \mathcal{C}_M are used. Indeed, otherwise, by the maximality of \mathcal{C}_{CD} and since the components of $T - r$ have size at most τk , we obtain

$$\begin{aligned} \sum_{S \in \mathcal{C}_{CD}} v(S) &\geq \deg_{G_{\nabla}}(x, (C \cup D) \setminus \text{ghost}(U)) - \left(\frac{\gamma}{\Omega^*}\right)^3 |C| - \tau k \\ &\stackrel{(5.1)}{\geq} \deg_{G_{\nabla}}(x, (C \cup D) \setminus \text{ghost}(U)) - 2\left(\frac{\gamma}{\Omega^*}\right)^3 |C|, \end{aligned} \quad (6.16)$$

⁹The issue of atypicality itself could be avoided by preprocessing each pair (S, T) of $\mathcal{M}_A \cup \mathcal{M}_B$ and making it super-regular. However this is not possible for atypicality with respect to a given (but unknown in advance) subpair (S', T') .

for each $(C, D) \in \mathcal{M}$. Then we have

$$\begin{aligned}
 \sum_{S \in \mathcal{C}_M} v(S) &> \sum_{(C, D) \in \mathcal{M}} \sum_{S \in \mathcal{C}_{CD}} v(S) \\
 &\stackrel{\text{(by (6.16))}}{\geq} \sum_{(C, D) \in \mathcal{M}} \left(\deg_{G_\nabla}(x, (C \cup D) \setminus \text{ghost}(U)) - 2\left(\frac{\gamma}{\Omega^*}\right)^3 |C| \right) \\
 &\stackrel{\text{(by Claim 6.18.1 and Fact 4.11)}}{\geq} \deg_{G_\nabla}(x, V(\mathcal{M}) \setminus \text{ghost}(U)) - 2\left(\frac{\gamma}{\Omega^*}\right)^3 \cdot \frac{2(\Omega^*)^2}{\gamma^2} k \\
 &\stackrel{\text{(as } x \notin \text{shadow}(\text{ghost}(U)))}{\geq} \deg_{G_\nabla}(x, V(\mathcal{M})) - \frac{\eta k}{1000} \\
 &\stackrel{\text{(by (6.14))}}{\geq} \sum_{S \in \mathcal{C}_M} v(S),
 \end{aligned}$$

a contradiction.

We use Lemma 6.7 to embed the components of \mathcal{C}_{CD} in $(C \cup D) \setminus \text{ghost}(U)$ with the following setting: $C_{L6.7} := C$, $D_{L6.7} := D$, $U_{L6.7} := \text{ghost}(U)$, $X_{L6.7}^* := (N_{G_\nabla}(x) \cap (C \cup D)) \setminus U_{L6.7}$, and (T_i, r_i) are the rooted trees from \mathcal{C}_{CD} with the roots being the neighbours of r . The constants in Lemma 6.7 are $\varepsilon_{L6.7} := \varepsilon'$, $\beta_{L6.7} := \sqrt{\varepsilon'}$, and $\ell_{L6.7} := |C| \geq \nu\pi k$. The rooted trees in \mathcal{C}_{CD} are smaller than $\varepsilon_{L6.7}\ell_{L6.7}$ by (5.1). Condition (6.2) is satisfied by (6.15), and since $(\gamma/\Omega^*)^3 \geq 50\sqrt{\varepsilon'}$.

It remains to deal with the components $\tilde{\mathcal{C}} \setminus \mathcal{C}_M$. In the sequel we shall assume that $\tilde{\mathcal{C}} \setminus \mathcal{C}_M \neq \emptyset$ (otherwise skip this step and go directly to the definition of T' and \mathcal{C}_1 , with $p = 0$). Thus, by our choice of \mathcal{C}_M , we have

$$\sum_{S \in \mathcal{C}_M} v(S) \geq \deg_{G_\nabla}(x, V(\mathcal{M})) - \frac{\eta k}{900}. \quad (6.17)$$

Let T_1, T_2, \dots, T_p be the trees of $\tilde{\mathcal{C}} \setminus \mathcal{C}_M$ rooted at the vertices $r_i \in \text{Ch}(r) \cap V(T_i)$. We shall sequentially extend our embedding of \mathcal{C}_M to subtrees $T'_i \subseteq T_i$. Let $U_i \subseteq V(G)$ be the union of the images of $\bigcup \mathcal{C}_M \cup \{r\}$ and of T'_1, \dots, T'_i under this embedding.

Suppose that we have embedded the trees T'_1, \dots, T'_i for some $i = 0, 1, \dots, p-1$. We claim that at least one of the following holds.

- (V1) $\deg_{G_\nabla}(x, V(G_{\text{exp}}) \setminus (U \cup U_i)) \geq \frac{\eta k}{1000}$,
- (V2) $\deg_{G_\nabla}(x, \mathfrak{A} \setminus (U \cup U_i)) \geq \frac{\eta k}{1000}$, or
- (V3) $\deg_{G_\nabla}(x, L' \setminus (V(G_{\text{exp}}) \cup \mathfrak{A} \cup U \cup U_i \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}))) \geq \frac{\eta k}{1000}$.

Indeed, suppose that none of (V1)–(V3) holds. Then, first note that since $U \subseteq \text{ghost}(U)$ and since $x \notin \text{shadow}(\text{ghost}(U), \eta k/1000)$, we have

$$\deg_{G_\nabla}(x, U) \leq \eta k/1000. \quad (6.18)$$

Also,

$$\deg_{G_D}(x, V(\mathcal{M}_A \cup \mathcal{M}_B)) \leq \deg_{G_D}(x, V(\mathcal{M}) \cup \mathfrak{A}). \quad (6.19)$$

Thus,

$$\begin{aligned}
 & \deg_{G_{\nabla}} \left(x, V_{\text{good}} \setminus \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}) \right) \\
 \stackrel{\text{(by (6.18) and (6.19), def of } V_{\text{good}})}{\leq} & \deg_{G_{\nabla}} \left(x, (V(\mathcal{M}) \cup V(G_{\text{exp}}) \cup \mathfrak{A} \cup L') \setminus (U \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000})) \right) \\
 & + \deg_{G_{\nabla}} \left(x, \mathbb{L}_{\frac{9}{10}\eta, k}(G_{\nabla}) \setminus (\Psi \cup L') \right) + \frac{\eta k}{1000} \\
 \stackrel{\text{(by (5.18))}}{\leq} & \deg_{G_{\nabla}} \left(x, (V(G_{\text{exp}}) \cup \mathfrak{A} \cup L') \setminus (V(\mathcal{M}) \cup U \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000})) \right) \\
 & + \deg_{G_{\nabla}}(x, V(\mathcal{M})) + \frac{\eta k}{100} + \frac{\eta k}{1000} \\
 \stackrel{\text{(by } \neg(\mathbf{V1}), \neg(\mathbf{V2}), \neg(\mathbf{V3}), \text{ by (6.17))}}{\leq} & 3 \cdot \frac{\eta k}{1000} + \sum_{j=1}^i v(T'_j) + \sum_{S \in \mathcal{C}_M} v(S) + \frac{\eta k}{900} + \frac{\eta k}{100} + \frac{\eta k}{1000} \\
 & < \sum_{S \in \tilde{\mathcal{C}}} v(S) + \frac{\eta k}{40},
 \end{aligned}$$

a contradiction to (6.12).

In cases **(V1)**–**(V2)** we shall embed the entire tree $T'_{i+1} := T_{i+1}$. In case **(V3)** we either embed the entire tree $T'_{i+1} := T_{i+1}$, or embed only one vertex $T'_{i+1} := r_{i+1}$ (that will only happen in case **(V3c)**). In the latter case, we keep track of the components of $T_{i+1} - r_{i+1}$ in the set $\mathcal{C}_{1,i+1}$ (we tacitly assume we set $\mathcal{C}_{1,i+1} := \emptyset$ in all cases other than **(V3c)**). The union of the sets $\mathcal{C}_{1,i}$ will later form the set \mathcal{C}_1 . Let us go through our three cases in detail.

In case **(V1)** we embed T_{i+1} rooted at r_{i+1} using Lemma 6.5 for one tree (i.e. $\ell_{L6.5} := 1$) with the following sets/parameters: $H_{L6.5} := G_{\text{exp}}$, $U_{L6.5} := U \cup U_i$, $U_{L6.5}^* := N_{G_{\nabla}}(x) \cap (V(G_{\text{exp}}) \setminus (U \cup U_i))$, $V_1 = V_2 := V(G_{\text{exp}})$, $Q_{L6.5} := 1$, $\zeta_{L6.5} := \rho$, and $\gamma_{L6.5} := \gamma$. Note that $|U \cup U_i| < k$, that $|N_{G_{\nabla}}(x) \cap (V(G_{\text{exp}}) \setminus (U \cup U_i))| \geq \eta k / 1000 > 32\gamma k / \rho + 1$, that $v(T_{i+1}) \leq \tau k < \rho k / 4$ and that $128\gamma < \rho^2$.

In case **(V2)** we embed T_{i+1} rooted at r_{i+1} using Lemma 6.4 for one tree (i.e. $\ell_{L6.4} := 1$) with the following setting: $H_{L6.4} := G - \Psi$, $\mathfrak{A}_{L6.4} := \mathfrak{A}$, $U_{L6.4} := U \cup U_i$, $U_{L6.4}^* := N_{G_{\nabla}}(x) \cap (\mathfrak{A} \setminus (U \cup U_i))$, $\Lambda_{L6.4} := \Lambda$, $\gamma_{L6.4} := \gamma$, $\varepsilon_{L6.4} := \varepsilon'$. Note that $|U \cup U_i| \leq k < \Lambda k$, that $|N_{G_{\nabla}}(x) \cap (\mathfrak{A} \setminus (U \cup U_i))| \geq \eta k / 1000 > 2\varepsilon' k$, and that $v(T_{i+1}) \leq \tau k < \gamma k / 2$.

We commence case **(V3)** with an auxiliary claim.

Claim 6.18.3. There exists $C_0 \in \mathbf{V}$ such that

$$\deg_{G_{\mathcal{D}}}(x, (C_0 \cap L') \setminus (V(G_{\text{exp}}) \cup U \cup U_i \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}))) \geq \frac{\varepsilon'}{\gamma^2} \mathbf{c}.$$

Proof of Claim 6.18.3. Observe that $L' \setminus (V(G_{\text{exp}}) \cup \mathfrak{A} \cup \Psi \cup U \cup U_i) \subseteq \bigcup \mathbf{V}$ and that (since $x \in \bigcup \mathbf{V}$)

$$E_{G_{\nabla}}[x, L' \setminus (V(G_{\text{exp}}) \cup \mathfrak{A} \cup U \cup U_i \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}))] \subseteq E(G_{\mathcal{D}}).$$

By Fact 4.8, there are at most $\frac{2(\Omega^*)^2 k}{\gamma^2 \mathbf{c}}$ clusters $C \in \mathbf{V}$ such that $\deg_{G_{\mathcal{D}}}(x, C) > 0$. Using the

assumption **(V3)**, there exists a cluster $C_0 \in \mathbf{V}$ such that

$$\begin{aligned} \deg_{G_{\mathcal{D}}} \left(x, (C_0 \cap L') \setminus (V(G_{\text{exp}}) \cup U \cup U_i \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000})) \right) &\geq \frac{\eta k}{1000} \cdot \frac{\gamma^2 \mathbf{c}}{2(\Omega^*)^2 k} \\ &\stackrel{(5.1)}{\geq} \frac{\varepsilon'}{\gamma^2} \mathbf{c}, \end{aligned}$$

as desired. \square

Let us take a cluster C_0 from Claim 6.18.3. We embed the root r_{i+1} of T_{i+1} in an arbitrary neighbour y of x in $(C_0 \cap L') \setminus (V(G_{\text{exp}}) \cup U \cup U_i \cup \text{shadow}(\text{ghost}(U), \frac{\eta k}{1000}))$.

Let $H \subseteq G$ be the subgraph of G consisting of all edges in dense spots \mathcal{D} , and all edges incident with Ψ' . As by (5.16), y has at most $\eta k/100$ neighbours in $\Psi \setminus \Psi'$, and since $y \in L' \subseteq \mathbb{L}_{9\eta/10, k}(G_{\nabla})$ and $y \notin \text{shadow}(U, \frac{\eta k}{100})$, we find that

$$\begin{aligned} \deg_H(y, V(G) \setminus ((U \cup U_i) \cup (\Psi \setminus \Psi'))) &\geq \left(1 + \frac{9\eta}{10}\right) k - \frac{\eta k}{1000} - |U_i| - \frac{\eta k}{100} \\ &> k - |U_i| + \frac{\eta k}{2}. \end{aligned}$$

Therefore, one of the three following subcases must occur. (Recall that $y \notin \mathfrak{A}$ as $y \in C_0 \in \mathbf{V}$.)

- (V3a)** $\deg_{G_{\nabla}}(y, \mathfrak{A} \setminus (U \cup U_i)) \geq \frac{\eta k}{6}$,
- (V3b)** $\deg_{G_{\text{reg}}}(y, \bigcup \mathbf{V} \setminus (U \cup U_i)) \geq \frac{\eta k}{6}$, or
- (V3c)** $\deg_{G_{\nabla}}(y, \Psi') \geq k - |U_i| + \frac{\eta k}{6}$.

In case **(V3a)** we embed the components of $T_{i+1} - r_{i+1}$ (as trees rooted at the children of r_{i+1}) using the same technique as in case **(V2)**, with Lemma 6.4.

In **(V3b)** we embed the components of $T_{i+1} - r_{i+1}$ (as trees rooted at the children of r_{i+1}). By Fact 4.8 there exists a cluster $D \in \mathbf{V}$ such that

$$\deg_{G_{\text{reg}}}(y, D \setminus (U \cup U_i)) \geq \frac{\eta k}{6} \cdot \frac{\gamma^2 \mathbf{c}}{2(\Omega^*)^2 k} > \frac{\gamma^2}{2} \mathbf{c}. \quad (6.20)$$

We use Lemma 6.6 with input $\varepsilon_{\text{L6.6}} := \varepsilon'$, $\beta_{\text{L6.6}} := \gamma^2$, $C_{\text{L6.6}} := D$, $D_{\text{L6.6}} := C_0$, $X_{\text{L6.6}}^* = X_{\text{L6.6}} := D \setminus (U \cup U_i)$ and $Y_{\text{L6.6}} := C_0 \setminus (U \cup U_i \cup \{y\})$ to embed the tree T_{i+1} into the pair (C_0, D) , by embedding the components of $T_{i+1} - r_{i+1}$ one after the other. The numerical conditions of Lemma 6.6 hold because of Claim (6.18.3) and because of (6.20).

In case **(V3c)** we set $T'_{i+1} := r_{i+1}$ and define $\mathcal{C}_{1, i+1}$ as set of all components of $T_{i+1} - r_{i+1}$. Then $\phi(\text{Par}(\bigcup \mathcal{C}_{1, i+1}) \cap V(T'_{i+1})) = \{y\}$ and

$$\deg_{G_{\nabla}}(y, \Psi') \geq k - |U_i| + \frac{\eta k}{6}. \quad (6.21)$$

When all the trees T_1, \dots, T_p are processed, we define $T' := \{r\} \cup \bigcup \mathcal{C}_M \cup \bigcup_{i=1}^p T'_i$, and set $\mathcal{C}_1 := \bigcup_{i=1}^p \mathcal{C}_{1, i}$. Thus also (a) is satisfied by (6.21) for $i = p$, since $|T'| = |U_p|$. This finishes the proof of the lemma. \square

It turns out that our techniques for embedding a tree $T \in \mathbf{trees}(k)$ for Configurations $(\diamond 2)$ – $(\diamond 5)$ are very similar. In Lemma 6.19 below we resolve these tasks at once. The proof of Lemma 6.19 follows the same basic strategy for each of the configurations $(\diamond 2)$ – $(\diamond 5)$ and deviates only in the elementary procedures of embedding shrubs of T .

Lemma 6.19. *Suppose that we are in Setting 5.1, and one of the following configurations can be found in G :*

- a) Configuration $(\diamond 2)$ $((\Omega^*)^2, 5(\Omega^*)^9, \rho^3)$,
- b) Configuration $(\diamond 3)$ $((\Omega^*)^2, 5(\Omega^*)^9, \gamma/2, \gamma^3/100)$,
- c) Configuration $(\diamond 4)$ $((\Omega^*)^2, 5(\Omega^*)^9, \gamma/2, \gamma^4/100)$, or
- d) Configuration $(\diamond 5)$ $((\Omega^*)^2, 5(\Omega^*)^9, \varepsilon', 2/(\Omega^*)^3, \frac{1}{(\Omega^*)^5})$,

Let (T, r) be a rooted tree of order k with a (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$. Then $T \subseteq G$.

Proof. First observe that each of the configurations given by a)–d) contains two sets $\Psi'' \subseteq \Psi$ and $V_1 \subseteq V(G) \setminus \Psi$ with

$$\deg_{G_{\nabla}}^{\min}(\Psi'', V_1) \geq 5(\Omega^*)^9 k, \quad (6.22)$$

$$\deg_{G_{\nabla}}^{\min}(V_1, \Psi'') \geq \varepsilon' k. \quad (6.23)$$

For any vertex $z \in W_A \cup W_B$ we define $T(z)$ as the forest consisting of all components of $T - (W_A \cup W_B)$ that contain children of z . Throughout the proof, we write ϕ for the current partial embedding of T into G .

Overview of the embedding procedure. As outlined in Section 6.1.1 the embedding scheme is the same for Configurations $(\diamond 2)$ – $(\diamond 5)$. The embedding ϕ is defined in two stages. In Stage 1, we embed $W_A \cup W_B$, all the internal shrubs, all the end shrubs of \mathcal{S}_A , and a part¹⁰ of the end shrubs of \mathcal{S}_B . In Stage 2 we embed the rest of \mathcal{S}_B . Which part of \mathcal{S}_B are embedded in Stage 1 and which part in Stage 2 will be determined during Stage 1. We first give a rough outline of both stages listing some conditions which we require to be met, and then we describe each of the stages in detail.

Stage 1 is defined in $|W_A \cup \{r\}|$ steps. First we map r to any vertex in Ψ'' . Then in each step we pick a vertex $x \in W_A$ for which the embedding ϕ has already been defined but such that ϕ is not yet defined for any of the children of x . In this step we embed $T(x)$, together with all the children and grandchildren of x in the knag which contains x . For each $y \in W_B \cap \text{Ch}(x)$, Lemma 6.18 determines a subforest $T'(y) \subseteq T(y)$ which is embedded in Stage 1, and sets $\mathcal{C}_1(y)$ and $\mathcal{C}_2(y)$, which will be embedded in Stage 2.

The embedding in each step of Stage 1 will be defined so that the following properties hold.

- (*1) All vertices from W_A are mapped to Ψ'' .
- (*2) All vertices except for W_A are mapped to $V(G) \setminus \Psi$.

¹⁰in the sense that individual shrubs \mathcal{S}_B may be embedded only in part




Figure 6.6: Stage 1 of the embedding in the proof of Lemma 6.19. Starting from an already embedded vertex $x \in W_A$ we extend the embedding to (in this order)

- (1) all the children $y \in W_B$ of x in the same knag (in grey),
- (2) a part $T'(y)$ of the forest $T(y)$,
- (3) all the grandchildren $x' \in W_A$ of x in the same knag,
- (4) the forest $T(x)$ together with the bordering cut-vertices $x^* \in W_A$.

(*3) For each $y \in W_B$, for each $v \in \text{Par}(V(\bigcup \mathcal{C}_1(y)))$ it holds that

$$\deg_G(\phi(v), \Psi') \geq k + \frac{\eta k}{100} - v(T'(y)) .$$

(*4) For each $y \in W_B$, for each $v \in \text{Par}(V(\bigcup \mathcal{C}_2(y)))$ it holds that

$$\deg_G(\phi(v), \Psi') \geq \frac{k}{2} + \frac{\eta k}{100} - v(T'(y) \cup \bigcup \mathcal{C}_1(y)) .$$

In Stage 2, we shall utilize properties (*3) and (*4) to embed $T_B^* := \bigcup S_B - \bigcup_{y \in W_B} T'(y)$. Stage 2 is substantially simpler than Stage 1; this is due to the fact that T_B^* consists only of end shrubs.

The embedding step of Stage 1. The embedding step is the same for Configurations ($\diamond 2$)–($\diamond 5$), except for the embedding of internal shrubs. The order of the embedding steps is illustrated in Figure 6.6.

In each step we have picked $x \in W_A$ already embedded in G but such that none of $\text{Ch}(x)$ are embedded. By (*1), or by the choice of $\phi(r)$, we have $\phi(x) \in \Psi''$. So by (6.22) we have

$$\deg_{G_{\nabla}}(\phi(x), V_1 \setminus U) \geq 5(\Omega^*)^9 k - k. \quad (6.24)$$

First, we embed successively in $|W_B \cap \text{Ch}(x)|$ steps the vertices $y \in W_B \cap \text{Ch}(x)$ together with components $T'(y) \subseteq T(y)$ which will be determined on the way. Suppose that in a certain step we are to embed $y \in W_B \cap \text{Ch}(x)$ and the (to be determined) tree $T'(y)$. Let $F := \bigcup_{i=0}^2 \text{shadow}_{G_{\nabla} - \Psi}^{(i)}(\text{ghost}(U), \frac{\eta k}{10^5})$, where U is the set of vertices used by the embedding ϕ in previous steps, so $|U| \leq k$. By Fact 4.12, $|F| \leq \frac{10^{10}(\Omega^*)^2}{\eta^2} k$. We embed y anywhere in $(N_G(\phi(x)) \cap V_1) \setminus F$, cf. (6.22). Note that then (*2) holds for y . We use Lemma 6.18 in order to embed $T'(y) \subseteq T(y)$ (the subtree $T'(y)$ is determined by Lemma 6.18). Lemma 6.18 ensures that (*3) and (*4) hold and that we have $\phi(V(T'(y))) \subseteq V(G) \setminus \Psi$.

6.5 Main embedding lemmas

Also, we map the vertices $x' \in W_A \cap \text{Ch}(y)$ to $\Psi'' \setminus U$. To justify this step, employing (*2), it is enough to prove that

$$\deg(\phi(y), \Psi'') \geq |W_A|. \quad (6.25)$$

Indeed, on one hand, we have $|W_A| \leq 336/\tau$ by Definition 3.3(c). On the other hand, we have that $\phi(y) \in V_1$, and thus (6.23) applies. We can thus embed x' as planned, ensuring (*1), and finishing the step for y .

Next, we sequentially embed the components \tilde{T} of $T(x)$. In the following, we describe such an embedding procedure only for an internal shrub \tilde{T} , with x^* denoting the other neighbour of \tilde{T} in W_A (cf. (*1)). The case when \tilde{T} is an end shrub is analogous: actually it is even easier as we do not have to worry about placing x^* well. The actual embedding of \tilde{T} together with x^* depends on the configuration we are in. We shall slightly abuse notation by letting U now denote everything embedded before the tree \tilde{T} .

For Configuration ($\diamond 2$), we use Lemma 6.5 for one tree, namely $\tilde{T} - x^*$, using the following setting: $Q_{L6.5} := 1, \gamma_{L6.5} := \gamma, \zeta_{L6.5} := \rho^3, H_{L6.5} := G_{\text{exp}}, U_{L6.5} := U$, and $U_{L6.5}^* := (N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$ (this last set is large enough by (6.24)). The child of x gets embedded in $(N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$, the vertices at odd distance from x get embedded in V_1 , and the vertices at even distance from x get embedded in V_2 . In particular, $\text{Par}_T(x^*)$ gets embedded in V_1 . After this, we accomodate x^* in a vertex in $\Psi'' \setminus U$ which is adjacent to $\phi(\text{Par}_T(x^*))$. This is possible by the same reasoning as in (6.25).

For Configuration ($\diamond 3$), we use Lemma 6.14 to embed \tilde{T} with the setting $\gamma_{L6.14} := \gamma, \delta_{L6.14} := \gamma^3/100, U_{L6.14} := U$ and $U_{L6.14}^* := (N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$ (this last set is large enough by (6.24)). Then the child of x gets embedded in $(N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$, vertices of \tilde{T} of odd distance to x (i.e. of even distance to the root of \tilde{T}) get embedded in $V_1 \setminus U$, and vertices of even distance get embedded in $V_2 \setminus U$. We extend the embedding by mapping x^* to a suitable vertex in $\Psi'' \setminus U$ adjacent to $\phi(\text{Par}_T(x^*))$ in the same way as above.

For Configuration ($\diamond 4$), we use Lemma 6.15 to embed \tilde{T} with the setting $\gamma_{L6.15} := \gamma, \delta_{L6.15} := \gamma^4/100, U_{L6.15} := U$ and $U_{L6.15}^* := (N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$ (this last set is large enough by (6.24)). The fruit $r'_{L6.15}$ in the lemma is chosen as $\text{Par}_T(x^*)$, note that this is indeed a fruit (in \tilde{T}) because of Definition 3.3 (i). Then the child of x gets embedded in $(N_{G_{\nabla}}(\phi(x)) \cap V_1) \setminus U$, the vertex $r'_{L6.15} = \text{Par}_T(x^*)$ gets embedded in $V_1 \setminus U$, and the rest of \tilde{T} gets embedded in $(\mathfrak{A}' \cup V_2) \setminus U$. This allows us to extend the embedding to x^* as above.

In Configuration ($\diamond 5$), let $\mathbf{W} \subseteq \mathbf{V}$ denote the set of those clusters, which have at least an $\frac{1}{2(\Omega^*)^5}$ -fraction of their vertices contained in the set $U' := U \cup \text{shadow}_{G_{\text{reg}}}(U, k/(\Omega^*)^3)$. We get from Fact 4.12 that $|U'| \leq 2(\Omega^*)^4 k$, and consequently $|U' \cup \bigcup \mathbf{W}| \leq 4(\Omega^*)^9 k$. By (6.24) we can find a vertex $v \in (N_G(\phi(x)) \cap V_1) \setminus (U' \cup \bigcup \mathbf{W})$.

We use the fact that $v \notin \text{shadow}_{G_{\text{reg}}}(U, k/(\Omega^*)^3)$ together with inequality (5.27) to see that $\deg_{G_{\text{reg}}}(v, V(G_{\text{reg}}) \setminus U) \geq k/(\Omega^*)^3$. Now, since there are only boundedly many clusters seen from v (cf. Fact 4.8), there must be a cluster $D \in \mathbf{V}$ such that

$$\deg_{G_{\text{reg}}}(v, D \setminus U) \geq \frac{\gamma^2}{2 \cdot (\Omega^*)^5} |D| \geq \gamma^3 |D|. \quad (6.26)$$

Let C be the cluster containing v . We have $|(C \cap V_1) \setminus U| \geq \frac{1}{2(\Omega^*)^5} |C| \geq \gamma^3 |C|$ because of (5.28) and since $C \notin \mathbf{W}$. Thus, by Fact 2.1, $((C \cap V_1) \setminus U, D \setminus U)$ is an $2\epsilon'/\gamma^3$ -regular pair of density at least

$\gamma^2/2$. We can therefore embed \tilde{T} in this pair using the regularity method. Moreover, by (6.26), we can do so by mapping the child z of x to v . Thus the parent of x^* (lying at even distance to z) will be embedded in $(C \cap V_1) \setminus U$. We can then extend our embedding to x^* as above.

This finishes our embedding of $T(x)$. Note that in all cases we have $\phi(x^*) \in \Psi''$ and $\phi(V(\tilde{T})) \subseteq V(G) \setminus \Psi$, as required by (*1) and (*2).

The embedding steps of Stage 2. For $i = 1, 2$, set $Z_i := \bigcup_{y \in W_B} \text{Ch}(T'(y)) \cap \bigcup \mathcal{C}_i(y)$.

First, we embed all the vertices $z \in Z_2$ in Ψ' . By (*2), until now, only vertices of $W_A \cup Z_2$ are mapped to Ψ' , and using (*4) and the properties (c), (k) and (l) of Definition 3.3, we see that

$$\begin{aligned} \deg_G(\phi(\text{Par}(z)), \Psi') &\geq \frac{\eta k}{100} + \left(\frac{k}{2} - \bigcup_{y \in W_B} (T'(y) \cup \bigcup \mathcal{C}_1(y))\right) \\ &> |W_A| + |Z_2|. \end{aligned}$$

So there is space for the vertex z in $\Psi' \cap \phi(N_G(\text{Par}(z)))$.

Next, we embed all the vertices $z \in Z_1$ in Ψ' . By (*2), until now, only vertices of $W_A \cup Z_2 \cup Z_1$ are mapped to Ψ' , and by (*3) we have, similarly as above,

$$\deg_G(\phi(\text{Par}(z)), \Psi') > |W_A| + |Z_2| + |Z_1|.$$

So z can be embedded in $\Psi' \cap N_G(\phi(\text{Par}(z)))$ as planned.

Finally, for $z \in Z_1 \cup Z_2$, denote by T_z the component of $\mathcal{C}_1 \cup \mathcal{C}_2$ that contains z . We use Lemma 6.17 to embed the rest of the rooted tree (T_z, z) . (Note that our parameters work because of (5.1).) Once all rooted trees (T_z, z) , $z \in Z_1 \cup Z_2$ have been processed, we have finished Stage 2 and thus the proof of the lemma. \square

6.5.3 Embedding in Configurations ($\diamond 6$)–($\diamond 10$)

We follow the schemes outlined in Sections 6.1.2, 6.1.3, 6.1.4, and 6.1.5.

Embedding a tree $T_{T_{1,2}} \in \mathbf{trees}(k)$ using Configurations ($\diamond 6$), ($\diamond 7$), ($\diamond 8$) has two parts: first the internal part of $T_{T_{1,2}}$ is embedded, and then this partial embedding is extended to end shrubs of $T_{T_{1,2}}$ as well. Lemma 6.20 (for configurations ($\diamond 6$) and ($\diamond 7$)) and Lemma 6.21 (for configuration ($\diamond 8$)) are used for the former part, and Lemmas 6.22 and 6.23 (depending on whether we have ($\heartsuit 1$) or ($\heartsuit 2$)) for the latter. Lemma 6.24 then puts these two pieces together.

Embedding using Configurations ($\diamond 9$) and ($\diamond 10$) is resolved in Lemmas 6.25 and 6.26, respectively.

Lemma 6.20. *Suppose we are in Setting 5.1 and 5.4, and we have one of the following two configurations:*

- Configuration ($\diamond 6$)($\delta_6, \tilde{\varepsilon}, d', \mu, 1, 0$), or
- Configuration ($\diamond 7$)($\delta_7, \frac{\eta\gamma}{400}, \tilde{\varepsilon}, d', \mu, 1, 0$),

with $10^5 \sqrt{\gamma}(\Omega^*)^2 \leq \delta_6^4 \leq 1$, $10^2 \sqrt{\gamma}(\Omega^*)^3/\Lambda \leq \delta_7^3 < \eta^3 \gamma^3/10^6$, $d' > 10\tilde{\varepsilon} > 0$, and $d'\mu\tau k \geq 4 \cdot 10^3$. Both configurations contain distinguished sets $V_0, V_1 \subseteq \mathfrak{P}_0$ and $V_2, V_3 \subseteq \mathfrak{P}_1$.

6.5 Main embedding lemmas

Suppose that $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ is a (τk) -fine partition of a rooted tree (T, r) of order at most k such that $|W_A \cup W_B| \leq k^{0.1}$. Let T' be the tree induced by all the cut-vertices $W_A \cup W_B$ and all the internal shrubs.

Then there exists an embedding ϕ of T' such that $\phi(W_A) \subseteq V_1$, $\phi(W_B) \subseteq V_0$, and $\phi(T' - (W_A \cup W_B)) \subseteq \mathfrak{P}_1$.

Proof. For simplicity, let us assume that $r \in W_A$. The case when $r \in W_B$ is similar. The (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ induces a (τk) -fine partition in T' . By Lemma 3.8, the tree T' has an ordered skeleton (X_0, X_1, \dots, X_m) where the X_i are either shrubs or knags (X_0 being a knag).

Our strategy is as follows. We sequentially embed the knags and the internal shrubs in the order given by the ordered skeleton. For embedding the knags we use Lemma 6.5 in Preconfiguration (**exp**), and Lemma 6.8 in Preconfiguration (**reg**). For embedding the internal shrubs, we use Lemmas 6.12 and 6.13 if we have Configurations ($\diamond 6$), and ($\diamond 7$), respectively.

Throughout, ϕ denotes the current (partial) embedding of (X_0, X_1, \dots, X_m) . In consecutive steps, we extend ϕ . We define auxiliary sets $D_i \subseteq V(G)$ which will serve for reserving space for the roots of the shrubs X_i . So the set $Z_{<i} := \bigcup_{j < i} (\phi(X_j) \cup D_j)$ contains what is already used and what should (mainly) be avoided.

Let $W_{A,i} := W_A \cap V(X_i)$, and $W_{B,i} := W_B \cap V(X_i)$. For each $y \in W_{A,j}$ with $j \leq i$ let

$$S_y := (V_2 \cap N_G(\phi(y))) \setminus Z_{<i},$$

except if the latter set has size $> k$, in that case we choose a subset of size k . This is a target set for the roots of shrubs adjacent to y .

Also, in the case X_i is a shrub, we write r_i for its root, and f_i for the only other vertex neighbouring $W_A \cup W_B$. Note that f_i is a fruit of (X_i, r_i) .

The value $h = 6$ or $h = 7$ indicates whether we have configuration ($\diamond 6$) or ($\diamond 7$). Define

$$F_i := \text{shadow}_{G-\Psi} \left(Z_{<i}, \frac{\delta_h k}{4} \right) \cup Z_{<i}. \quad (6.27)$$

Define $U_i := F_i$ if we have Preconfiguration (**exp**) (note that in that case we have Configuration ($\diamond 6$)). To define U_i in case of Preconfiguration (**reg**) we make use of the super-regular pairs $(Q_0^{(j)}, Q_1^{(j)})$ ($j \in \mathcal{Y}$). Set

$$U_i := F_i \cup \bigcup \left\{ Q_1^{(j)} : j \in \mathcal{Y}, |Q_1^{(j)} \cap F_i| \geq \frac{|Q_1^{(j)}|}{2} \right\}. \quad (6.28)$$

In either case, we have $|U_i| \leq 2|F_i|$.

Finally, set

$$W_i := \text{shadow}_{G-\Psi} \left(U_i, \frac{\delta_h k}{2} \right) \cup Z_{<i}. \quad (6.29)$$

We will now show how to embed successively all X_i . At each step i , our embedding ϕ will have the following properties:

- (a) $\phi(W_{A,i}) \subseteq V_1 \setminus F_i$ and $\phi(W_{B,i}) \subseteq V_0$,
- (b) for each $y \in W_{A,j}$ with $j \leq i$ we have $|S_y \cap \phi(X_i)| \leq |S_y \cap D_i| + k^{3/4}$,

- (c) $|Z_{<i+1}| \leq 2k$,
- (d) $D_i \subseteq \mathfrak{P}_1 \setminus (\phi(X_i) \cup Z_{<i})$,
- (e) $\phi(X_i - r_i)$ is disjoint from $\bigcup_{j < i} \cup D_j$,
- (f) $\phi(f_i) \in V_2 \setminus W_i$ if X_i is a shrub,
- (g) $\phi(X_i) \subseteq \mathfrak{P}_1$ if X_i is a shrub.

(We remark that since r_i is not defined for knags X_i , condition (e) means that $\phi(X_i)$ is disjoint from $\bigcup_{j < i} \cup D_j$ for knags X_i .)

It is clear that the first together with the last condition ensures that in step m we have found the desired embedding for T' .

Before we show how to embed each X_i fulfilling the properties above, let us quickly calculate a useful bound. By Fact 4.12 and (c), we have that $|F_i| \leq \frac{9\Omega^*}{\delta_h} k$ for all $i \leq m$. Thus, using $|U_i| \leq 2|F_i|$ and again Fact 4.12 and (c), this shows

$$|W_i| \leq \frac{38(\Omega^*)^2}{\delta_h^2} k. \quad (6.30)$$

Now suppose we are at step i with $0 \leq i \leq m$. That is, we have already embedded all X_j with $j < i$, and are about to embed X_i .

First assume that X_i is a knag. Note that if $i \neq 0$, then there is exactly one fruit f_ℓ with $\ell < i$ which neighbours X_i . Set $N_i := N_G(\phi(f_\ell))$ in this case, and let $N_i := V(G)$ for $i = 0$. We distinguish between the two preconfigurations we might be in.

Suppose first we are in Preconfiguration (**exp**). Recall that then we are in Configuration ($\diamond 6$).

We use Lemma 6.5 to embed the single tree X_i with the following setting: $\ell_{L6.5} := 1$, $V_{1,L6.5} := V_1$, $V_{2,L6.5} := V_0$, $U_{L6.5}^* := (N_i \cap V_1) \setminus U_i = (N_i \cap V_1) \setminus F_i$, $U_{L6.5} := U_i = F_i$, $Q_{L6.5} := \frac{18\Omega^*}{\delta_6}$, $\zeta_{L6.5} := \delta_6$, and $\gamma_{L6.5} := \gamma$. Note that $U_{L6.5}^*$ is large enough by (f) for ℓ and by (5.37) and (5.41), respectively. Lemma 6.5 gives an embedding of the tree X_i such that $\phi(V_{\text{even}}(X_i)) \subseteq V_1 \setminus F_i$ and $\phi(V_{\text{odd}}(X_i)) \subseteq V_0 \setminus F_i$, which maps the root of X_i to the neighbourhood of its parent's image. Note that this ensures (a) and (e) for step i , and setting $D_i := \emptyset$ we also ensure (c) and (d). Property (b) holds since $V_2 \cap \phi(X_i) = \emptyset$. Since X_i is a knag, (f) and (g) are empty.

Suppose now we are in Preconfiguration (**reg**). Then let $j \in \mathcal{Y}$ be such that $(N_i \cap Q_1^{(j)}) \setminus U_i \neq \emptyset$. Such an index j exists by (f) for ℓ and by (5.37) and (5.41), respectively, if $i \neq 0$, and trivially if $i = 0$. We shall use Lemma 6.8 to embed X_i in $(Q_0^{(j)}, Q_1^{(j)})$. More precisely, we use Lemma 6.8 with $A_{L6.8} := Q_1^{(j)}$, $B_{L6.8} := Q_0^{(j)}$, $\varepsilon_{L6.8} := \tilde{\varepsilon}$, $d_{L6.8} := d'$, $\ell_{L6.8} := \mu k$, $U_A := U_i \cap A$, $U_B := \phi(W_{B,<i}) \cap B$ (then $|U_A| \leq |A|/2$ by the definition of U_i and the choice of j).

Lemma 6.8 yields a $(V_{\text{even}}(X_i) \hookrightarrow V_1 \setminus F_i, V_{\text{odd}}(X_i) \hookrightarrow V_0)$ -embedding of X_i , which maps the root of X_i to the neighbourhood of its parent's image. Setting $D_i := \emptyset$, we have (a)–(g).

So let us now assume that X_i is a shrub. The parent y of the root r_i of X_i lies in $W_{A,\ell}$ for some $\ell < i$. By (a) for ℓ , we mapped y to a vertex $\phi(y) \in V_1 \setminus F_\ell$. As $\deg_G(\phi(y), V_2) \geq \delta_h k$ (by (5.36) and (5.40), respectively), and since $\phi(y) \notin F_\ell$, we have

$$|S_y| \geq \frac{3\delta_h k}{4}. \quad (6.31)$$

6.5 Main embedding lemmas

Using (b) for all j with $\ell \leq j < i$, and using that the sets D_j are pairwise disjoint by (d), we see that

$$|S_y \cap \phi(X_0 \cup \dots \cup X_{i-1})| = |S_y \cap \phi(X_\ell \cup \dots \cup X_{i-1})| \leq |S_y \cap \bigcup_{\ell \leq j < i} D_j| + m \cdot k^{3/4} \leq |S_y \cap \bigcup_{0 \leq j < i} D_j| + m \cdot k^{3/4}.$$

Therefore, and as by (d) and (e), the sets $\phi(X_0 \cup \dots \cup X_{i-1})$ and $\bigcup_{0 \leq j < i} D_j$ are disjoint except for the at most $m \leq |W_A \cup W_B| \leq k^{0.1}$ roots r_j of shrubs X_j , and since $k \gg 1$, we have

$$|S_y| \geq |S_y \cap \phi(X_0 \cup \dots \cup X_{i-1})| + |S_y \cap \bigcup_{0 \leq j < i} D_j| - m \geq 2|S_y \cap \phi(X_0 \cup \dots \cup X_{i-1})| - k^{0.9}.$$

Thus,

$$|S_y \setminus \phi(X_0 \cup \dots \cup X_{i-1})| \geq \frac{|S_y| - k^{0.9}}{2} \stackrel{(6.31)}{\geq} \frac{3\delta_h k}{8} - \frac{k^{0.9}}{2} > \frac{\delta_h k}{3}.$$

So for $U^* := S_y \setminus \phi(X_0 \cup \dots \cup X_{i-1})$ we have that $|U^*| \geq \frac{\delta_h k}{3}$. If we have Configuration $(\diamond 6)$ or $(\diamond 7)$ we use Lemma 6.12 or 6.13, respectively, with input $U_{L6.12-6.13} := W_i$, $U_{L6.12-6.13}^* := U^*$, $L_{L6.12-6.13} := |W_{A,i}|$, $\gamma_{L6.12-6.13} := \gamma$, the family $\{P_t\}_{L6.12-6.13} := \{S_y\}_{y \in W_{A,i}, j < i}$, and the rooted tree (X_i, r_i) with fruit f_i . Further, for Configuration $(\diamond 6)$, set $\delta_{L6.12} := \delta_6$, $V_{2,L6.12} := V_2$ and $V_{3,L6.12} := V_3$ and for Configuration $(\diamond 7)$, set $\delta_{L6.13} := \delta_7$, $\ell_{L6.13} := 1$, $Y_{1,L6.13} := V_2$ and $Y_{2,L6.13} := V_3$. The output of Lemma 6.12 or 6.13, respectively, is the extension of our embedding ϕ to X_i , and a set $D_i := C_{L6.12-6.13} \subseteq (V_2 \cup V_3) \setminus (W_i \cup \phi(X_i))$ for which properties (a) (which is empty) and properties (b)–(g) hold. \square

Lemma 6.21. *Suppose we are in Setting 5.1 and 5.4 and suppose further we have Configuration $(\diamond 8)$ $(\delta, \frac{\eta\gamma}{400}, \varepsilon_1, \varepsilon_2, d_1, d_2, \mu_1, \mu_2, h_1, 0)$, with $2 \cdot 10^5 (\Omega^*)^6 / \Lambda \leq \delta^6$, $\delta < \gamma^2 \eta^4 / (10^{16} (\Omega^*)^2)$, $d_2 > 10\varepsilon_2 > 0$, $d_2 \mu_2 \tau k \geq 4 \cdot 10^3$, and $\max\{\varepsilon_1, \tau / \mu_1\} \leq \eta^2 \gamma^2 d_1 / (10^{10} (\Omega^*)^3)$. Recall that we have distinguished sets V_0, \dots, V_4 and a semiregular matching \mathcal{N} .*

Let $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ be a (τk) -fine partition of a rooted tree (T, r) of order at most k . Let T' be the tree induced by all the cut-vertices $W_A \cup W_B$ and all the internal shrubs. Suppose that

$$v(T') < h_1 - \frac{\eta^2 k}{10^5}. \tag{6.32}$$

Then there exists an embedding ϕ of T' such that $\phi(W_A) \subseteq V_1$, $\phi(W_B) \subseteq V_0$, and $\phi(T') \subseteq \mathfrak{P}_0 \cup \mathfrak{P}_1$.

Proof. We assume that $r \in W_A$. The case when $r \in W_B$ is similar.

Let \mathcal{K} be the set of all knags of the (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of T . For each such knag $K \in \mathcal{K}$ set $Y_K := K \cup \text{Ch}_{T'}(K)$. We call the subgraphs Y_K *extended knags*. Set $\mathcal{Y} := \{Y_K : K \in \mathcal{K}\}$ and $W_C := V(\bigcup \mathcal{Y} \setminus \bigcup \mathcal{K})$. Since $W_C \subseteq V(T')$, we clearly have that $|W_C| \leq |W_A \cup W_B|$.

Note that the forest $T' - \bigcup \mathcal{Y}$ consists of the set \mathcal{P} of peripheral subshrubs of internal shrubs of $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$, and the set \mathcal{S} of principal subshrubs of internal shrubs of $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$. It is not difficult to observe that there is a sequence (X_0, X_1, \dots, X_m) such that $X_i = (M_i, Y_i, P_i)$ such that $M_i \in \mathcal{S}$ and $P_i \subseteq \mathcal{P}$ for each $i \leq m$, and such that the following holds.

- (I) $M_0 = \emptyset$ and Y_0 contains r .

- (II) \mathcal{P}_i are exactly those peripheral subshrubs whose parents lie in Y_i .
- (III) The parent f_i of Y_i lies in M_i (unless $i = 0$).
- (IV) The parent r_i of M_i lies in some Y_j with $j < i$ (unless $i = 0$),
- (V) $\bigcup_{i \leq m} V(M_i \cup Y_i \cup \bigcup \mathcal{P}_i) = V(T')$.

See Figure 6.7 for an illustration.




Figure 6.7: An example of a sequence $(X_0, X_1, X_2, X_3, \dots)$ in Lemma 6.21.

We now successively embed the elements of X_i , except possibly for a part of the subshrubs in \mathcal{P}_i . The omitted peripheral subshrubs will be embedded at the very end, after having completed the inductive procedure we are about to describe now.

We shall make use of the following lemmas: Lemma 6.8 (for embedding knags), Lemmas 6.9 and 6.6 (for embedding peripheral subshrubs in \mathcal{N}), Lemma 6.13 (for embedding principal subshrubs in $V_3 \cup V_4$).

Throughout, ϕ denotes the current (partial) embedding of T' . In each step i we embed $M_i \cup Y_i$ and a subset of \mathcal{P}_i , and denote by $\phi(X_i)$ the image of these sets (as far as it is defined). We also define an auxiliary set $D_i \subseteq V(G)$ which will serve to ensure there is enough space for the roots of the subshrubs M_ℓ with $\ell > i$. Set

$$Z_{<i} := \bigcup_{j < i} (\phi(X_j) \cup D_j).$$

Our plan for embedding the various parts of X_i is depicted in Figure 6.8, which is a refined version of Figure 6.3.

Let $W_{O,i} := W_O \cap V(Y_i)$ for $O = A, B, C$. For each $y \in W_{C,i}$ let

$$S_y := (V_3 \cap N_G(\phi(y))) \setminus Z_{<i},$$

except if this set has size more than k , in which case we choose any subset of size k . Similar as in the preceding lemma, this is a target set for the roots of the principal subshrub adjacent to y .




Figure 6.8: Embedding a part of the internal tree in Lemma 6.21.

Fix a matching involution \mathfrak{d} for \mathcal{N} , and for $\ell = 1, 2$ define

$$F_i^{(\ell)} := Z_{<i} \cup \text{shadow}_{G-\Psi}^{(\ell)} \left(\text{ghost}_{\mathfrak{d}}(Z_{<i}), \frac{\delta k}{8} \right). \quad (6.33)$$

We use the super-regular pairs $(Q_0^{(j)}, Q_1^{(j)})$ ($j \in \mathcal{Y}$) to define

$$U_i := F_i^{(2)} \cup \bigcup \left\{ Q_1^{(j)} : j \in \mathcal{Y}, |Q_1^{(j)} \cap F_i^{(2)}| \geq \frac{|Q_1^{(j)}|}{2} \right\}. \quad (6.34)$$

We have

$$|U_i| \leq 2|F_i^{(2)}|. \quad (6.35)$$

Finally, for $\ell = 1, 2$ set

$$W_i^{(\ell)} := \text{shadow}_{G-\Psi}^{(\ell)} \left(U_i, \frac{\delta k}{2} \right). \quad (6.36)$$

We will now show how to define successively our embedding. At each step i , the embedding ϕ will be defined for $M_i \cup Y_i$ and a subset of \mathcal{P}_i , and it will have the following properties:

- (a) $\phi(W_{A,i}) \subseteq V_1 \setminus F_i^{(2)}$ and $\phi(W_{B,i}) \subseteq V_0$,
- (b) $\phi(W_{C,i}) \subseteq V_2 \setminus F_i^{(1)}$,
- (c) $\phi(f_i) \in V_2 \setminus (F_i^{(1)} \cup W_i^{(1)})$,
- (d) for each $y \in W_{C,j}$ with $j \leq i$ we have $|S_y \cap \phi(X_i)| \leq |S_y \cap D_i| + 2k^{3/4}$,
- (e) $|Z_{<i+1}| \leq 2k$,
- (f) $D_i \subseteq V_3 \setminus (\phi(X_i) \cup Z_{<i})$,

- (g) $\phi(X_i \setminus (V(M_i) \cap \text{Ch}(W_C)))$ is disjoint from $\bigcup_{j < i} D_j$,¹¹
- (h) $\phi(X_i) \subseteq \mathfrak{P}_1 \cup \phi(Y_i \cup f_i)$,
- (i) if $P \in \mathcal{P}_i$ is not embedded in step i then for its parent $w \in W_C$ we have that $\deg_{G_D}(\phi(w), V_3) \geq h_1 - |\phi(X_i) \cap V(\mathcal{N})| - \frac{\eta^2 k}{10^6}$.

Note that for (h), since f_0 is not defined, we assume $\phi(f_0) = \emptyset$.

Before we go on let us remark that (h) together with (f) implies that at each step i we have

$$|Z_{<i} \cap \mathfrak{P}_0| \leq 3 \cdot (|W_A| + |W_B|) \stackrel{\text{D3.3(e)}}{\leq} \frac{2016}{\tau} < \frac{\delta k}{8}. \quad (6.37)$$

Also note that by Fact 4.12 and by (e), we have

$$|F_i^{(2)}| \leq \frac{65(\Omega^*)^2}{\delta^2} k, \quad (6.38)$$

and

$$|W_i^{(2)}| \leq \frac{520(\Omega^*)^4}{\delta^4} k. \quad (6.39)$$

By (b) and by (5.46) we have that $|S_y| \geq \frac{7\delta k}{8}$. Now, using (d), (f) and (g), we can calculate similarly as in the previous lemma that at each step i we have

$$|S_y \setminus \bigcup_{\ell < i} \phi(X_\ell)| \geq \frac{3\delta k}{8}. \quad (6.40)$$

Now assume we are at step i of the inductive procedure, that is, we have already dealt with X_0, \dots, X_{i-1} and wish to embed (parts of) X_i .

We start with embedding M_i , except if $i = 0$, in that case we go directly to embedding Y_0 . We shall embed M_i in $V_3 \cup V_4$, except for the fruit f_i , which will be mapped to V_2 . The embedding has three stages. First we embed $M_i - M_i(\uparrow f_i)$, then we embed f_i , and finally we embed the forest $M_i(\uparrow f_i) - f_i$. The embedding of $M_i - M_i(\uparrow f_i)$ is an application of Lemma 6.13 analogous to the case of Configuration ($\diamond 7$) in the previous Lemma 6.20. That is, set $Y_{1, \text{L6.13}} := V_3$, $Y_{2, \text{L6.13}} := V_4$, let

$$U_{\text{L6.13}}^* := S_{r_i} \setminus \bigcup_{\ell < i} \phi(X_\ell),$$

where r_i lies in W_C by (IV), and

$$U_{\text{L6.13}} := F_i^{(2)} \cup W_i^{(2)}.$$

Note that

$$|U_{\text{L6.13}}| \leq \frac{10^3(\Omega^*)^4}{\delta^4} k \leq \frac{\delta \Lambda}{2\Omega^*} k,$$

and by (6.40) (which we use for $i - 1$), also

$$|U_{\text{L6.13}}^*| \geq \frac{3\delta k}{8}.$$

¹¹Note that $V(M_i) \cap \text{Ch}(W_C)$ contains a single vertex, the root of M_i .

6.5 Main embedding lemmas

The family $\{P_1, \dots, P_L\}_{L6.13}$ is $\{S_y\}_{y \in \bigcup_{j < i} W_{C,j}}$. There is only one tree to be embedded, namely $M_i - M_i(\uparrow f_i)$. It is not difficult to check that all the conditions of Lemma 6.13 are fulfilled. Lemma 6.13 gives an embedding of $M_i - M_i(\uparrow f_i)$ in $V_3 \cup V_4 \subseteq \mathfrak{P}_1$ with the property that $\text{Par}(f_i)$ is mapped to $V_3 \setminus (F_i^{(2)} \cup W_i^{(2)})$. The lemma further gives a set $D' := C_{L6.13}$ of size $v(M_i - M_i(\uparrow f_i))$ such that

$$|S_y \cap \phi(M_i - M_i(\uparrow f_i))| \leq |S_y \cap D'| + k^{0.75}$$

for each $y \in \bigcup_{j < i} W_{C,j}$.

Using the degree condition (5.47) we can embed f_i to

$$V_2 \setminus (F_i^{(1)} \cup W_i^{(1)})$$

(recall that (6.37) asserts that only very little space in V_2 is occupied). This ensures (c) for i .

To embed $M_i(\uparrow f_i) - f_i$ we use again Lemma 6.13. The parameters are this time $Y_{1,L6.13} := V_3$, $Y_{2,L6.13} := V_4$,

$$\begin{aligned} U_{L6.13}^* &:= (\text{N}_G(\phi(f_i)) \cap V_3) \setminus (Z_{<i} \cup \phi(M_i - M_i(\uparrow f_i))), \text{ and} \\ U_{L6.13} &:= Z_{<i} \cup \phi(M_i - M_i(\uparrow f_i)) \cup D'. \end{aligned}$$

Note that $|U_{L6.13}^*| \geq \frac{\delta k}{4}$ by (5.46), by the fact that $\phi(f_i) \notin W_i^{(1)}$, and as $v(T_i) + i < \delta k/8$. The family $\{P_1, \dots, P_L\}_{L6.13}$ is $\{S_y\}_{y \in \bigcup_{j < i} W_{C,j}}$. The trees to be embedded are the components of $M_i(\uparrow f_i) - f_i$ rooted at the children of f_i . All the conditions of Lemma 6.13 are fulfilled. The lemma provides an embedding in $V_3 \cup V_4 \subseteq \mathfrak{P}_1$. It further gives a set $D'' := C_{L6.13}$ of size $v(M_i(\uparrow f_i)) - 1$ such that

$$|S_y \cap \phi(M_i(\uparrow f_i) - f_i)| \leq |S_y \cap D''| + k^{0.75}$$

for each $y \in \bigcup_{j < i} W_{C,j}$. Then $D_i := V_3 \cap (D' \cup D'')$ is such that for each $y \in \bigcup_{j < i} W_{C,j}$,

$$|S_y \cap \phi(M_i)| \leq |S_y \cap D_i| + 2k^{0.75}, \quad (6.41)$$

as $S_y \subseteq V_3$ and $\phi(f_i) \notin V_3$. Note that this choice of D_i also ensures (e) for i , and we have by the choices of $U_{L6.13}^*$ and $U_{L6.13}$ in both applications of Lemma 6.13 that

$$D_i \subseteq V_3 \setminus (\phi(M_i) \cup Z_{<i}) \quad \text{and} \quad \phi(X_i \setminus (V(M_i) \cap \text{Ch}(W_C))) \cap \bigcup_{j < i} D_j = \emptyset. \quad (6.42)$$

We now turn to embedding Y_i . Our plan is to use first Lemma 6.8 to embed $Y_i \setminus W_C$ in $(Q_0^{(j)}, Q_1^{(j)})$ for an appropriate index j . After that, we shall show how to embed $W_{C,i}$.

If $i = 0$ then take an arbitrary $j \in \mathcal{Y}$. Otherwise note that by (III), the parent f_i of the root of Y_i lies in M_i . Note that f_i is a fruit in M_i . Let $j \in \mathcal{Y}$ be such that $(\text{N}_G(\phi(f_i)) \cap Q_1^{(j)}) \setminus U_i \neq \emptyset$. Such an index j exists by (5.45) and the fact that $\phi(f_i) \notin W_i^{(1)}$ by (c) for i .

We use Lemma 6.8 with $A_{L6.8} := Q_1^{(j)}$, $B_{L6.8} := Q_0^{(j)}$, $\varepsilon_{L6.8} := \varepsilon_2$, $d_{L6.8} := d_2$, $\ell_{L6.8} := \mu_2 k$, $U_A := U_i \cap A_{L6.8}$, $U_B := Z_{<i} \cap B_{L6.8}$. By the choice of j and the definition of U_i , we find that U_A is small enough, and using (6.37) we see that U_B is also small enough. Lemma 6.8 yields a $(V_{\text{even}}(Y_i - W_C) \hookrightarrow V_1 \setminus F_i^{(2)}, V_{\text{odd}}(Y_i - W_C) \hookrightarrow V_0)$ -embedding of $Y_i - W_C$. We clearly see condition (a) satisfied for i .

6.5 Main embedding lemmas

We now embed successively the vertices of the set $W_{C,i} = \{w_\ell : \ell = 1, \dots, |W_{C,i}|\}$. By the definition of the set W_C , we know that the parent x of w_ℓ lies in $W_{A,i}$. Combining (5.44) with the fact that $\phi(x) \in V_1 \setminus F_i^{(2)}$ by (a) for i , we have that

$$\left| N_G \left(\phi(x), V_2 \setminus (F_i^{(1)} \setminus Z_{<i}) \right) \right| \geq \frac{7\delta k}{8}.$$

Thus by (6.37) and since $V_2 \subseteq \mathfrak{P}_0$, we can accommodate w_ℓ in $V_2 \setminus F_i^{(1)}$. This is as desired for (b) in step i .

We now turn to \mathcal{P}_i . We will embed a subset of these peripheral subshrubs in \mathcal{N} . This procedure is divided into two stages. First we shall aim to embed as many subshrubs as possible in \mathcal{N} in a balanced way, with the help of Lemma 6.9. When it is no longer possible to embed any subshrub in a balanced way in \mathcal{N} , we embed in \mathcal{N} as many of the leftover subshrubs as possible, in an unbalanced way. For this part of the embedding we use Lemma 6.6.

By (II) all the parents of the subshrubs in \mathcal{P}_i lie in $W_{C,i}$. For $w_\ell \in W_{C,i}$, let $\mathcal{P}_{i,\ell}$ denote the set of all subshrubs in \mathcal{P}_i adjacent to w_ℓ . In the first stage, we shall embed, successively for $j = 1, \dots, |W_{C,i}|$, either all or none of $\mathcal{P}_{i,j}$ in a balanced way in \mathcal{N} . Assume inductively that

$$\phi \left(\bigcup_{p < j} \mathcal{P}_{i,p} \right) \text{ is } (\tau k)\text{-balanced with respect to } \mathcal{N}. \quad (6.43)$$

Construct a semiregular matching \mathcal{N}_j absorbed by \mathcal{N} as follows. Let $\mathcal{N}_j := \{(X'_1, X'_2) : (X_1, X_2) \in \mathcal{N}\}$, where for $(X_1, X_2) \in \mathcal{N}$ we define (X'_1, X'_2) as the maximal balanced unoccupied subpair seen from $\phi(w_j)$, i.e., for $b = 1, 2$, we take

$$X'_b \subseteq (X_b \cap N_{G_{\text{reg}}}(\phi(w_j))) \setminus \left(\phi \left(\bigcup_{p < j} \mathcal{P}_{i,p} \right) \cup \bigcup_{\ell < i} \phi(X_\ell) \right)$$

maximal subject to $|X'_1| = |X'_2|$. If $|V(\mathcal{N}_j)| \geq \frac{\eta^2 k}{10^7 \Omega^*}$ then we shall embed $\mathcal{P}_{i,j}$, otherwise we do not embed $\mathcal{P}_{i,j}$ in this step. So assume we decided to embed $\mathcal{P}_{i,j}$. Recall that the total order of the subshrubs in this set is at most τk . Using the same argument as for Claim 6.18.1 we have

$$\left| \bigcup \{X \cup Y : (X, Y) \in \mathcal{N}, \deg_{G_D}(\phi(w_j), X \cup Y) > 0\} \right| \leq \frac{4(\Omega^*)^2}{\gamma^2} k.$$

Thus, there exists a subpair $(X'_1, X'_2) \in \mathcal{N}_j$ of some $(X_1, X_2) \in \mathcal{N}$ with

$$\frac{|X'_1|}{|X_1|} \geq \frac{\frac{\eta^2}{10^7 \Omega^*} k}{\frac{4(\Omega^*)^2}{\gamma^2} k} \geq \frac{\gamma^2 \eta^2}{10^8 (\Omega^*)^3}. \quad (6.44)$$

In particular, (X'_1, X'_2) forms a $\frac{2 \cdot 10^8 \varepsilon_1 (\Omega^*)^3}{\gamma^2 \eta^2}$ -regular pair of density at least $d_1/2$ by Fact 2.1. We use Lemma 6.9 to embed $\mathcal{P}_{i,j}$ in $\mathcal{M}_{L6.9} := \{(X'_1, X'_2)\}$. The family $\{f_{CD}\}_{L6.9}$ comprises of a single number $f_{(X'_1, X'_2)}$ which is the discrepancy of $\bigcup_{p < j} \phi(\mathcal{P}_{i,p})$ with respect to (X_1, X_2) . This guarantees that (6.43) is preserved. This finishes the j -th step. We repeat this step until $j = |W_{C,i}|$, then we go to the next stage.

6.5 Main embedding lemmas

Denote by \mathcal{Q}_i the set of all $P \in \mathcal{P}_i$ that have not been embedded in the first stage. Note that for each $Q \in \mathcal{Q}_i$, with $Q \in \mathcal{P}_{i,j}$, say, and for each $(X_1, X_2) \in \mathcal{N}$ there is a $b_{(X_1, X_2)} \in \{1, 2\}$ such that for

$$O_j := \bigcup_{(X_1, X_2) \in \mathcal{N}} \left(X_{b_{(X_1, X_2)}} \cap N_{G_{\text{reg}}}(\phi(w_j)) \right) \setminus \left(\phi\left(\bigcup_{p < j} \mathcal{P}_{i,p}\right) \cup \bigcup_{\ell < i} \phi(X_\ell) \right)$$

we have that

$$|O_j| < \frac{\eta^2 k}{10^7 \Omega^*}. \quad (6.45)$$

The fact that O_j is small implies that there an \mathcal{N} -cover such that the G_{reg} -neighborhood of w_j restricted to this cover is essentially exhausted by the image of T' .

In the second stage, we shall embed some of the peripheral subshrubs of \mathcal{Q}_i . They will be mapped in an unbalanced way to \mathcal{N} . We will do this in steps $j = 1, \dots, |W_{C,i}|$, and denote by \mathcal{R}_j the set of those $\mathcal{P} \subseteq \mathcal{Q}_i$ embedded until step j . At step j , we decide to embed $\mathcal{P}_{i,j}$ if $\mathcal{P}_{i,j} \subseteq \mathcal{Q}_i$ and

$$\deg_{G_{\text{reg}}}(\phi(w_j), V(\mathcal{N}) \setminus \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i)) - |\bigcup \mathcal{R}_{j-1}| \geq \frac{\eta^2 k}{10^6}. \quad (6.46)$$

Let

$$\tilde{\mathcal{N}} := \left\{ (X, Y) \in \mathcal{N} : |(X \cup Y) \cap Z_{<i}| < \frac{\gamma^2 \eta^2}{10^9 (\Omega^*)^2} |X| \right\}.$$

As by (b) we know that w_j was embedded in $V_2 \setminus F_i^{(1)}$, we have

$$\deg_{G_{\text{reg}}}(\phi(w_j), V(\mathcal{N} \setminus \tilde{\mathcal{N}})) \leq \frac{2 \cdot 10^9 (\Omega^*)^2}{\gamma^2 \eta^2} \cdot \frac{\delta k}{8} \leq \frac{\eta^2}{10^7} k. \quad (6.47)$$

Using (6.45), (6.46) and (6.47), similar calculations as in (6.44) show the existence of a pair $(X, Y) \in \tilde{\mathcal{N}}$ with

$$\deg_{G_{\text{reg}}}(\phi(w_j), (X \cup Y) \setminus (O_j \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i))) - |(X \cup Y) \cap \phi(\bigcup \mathcal{R}_{j-1})| \geq \frac{\gamma^2 \eta^2}{10^8 (\Omega^*)^2} |X \cup Y|.$$

Then by the definition of $\tilde{\mathcal{N}}$, and setting $Z_{<i}^+ := \text{ghost}_0(Z_{<i})$ we get that

$$\begin{aligned} & \deg_{G_{\text{reg}}}(\phi(w_j), (X \cup Y) \setminus (Z_{<i}^+ \cup O_j \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i))) - |(X \cup Y) \cap \phi(\bigcup \mathcal{R}_{j-1})| \\ & \geq \frac{\gamma^2 \eta^2}{10^9 (\Omega^*)^2} |X \cup Y|. \end{aligned}$$

By the definition of O_j , all of the degree counted here goes to one side of the matching edge (X, Y) , say to X . So

$$\deg_{G_{\text{reg}}}(\phi(w_j), X \setminus (Z_{<i}^+ \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i \cup \bigcup \mathcal{R}_{j-1}))) - |Y \cap \phi(\bigcup \mathcal{R}_{j-1})| \geq \frac{\gamma^2 \eta^2}{10^9 (\Omega^*)^2} |X| \quad (6.48)$$

$$\geq 12 \frac{\varepsilon_1}{d_1} |X| + \tau k. \quad (6.49)$$

We claim that furthermore,

$$|Y \setminus (Z_{<i}^+ \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i \cup \bigcup \mathcal{R}_{j-1}))| \geq \frac{\gamma^2 \eta^2}{10^{10}(\Omega^*)^2} |Y| \geq 12 \frac{\varepsilon_1}{d_1} |Y| + \tau k. \quad (6.50)$$

Indeed, otherwise we get by (6.48) that

$$|X \setminus (Z_{<i}^+ \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i))| > |Y \setminus (Z_{<i}^+ \cup \phi(\bigcup \mathcal{P}_i \setminus \mathcal{Q}_i))| + \frac{\gamma^2 \eta^2}{10^{10}(\Omega^*)^2} |X|,$$

which is impossible by (6.43) and since $|X| \geq \mu_1 k$.

Hence, by (6.49) and (6.50), we can embed $\mathcal{P}_{i,j}$ into the unoccupied part (X, Y) using Lemma 6.6 repeatedly.¹²

Note that if some $\mathcal{P}_{i,j}$ has not been embedded in either of the two stages, then the vertex w_j must have a somewhat insufficient degree in \mathcal{N} . More precisely, employing (6.46) we see that $\deg_{G_{\text{reg}}}(\phi(w_j), V(\mathcal{N})) - |\phi(X_i) \cap V(\mathcal{N})| < \frac{\eta^2 k}{10^6}$. Combined with (5.50), we find that

$$\deg_{G_{\mathcal{D}}}(\phi(w_j), V_3) \geq h_1 - |\phi(X_i) \cap V(\mathcal{N})| - \frac{\eta^2 k}{10^6},$$

in other words, (i) holds for i .

This finishes step i of the embedding procedure. Recall that the sets V_3 and $V(\mathcal{N})$ are disjoint. Hence, by (a) and (b), the principal subshrubs are the only parts of T' that were embedded in V_3 (and possibly elsewhere). Thus, using (6.42), we see that (f), (g) and (h) are satisfied for i . Also, by (6.41), (d) holds for i .

After having completed the inductive procedure, we still have to embed some peripheral subshrubs. Let us take sequentially those $P \in \mathcal{P}$ which were not embedded. Say w is the parent of P . By (i) we have

$$\deg_{G_{\mathcal{D}}}(\phi(w), V_3 \setminus \text{Im}(\phi)) \geq h_1 - |\text{Im}(\phi) \cap V(\mathcal{N})| - |\text{Im}(\phi) \cap V_3| - \frac{\eta^2 k}{10^6} \stackrel{(6.32)}{\geq} \frac{\eta^2 k}{10^6}.$$

An application of Lemma 6.13 in which $Y_{1,L6.13} := V_3$, $Y_{2,L6.13} := V_4$, $U_{L6.13} := \text{Im}(\phi)$, $U_{L6.13}^* := N_{G_{\mathcal{D}}}(\phi(w)) \cap V_3 \setminus \text{Im}(\phi)$, and $\{P_1, \dots, P_L\}_{L6.13} := \emptyset$ gives an embedding of P in $V_3 \cup V_4 \subseteq \mathfrak{P}_1$.

By conditions (a), (b), (c) and (h) we have thus found the desired embedding for T' . \square

Lemma 6.22. *Suppose we are in Setting 5.1 and 5.4, and that the sets V_0 and V_1 witness Pre-configuration $(\heartsuit 1)(2\eta^3 k/10^3, h)$. Suppose that $U \subseteq \mathfrak{P}_0 \cup \mathfrak{P}_1$. Suppose that $\{x_j\}_{j=1}^\ell \subseteq V_0$ and $\{y_j\}_{j=1}^{\ell'} \subseteq V_1$ are sets of mutually distinct vertices. Let $\{(T_j, r_j)\}_{j=1}^\ell$ and $\{(T'_j, r'_j)\}_{j=1}^{\ell'}$ be families of rooted trees such that each component of $T_j - r_j$ and of $T'_j - r'_j$ has order at most τk .*

If

$$\sum_j v(T_j) \leq \frac{h}{2} - \frac{\eta^2 k}{1000}, \quad (6.51)$$

$$\sum_j v(T_j) + \sum_j v(T'_j) \leq h - \frac{\eta^2 k}{1000}, \text{ and} \quad (6.52)$$

$$|U| + \sum_j v(T_j) + \sum_j v(T'_j) \leq k, \quad (6.53)$$

¹²Recall that the total order of $\mathcal{P}_{i,j}$ is at most τk .

6.5 Main embedding lemmas

then there exist $(r_j \hookrightarrow x_j, V(T_j) \setminus \{r_j\} \hookrightarrow V(G) \setminus U)$ -embeddings of T_j and $(r'_j \hookrightarrow y_j, V(T'_j) \setminus \{r'_j\} \hookrightarrow V(G) \setminus U)$ -embeddings of T'_j in G , all mutually disjoint.

Proof. The embedding has three stages. In Stage I we embed some components of $T_j - r_j$ (for all $j = 1, \dots, \ell$) in the parts of $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edges which are “seen in a balanced way from x_j ”. In Stage II we embed the remaining components of $T_j - r_j$. Last, in Stage III we embed all the components $T'_j - r'_j$ (for all $j = 1, \dots, \ell'$).

Let us first give a bound on the total size of $(\mathcal{M}_A \cup \mathcal{M}_B)$ -vertices $C \in \mathcal{V}(\mathcal{M}_A \cup \mathcal{M}_B)$, $C \subseteq \bigcup \mathbf{V}$ seen from a given vertex via edges of $G_{\mathcal{D}}$. This bound will be used repeatedly.

Claim 6.22.1. Let $v \in V(G)$. Then for $\mathcal{U} := \{C \in \mathcal{V}(\mathcal{M}_A \cup \mathcal{M}_B) : C \subseteq \bigcup \mathbf{V}, \deg_{G_{\mathcal{D}}}(x, C) > 0\}$ we have

$$|\bigcup \mathcal{U}| \leq \frac{2(\Omega^*)^2 k}{\gamma^2}, \text{ and} \quad (6.54)$$

$$|\mathcal{U}| \leq \frac{2(\Omega^*)^2 k}{\gamma^2 \pi \mathbf{c}}. \quad (6.55)$$

Proof of Claim 6.22.1. Let $\mathbf{U} \subseteq \mathbf{V}$ be the set of those clusters which intersect $N_{G_{\mathcal{D}}}(x_j)$. Using the same argument as in the proof of Claim 6.18.1 we get that $|\bigcup \mathbf{U}| \leq \frac{2(\Omega^*)^2 k}{\gamma^2}$, i.e. (6.54) holds. Also, (6.55) follows since $\mathcal{M}_A \cup \mathcal{M}_B$ is $(\varepsilon, d, \pi \mathbf{c})$ -semiregular. \square

Stage I: We proceed inductively for $j = 1, \dots, \ell$. Suppose that we embedded some components $\mathcal{F}_1, \dots, \mathcal{F}_{j-1}$ of the forests $T_1 - r_1, \dots, T_{j-1} - r_{j-1}$. We write F_{j-1} for the partial images of this embedding. We inductively assume that

$$F_{j-1} \text{ is } \tau k\text{-balanced w.r.t. } \mathcal{M}_A \cup \mathcal{M}_B. \quad (6.56)$$

For each $(A, B) \in \mathcal{M}_A \cup \mathcal{M}_B$ with $\deg_{G_{\mathcal{D}}}(x_j, (A \cup B) \setminus \mathfrak{A}) > 0$ take a subpair (A', B') ,

$$A' \subseteq (A \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\setminus 2} \setminus F_{j-1} \quad \text{and} \quad B' \subseteq (B \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\setminus 2} \setminus F_{j-1},$$

such that

$$|A'| = |B'| = \min \left\{ |(A \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\setminus 2} \setminus F_{j-1}|, |(B \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\setminus 2} \setminus F_{j-1}| \right\}.$$

These pairs comprise a semiregular matching \mathcal{N}_j . (Pairs $(A, B) \in \mathcal{M}_A \cup \mathcal{M}_B$ with $\deg_{G_{\mathcal{D}}}(x_j, (A \cup B) \setminus \mathfrak{A}) = 0$ are not considered for the construction of \mathcal{N}_j .)

Let $\mathcal{M}_j := \{(A', B') \in \mathcal{N}_j : |A'| > \alpha |A|\}$, for

$$\alpha := \frac{\eta^3 \gamma^2}{10^{10} (\Omega^*)^2}.$$

By Fact 2.1 \mathcal{M}_j is a $(2\varepsilon/\alpha, d/2, \alpha \pi \mathbf{c})$ -semiregular matching.

Claim 6.22.2. We have $|V(\mathcal{M}_j)| \geq |V(\mathcal{N}_j)| - \frac{\eta^3 k}{10^9}$.

Proof of Claim 6.22.2. By (6.54), and by Property 4 of Setting 5.1, we have $|V(\mathcal{M}_j)| \geq |V(\mathcal{N}_j)| - \alpha \cdot 2 \cdot \frac{2(\Omega^*)^2 k}{\gamma^2}$. \square

Let \mathcal{F}_j be a maximal set of components of $T_j - r_j$ such that

$$v(\mathcal{F}_j) \leq |V(\mathcal{M}_j)| - \frac{\eta^3 k}{10^9}. \quad (6.57)$$

Observe that if \mathcal{F}_j does not contain all the components of $T_j - r_j$ then

$$v(\mathcal{F}_j) \geq |V(\mathcal{M}_j)| - \frac{\eta^3 k}{10^9} - \tau k \geq |V(\mathcal{M}_j)| - \frac{2\eta^3 k}{10^9}. \quad (6.58)$$

Lemma 6.9 yields an embedding of \mathcal{F}_j in \mathcal{M}_j . Further the lemma together with the induction hypothesis (6.56) guarantees that the embedding can be chosen so that the new image set F_j is τk -balanced w.r.t. $\mathcal{M}_A \cup \mathcal{M}_B$. We fix this embedding, thus ensuring (6.56) for step i . If \mathcal{F}_j does not contain all the components of $T_j - r_j$ then (6.58) gives

$$|V(\mathcal{M}_j) \setminus F_j| \leq \frac{2\eta^3 k}{10^9}. \quad (6.59)$$

After Stage I: Let \mathcal{N}^* be a maximal semiregular matching contained in $(\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2}$ which avoids F_ℓ . We need two auxiliary claims.

Claim 6.22.3. We have

$$\deg_{G_{\mathcal{D}}}^{\max} \left(V_0 \cup V_1, V(\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2} \setminus (V(\mathcal{N}^*) \cup F_\ell \cup \mathfrak{A}) \right) < \frac{\eta^3 k}{10^9}.$$

Proof of Claim 6.22.3. Let us consider an arbitrary vertex $x \in V_0 \cup V_1$. By (6.55) the number of $(\mathcal{M}_A \cup \mathcal{M}_B)$ -vertices $C \subseteq \bigcup \mathbf{V}$ such that $\deg_{G_{\mathcal{D}}}(x, C) > 0$ is at most $\frac{2(\Omega^*)^2 k}{\gamma^2 \pi \epsilon}$.

Due to (6.56), we have for each $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edge (A, B) that

$$\left| (A \cup B)^{\uparrow 2} \setminus (V(\mathcal{N}^*) \cup F_\ell) \right| \leq \tau k. \quad (6.60)$$

Thus summing (6.60) over all $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edges (A, B) with $\deg_{G_{\mathcal{D}}}(x, (A \cup B) \setminus \mathfrak{A}) > 0$ we get

$$\deg_{G_{\mathcal{D}}}(x, V(\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2} \setminus (V(\mathcal{N}^*) \cup F_\ell \cup \mathfrak{A})) \leq \frac{4(\Omega^*)^2 k}{\gamma^2 \pi \epsilon} \cdot \tau k.$$

The claim now follows by (5.1). \square

Claim 6.22.4. Let $j \in [\ell]$ be such that \mathcal{F}_j does not consist of all the components of $T_j - r_j$. Then there exists an \mathcal{N}^* -cover \mathcal{X}_j such that $\deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathcal{X}_j) \leq \frac{3\eta^3 k}{10^9}$.

Proof of Claim 6.22.4. First, we define an $(\mathcal{M}_A \cup \mathcal{M}_B)$ -cover \mathcal{R}_j as follows. For an $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edge (A, B) let \mathcal{R}_j contain A if

$$|(A \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\uparrow 2} \setminus F_{j-1}| \leq |(B \cap N_{G_{\mathcal{D}} \cup G_{\nabla}}(x_j))^{\uparrow 2} \setminus F_{j-1}|,$$

and B otherwise. Observe that by the definition of \mathcal{N}_j , we have

$$\deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathcal{R}_j \setminus V(\mathcal{N}_j)) = 0. \quad (6.61)$$

6.5 Main embedding lemmas

Also, we have $V(\mathcal{N}^*) \cap \bigcup \mathcal{R}_j \cap V(\mathcal{M}_j) \subseteq V(\mathcal{N}^*) \cap V(\mathcal{M}_j) \subseteq V(\mathcal{M}_j) \setminus F_j$. In particular, (6.59) gives that

$$\left| V(\mathcal{N}^*) \cap \bigcup \mathcal{R}_j \cap V(\mathcal{M}_j) \right| \leq \frac{2\eta^3 k}{10^9}. \quad (6.62)$$

Let \mathcal{X}_j be the restriction of \mathcal{R}_j to \mathcal{N}^* . We then have

$$\begin{aligned} \deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathcal{X}_j) &= \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}^*) \cap \bigcup \mathcal{R}_j) \\ &\stackrel{\text{(by (6.61))}}{\leq} \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}^*) \cap \bigcup \mathcal{R}_j \cap V(\mathcal{M}_j)) + \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}_j) \setminus V(\mathcal{M}_j)) \\ &\stackrel{\text{(by (6.62), Claim 6.22.2)}}{\leq} \frac{3\eta^3 k}{10^9}. \end{aligned}$$

□

For every $j \in [\ell]$ we define $\mathcal{N}_j^* \subseteq \mathcal{N}^*$ as those \mathcal{N}^* -edges (A, B) for which we have

$$((A \cup B) \setminus \bigcup \mathcal{X}_j) \cap \mathfrak{A} = \emptyset.$$

Stage II: We shall inductively for $j = 1, \dots, \ell$ embed those components of $T_j - r_j$ that are not included in \mathcal{F}_j ; let us denote the set of these components by \mathcal{K}_j . There is nothing to do when $\mathcal{K}_j = \emptyset$, so let us assume otherwise.

We write $\mathbf{L} := \{C \in \mathbf{V} : C \subseteq \mathbb{L}_{\eta, k}(G)\}$. Let $K \in \mathcal{K}_j$ be a component that has not been embedded yet. We write U' for the total image of what has been embedded (in Stage I, and Stage II so far), combined with U . We claim that x_j has a substantial degree into one of four specific vertex sets.

Claim 6.22.5. At least one of the following four cases occurs.

$$\text{(U1)} \quad \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}_j^*) \setminus \bigcup \mathcal{X}_j) - |U' \cap V(\mathcal{N}_j^*)| \geq \frac{\eta^2 k}{10^4},$$

$$\text{(U2)} \quad \deg_{G_{\mathcal{D}}}(x_j, \mathfrak{A} \setminus U') \geq \frac{\eta^2 k}{10^4},$$

$$\text{(U3)} \quad \deg_{G_{\nabla}}(x_j, V(G_{\text{exp}}) \setminus U') \geq \frac{\eta^2 k}{10^4},$$

$$\text{(U4)} \quad \deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathbf{L} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup U')) \geq \frac{\eta^2 k}{10^4}.$$

6.5 Main embedding lemmas

Proof. Write $U'' := (U')^{\setminus 2} = U' \setminus U$. By (5.29), we have

$$\begin{aligned}
\frac{h}{2} &\leq \deg_{G_{\nabla}}(x_j, V_{\text{good}}^{\setminus 2}) \\
&\leq \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}_j^*)^{\setminus 2} \setminus \bigcup \mathcal{X}_j) + \deg_{G_{\mathcal{D}}}(x_j, \mathfrak{A}^{\setminus 2} \setminus (V(\mathcal{N}_j^*) \cup V(G_{\text{exp}}) \cup \bigcup \mathcal{X}_j)) \\
&\quad + \deg_{G_{\nabla}}(x_j, V(G_{\text{exp}})^{\setminus 2}) + \deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathbf{L}^{\setminus 2} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V(\mathcal{N}_j^*))) \\
&\quad + \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{M}_A \cup \mathcal{M}_B)^{\setminus 2} \setminus (V(\mathcal{N}_j^*) \cup \mathfrak{A})) + \deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathcal{X}_j) \\
\text{(by C6.22.3, C6.22.4)} &\leq \deg_{G_{\mathcal{D}}}(x_j, V(\mathcal{N}_j^*) \setminus \bigcup \mathcal{X}_j) - |U' \cap V(\mathcal{N}_j^*)| \\
&\quad + \deg_{G_{\mathcal{D}}}(x_j, \mathfrak{A}^{\setminus 2} \setminus (V(\mathcal{N}_j^*) \cup \bigcup \mathcal{X}_j \cup U'')) + \deg_{G_{\nabla}}(x_j, V(G_{\text{exp}})^{\setminus 2} \setminus U'') \\
&\quad + \deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathbf{L}^{\setminus 2} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V(\mathcal{N}_j^*) \cup U'')) \\
&\quad + \frac{4\eta^3 k}{10^9} + |U''|.
\end{aligned}$$

The claim follows since $|U''| \leq \frac{h}{2} - \frac{\eta^2 k}{1000}$ by (6.51). \square

We now briefly describe how to embed K in each of the cases **(U1)**–**(U4)**.

- In case **(U1)** recall that each $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edge contains at most one \mathcal{N}_j^* -edge. Thus by (6.54) we get that there is an $(\mathcal{M}_A \cup \mathcal{M}_B)$ -edge (A, B) with

$$\deg_{G_{\mathcal{D}}}(x_j, (V(\mathcal{N}_j^*) \cap (A \cup B)) \setminus \bigcup \mathcal{X}_j) - |V(\mathcal{N}_j^*) \cap U' \cap (A \cup B)| \geq \frac{\eta^2 k}{10^4} \cdot \frac{\gamma^2}{2(\Omega^*)^2 k} \cdot |A|. \quad (6.63)$$

Let us fix this edge (A, B) , and let (A', B') be the corresponding edge in \mathcal{N}_j^* . Suppose without loss of generality that $B \in \mathcal{X}_j$. We can now embed K in (A', B') using Lemma 6.6 with the following input: $C_{L6.6} := A', D_{L6.6} := B', X_{L6.6} := A' \setminus U', X_{L6.6}^* := N_{G_{\mathcal{D}}}(x_j, A' \setminus U'), Y_{L6.6} := B' \setminus U', \varepsilon_{L6.6} := \frac{8 \cdot 10^4 (\Omega^*)^2 \varepsilon}{\gamma^2 \eta^2}, \beta_{L6.6} := d/6$. With the help of (6.63), we calculate that $\min\{X_{L6.6}, Y_{L6.6}\} \geq |X_{L6.6}^*| \geq \frac{\gamma^2 \eta^2 |A|}{4 \cdot 10^4 (\Omega^*)^2} \geq 4 \frac{\varepsilon_{L6.6}}{\beta_{L6.6}} |A'|$.

- In Case **(U2)** we embed K using Lemma 6.4 with the following input: $\varepsilon_{L6.4} := \varepsilon', U_{L6.4} := U', U_{L6.4}^* := N_{G_{\mathcal{D}}}(x_j, \mathfrak{A} \setminus U'), \ell := 1$.
- In Case **(U3)** we embed K using Lemma 6.5 with the following input: $H_{L6.5} := G_{\text{exp}}, V_{1,L6.5} := V_{2,L6.5} := V(G_{\text{exp}}), U_{L6.5} := U', U_{L6.5}^* := N_{G_{\text{exp}}}(x_j, V(G_{\text{exp}}) \setminus U'), Q_{L6.5} := 1, \zeta_{L6.5} := \rho, \ell_{L6.5} := 1$.
- In Case **(U4)** we proceed as follows. As $\deg_{G_{\mathcal{D}}}(x_j, V_{\neq \Psi}) < \frac{\eta^2 k}{10^5}$ (cf. Definition 5.12), we have

$$\deg_{G_{\mathcal{D}}}(x_j, \bigcup \mathbf{L} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V_{\neq \Psi} \cup U')) \geq \frac{2\eta^2 k}{10^5}.$$

As for (6.63), we use (6.54) to find a cluster $A \in \mathbf{L}$ with

$$\deg_{G_{\mathcal{D}}}(x_j, A \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V_{\neq \Psi} \cup U')) \geq \frac{2\eta^2 k}{10^5} \cdot \frac{\gamma^2}{2(\Omega^*)^2 k} \cdot |A| = \frac{\eta^2 \gamma^2}{10^5 (\Omega^*)^2} \cdot |A|. \quad (6.64)$$

6.5 Main embedding lemmas

Recall that by the definition of $L_\#$ and $V_{\neq \Psi}$, we have that $\deg_{G_\nabla}^{\min}(A \setminus (L_\# \cup V_{\neq \Psi}), V(G) \setminus \Psi) \geq (1 + \frac{4\eta}{5})k$. Thus at least one of the following subcases must occur for the set $A^* := (N_{G_D}(x_j) \cap A) \setminus (L_\# \cup V(G_{\text{exp}}) \cup V_{\neq \Psi} \cup U')$:

(U4a) For at least $\frac{1}{2}|A^*|$ vertices $v \in A^*$ we have $\deg_{G_\nabla}(v, \mathfrak{A} \setminus U') \geq \frac{2\eta k}{5}$.

(U4b) For at least $\frac{1}{2}|A^*|$ vertices $v \in A^*$ we have $\deg_{G_{\text{reg}}}(v, \bigcup \mathbf{V} \setminus U') \geq \frac{2\eta k}{5}$.

In case **(U4a)** we embed K using Lemma 6.4. Details are very similar to **(U2)**. As for case **(U2b)**, let us take an arbitrary vertex $v \in A^*$ with $\deg_{G_{\text{reg}}}(v, \bigcup \mathbf{V} \setminus U') \geq \frac{2\eta k}{5}$. In particular, using (6.54), we find a cluster $B \in \mathbf{V}$ with

$$\deg_{G_{\text{reg}}}(v, B \setminus U') \geq \frac{\gamma^2 \eta}{10(\Omega^*)^2} |B|. \quad (6.65)$$

Map the root r_K of K to v and embed $K - r_K$ in (A, B) using Lemma 6.6¹³ with the following input: $C_{L6.6} := B, D_{L6.6} := A, X_{L6.6} := B \setminus U', Y_{L6.6} := A \setminus U', X_{L6.6}^* := N_{G_{\text{reg}}}(v, B \setminus U'), \beta_{L6.6} := \gamma^2 \eta / (10(\Omega^*)^2), \varepsilon_{L6.6} := \varepsilon'$. By (6.64) and (6.65) we see that $X_{L6.6}, Y_{L6.6}$ and $X_{L6.6}^*$ are large enough.

Stage III: In this stage we embed the trees $\{T'_j\}_{j=1}^{\ell'}$. The embedding techniques are as in Stage II. The cover \mathcal{F}' from Definition 5.12 plays the same role as the covers \mathcal{X}_j in Stage II. Observe that \mathcal{F}' is universal whereas the covers \mathcal{X}_j are specific for each vertex x_j . A second simplification is that in Stage III we use the semiregular matching $(\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2}$ for embedding (in a counterpart of **(U1)**) instead of \mathcal{N}_j^* .

Again we proceed inductively for $j = 1, \dots, \ell$ with embedding the components of $T'_j - r'_j$, which we denote by \mathcal{K}'_j . Let $K \in \mathcal{K}'_j$ be a component that has not been embedded yet. We write U' for the total image of what has been embedded (in Stage I, II, and Stage III so far), combined with U and let $U'' = U' \cap \mathfrak{B}_2$. We claim that y_j has a substantial degree into one of four specific vertex sets.

Claim 6.22.6. At least one of the following four cases occurs.

$$\begin{aligned} \text{(U1')} \quad & \deg_{G_D}(y_j, V((\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2}) \setminus (\mathfrak{A} \cup \bigcup \mathcal{F}')) \\ & - |U'' \cap (\bigcup \mathcal{F}' \cup (V((\mathcal{M}_A \cup \mathcal{M}_B)^{\uparrow 2}) \setminus \mathfrak{A})| \geq \frac{\eta^2 k}{10^4}, \end{aligned}$$

$$\text{(U2')} \quad \deg_{G_D}(y_j, \mathfrak{A} \setminus U') \geq \frac{\eta^2 k}{10^4},$$

$$\text{(U3')} \quad \deg_{G_\nabla}(y_j, V(G_{\text{exp}}) \setminus U') \geq \frac{\eta^2 k}{10^4},$$

$$\text{(U4')} \quad \deg_{G_D}(y_j, \bigcup \mathbf{L} \setminus (L_\# \cup V(G_{\text{exp}}) \cup U')) \geq \frac{\eta^2 k}{10^4}.$$

¹³Lemma 6.6 deals with embedding a single tree in a regular pair, whereas $K - r_K$ has several components. We therefore apply the lemma repeatedly for each component.

6.5 Main embedding lemmas

Proof. As $y_j \in V_1$, we have that

$$\begin{aligned}
h &\leq \deg_{G_{\nabla}}(y_j, V_{\text{good}}^{|2}) \\
&\leq \deg_{G_{\mathcal{D}}}(y_j, V((\mathcal{M}_A \cup \mathcal{M}_B)^{|2}) \setminus (\mathfrak{A} \cup V(G_{\text{exp}}) \cup \bigcup \mathcal{F}')) + \deg_{G_{\mathcal{D}}}(y_j, \mathfrak{A}^{|2} \setminus (V(G_{\text{exp}}) \cup \bigcup \mathcal{F}')) \\
&\quad + \deg_{G_{\mathcal{D}}}(y_j, \bigcup \mathcal{F}') + \deg_{G_{\mathcal{D}}}(y_j, \bigcup \mathbf{L}^{|2} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V(\mathcal{M}_A \cup \mathcal{M}_B))) \\
&\quad + \deg_{G_{\nabla}}(y_j, V(G_{\text{exp}})^{|2}) + \deg_{G_{\mathcal{D}}}(y_j, V(\mathcal{M}_A \cup \mathcal{M}_B)^{|2} \setminus V((\mathcal{M}_A \cup \mathcal{M}_B)^{|2})) \\
(\text{by L 5.5}) \quad &\leq \deg_{G_{\mathcal{D}}}(y_j, V((\mathcal{M}_A \cup \mathcal{M}_B)^{|2}) \setminus (\mathfrak{A} \cup V(G_{\text{exp}}) \cup \bigcup \mathcal{F}')) \\
&\quad - \left| U'' \cap \left(\bigcup \mathcal{F}' \cup (V((\mathcal{M}_A \cup \mathcal{M}_B)^{|2}) \setminus \mathfrak{A}) \setminus V(G_{\text{exp}}) \right) \right| \\
&\quad + \deg_{G_{\mathcal{D}}}(y_j, \mathfrak{A}^{|2} \setminus (U'' \cup V(G_{\text{exp}}) \cup \bigcup \mathcal{F}')) + \deg_{G_{\nabla}}(y_j, V(G_{\text{exp}})^{|2} \setminus U'') \\
&\quad + \deg_{G_{\mathcal{D}}}(y_j, \bigcup \mathbf{L}^{|2} \setminus (L_{\#} \cup V(G_{\text{exp}}) \cup V(\mathcal{M}_A \cup \mathcal{M}_B) \cup U'')) + \frac{2\eta^3 k}{10^3} + \frac{\eta^2 k}{10^5} + |U''|.
\end{aligned}$$

The claim follows since $|U''| \leq \sum_j T_j + \sum_j T'_j \leq h - \frac{\eta^2 k}{1000}$. □

Cases **(U1')**–**(U4')** are treated analogously as Cases **(U1)**–**(U4)**. □

Lemma 6.23. *Suppose we are in Setting 5.1 and 5.4, and that the sets V_0 and V_1 witness Preconfiguration $(\heartsuit 2)(h)$. Suppose that $U \subseteq \mathfrak{P}_0 \cup \mathfrak{P}_1$, such that $|U| \leq k$. Suppose that $\{x_j\}_{j=1}^{\ell} \subseteq V_0 \cup V_1$ are distinct vertices. Let $\{(T_j, r_j)\}_{j=1}^{\ell}$ be a family of rooted trees such that each component of $T_j - r_j$ has order at most τk .*

If $\sum_j v(T_j) \leq h - \eta^2 k / 1000$ and $|U| + \sum_j v(T_j) \leq k$ then there exist disjoint $(r_j \leftrightarrow x_j, V(T_j) \setminus \{r_j\} \leftrightarrow V(G) \setminus U)$ -embeddings of T_j in G .

Proof. The proof is contained in the proof of Lemma 6.22. It suffices to repeat the first two stages of the embedding process in the proof. In that setting, we use $h_{\text{L6.22}} = 2h$. Note that the condition $\{x_j\} \subseteq V_0$ in the setting of Lemma 6.22 gives us the same possibilities for embedding as the condition $\{x_j\} \subseteq V_0 \cup V_1$ in the current setting (cf. (5.29) and (5.32)). □

Lemma 6.24. *Suppose we are in Setting 5.1 and 5.4, and at least one of the following configurations occurs:*

- Configuration $(\diamond 6)$ $(\frac{\eta^3 \rho^4}{10^{14}(\Omega^*)^4}, 4\pi, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta^2 \nu}{2 \cdot 10^4}, \frac{3\eta^3}{2 \cdot 10^3}, h)$,
- Configuration $(\diamond 7)$ $(\frac{\eta^3 \gamma^3 \rho}{10^{12}(\Omega^*)^4}, \frac{\eta \gamma}{400}, 4\pi, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta^2 \nu}{2 \cdot 10^4}, \frac{3\eta^3}{2 \cdot 10^3}, h)$, or
- Configuration $(\diamond 8)$ $(\frac{\eta^4 \gamma^4 \rho}{10^{15}(\Omega^*)^5}, \frac{\eta \gamma}{400}, \frac{4\epsilon}{p_1}, 4\pi, \frac{d}{2}, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{p_1 \pi c}{2k}, \frac{\eta^2 \nu}{2 \cdot 10^4}, h_1, h)$.

Suppose that $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ is a (τk) -fine partition of a rooted tree (T, r) of order k . If the total order of the end shrubs is at most $h - 2\frac{\eta^2 k}{10^3}$ and the total order of the internal shrubs is at most $h_1 - 2\frac{\eta^2 k}{10^5}$, then $T \subseteq G$.

6.5 Main embedding lemmas

Proof. Let T' be the tree induced by all the cut-vertices $W_A \cup W_B$ and all the internal shrubs. Summing up the order of the internal shrub and the cut-vertices, we get that $v(T') < h_1 - \frac{2\eta^2 k}{10^5}$. Fix an embedding of T' as in Lemma 6.20 (in configurations $(\diamond 6)$ and $(\diamond 7)$), or as in Lemma 6.21 (in configuration $(\diamond 8)$). This embedding now extends to external shrubs by Lemma 6.22 (in Preconfiguration $(\heartsuit 1)$, which can only occur in Configuration $(\diamond 6)$ and $(\diamond 7)$), or by Lemma 6.23 (in Preconfiguration $(\heartsuit 2)$). It is important to remember here that by Definition 3.3(1), the total order of end shrubs in \mathcal{S}_B is at most half the size of the total order of end shrubs. \square

The next lemma completely resolves Theorem 1.2 in the presence of Configuration $(\diamond 9)$.

Lemma 6.25. *Suppose we are in Setting 5.1 and 5.4, and assume we have Configuration $(\diamond 9)$ $(\delta, \frac{2\eta^3}{10^3}, h_1, h_2, \varepsilon_1, d_1, \mu_1, \varepsilon_2, d_2, \mu_2)$ with $d_2 > 10\varepsilon_2 > 0$, $4 \cdot 10^3 \leq d_2 \mu_2 \tau k$, $\max\{d_1, \tau/\mu_1\} \leq \gamma^2 \eta^2 / (4 \cdot 10^7 (\Omega^*)^2)$, $d_1^2/6 > \varepsilon_1 \geq \tau/\mu_1$ and $\delta k > 10^3/\tau$.*

Suppose that $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ is a (τk) -fine partition of a rooted tree (T, r) of order k . If the total order of the internal shrubs of $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ is at most $h_1 - \frac{\eta^2 k}{10^5}$, and the total order of the end shrubs is at most $h_2 - 2\frac{\eta^2 k}{10^3}$ then $T \subseteq G$.

Proof. Let $V_0, V_1, V_2, \mathcal{N}, \{Q_0^{(j)}, Q_1^{(j)}\}_{j \in \mathcal{Y}}$ and \mathcal{F}' witness $(\diamond 9)$. The embedding process has two stages. In the first stage we embed the knags and the internal shrubs of T . In the second stage we embed the end shrubs. The knags will be embedded in $V_0 \cup V_1$, and the internal shrubs will be embedded in $V(\mathcal{N})$. Lemma 6.22 will be used to embed the end shrubs.

The knags of $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ are embedded in such a way that W_A is embedded in V_1 and W_B is embedded in V_0 . Since no other part of T is embedded in $V_0 \cup V_1$ in the first stage, each knag can be embedded greedily using the minimum degree condition arising from the super-regularity of the pairs $\{(Q_0^{(j)}, Q_1^{(j)})\}_{j \in \mathcal{Y}}$ using the bound on the total order of knags coming from Definition 3.3(c), and using Lemma 6.8 with the following input: $\varepsilon_{L6.8} := \varepsilon_2$, $d_{L6.8} := d_2$, $\ell_{L6.8} := \mu_2 k$, $U_A \cup U_B$ is the image of $W_A \cup W_B$ embedded so far and $\{A_{L6.8}, B_{L6.8}\} := \{Q_0^{(j)}, Q_1^{(j)}\}$, where $j \in \mathcal{Y}$ is arbitrary for the first knag, and for all other knags P has the property that

$$N_{G_{\mathcal{D}}}(\phi(\text{Par}(P))) \cap Q_1^{(j)} \setminus U_A \neq \emptyset.$$

The existence of such an index j follows from the fact that

$$\phi(\text{Par}(P)) \in V_2, \tag{6.66}$$

together with condition (5.52). We shall ensure (6.66) during our embedding of the internal shrubs, see below.

We now describe how to embed an internal shrub $T^* \in \mathcal{S}_A$ whose parent $u \in W_A$ is embedded in a vertex $x \in V_1$. Let $w \in V(T^*)$ be the unique neighbor of a vertex from $W_A \setminus \{u\}$ (cf. Definition 3.3(h)). Let U be the image of the part of T embedded so far. The next claim will be useful for finding a suitable \mathcal{N} -edge for accommodating T^* .

Claim 6.25.1. There exists an \mathcal{N} -edge (A, B) , or an \mathcal{N} -edge (B, A) such that

$$\min \{ |N_{G_{\mathcal{D}}}(x) \cap V_2 \cap (A \setminus U)|, |B \setminus U| \} \geq 100d_1|A| + \tau k.$$

6.5 Main embedding lemmas

Proof of Claim 6.25.1. For the purpose of this claim we reorient \mathcal{N} so that $V_2(\mathcal{N}) \subseteq \bigcup \mathcal{F}'$.

Suppose the claim fails to be true. Then for each $(A, B) \in \mathcal{N}$ we have $|\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap (A \setminus U)| < 100d_1|A| + \tau k$ or $|B \setminus U| < 100d_1|A| + \tau k$. In either case we get

$$|\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap A| - |U \cap (A \cup B)| < 100d_1|A| + \tau k. \quad (6.67)$$

We write $S := \bigcup \{V(D) : D \in \mathcal{D}, x \in V(D)\}$. Combining Fact 4.3 and Fact 4.4 we get that

$$|S| \leq \frac{2(\Omega^*)^2 k}{\gamma^2}. \quad (6.68)$$

Let us look at the number

$$\lambda := \sum_{(A,B) \in \mathcal{N}} (|\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap A| - |U \cap (A \cup B)|). \quad (6.69)$$

For a lower bound on λ , we write $\lambda = |\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2| - |U \cap V(\mathcal{N})|$. (Note that $V_2 \subseteq V(\mathcal{N})$ as we are in Configuration $(\diamond 9)$.) The first term is at least h_1 by (5.51), while the second term is at most $h_1 - \frac{\eta^2 k}{10^5}$ by the assumptions of the lemma. Thus $\lambda \geq \frac{\eta^2 k}{10^5}$.

For an upper bound on λ we only consider those \mathcal{N} -edges (A, B) for which $\mathbb{N}_{G_{\mathcal{D}}}(x) \cap A \neq \emptyset$. In that case $A \subseteq S$ (cf. 3 of Setting 5.1). Thus, since \mathcal{N} is $(\varepsilon_1, d_1, \mu_1 k)$ -semiregular we get that

$$|\{(A, B) \in \mathcal{N} : \mathbb{N}_{G_{\mathcal{D}}}(x) \cap A \neq \emptyset\}| \leq \frac{|S|}{\mu_1 k}. \quad (6.70)$$

Thus,

$$\begin{aligned} \lambda &\leq \sum_{(A,B) \in \mathcal{N}, \mathbb{N}_{G_{\mathcal{D}}}(x) \cap A \neq \emptyset} (|\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap A| - |U \cap (A \cup B)|) \\ &\stackrel{\text{(by (6.67), (6.70))}}{\leq} 100d_1|S| + \frac{|S|}{\mu_1 k} \tau k \\ &\stackrel{\text{(by (6.68))}}{<} \frac{\eta^2 k}{10^5}, \end{aligned}$$

a contradiction. This finishes the proof of the claim. \square

By symmetry we suppose that Claim 6.25.1 gives an \mathcal{N} -edge (A, B) such that $\min \{|\mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap (A \setminus U)|, |B \setminus U|\} \geq 100d_1|A| + \tau k$. We apply Lemma 6.6 with input $C_{L6.6} := A$, $D_{L6.6} := B$, $X_{L6.6} = X_{L6.6}^* := \mathbb{N}_{G_{\mathcal{D}}}(x) \cap V_2 \cap (A \setminus U)$, $Y_{L6.6} := B \setminus U$, $\varepsilon_{L6.6} := \varepsilon_1$, $\beta_{L6.6} := d_1/3$. Then there exists an embedding of T^* in $V(\mathcal{N}) \setminus U$ such that w is embedded in V_2 . This ensures (6.66).

We remark that there may be several internal shrubs extending from $u \in W_A$. However Claim 6.25.1 and the subsequent application of Lemma 6.6 allows a sequential embedding of these shrubs. This finishes the first stage of the embedding process.

For the second stage, i.e., the embedding of the end shrubs of $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$, we first recall that the total order of end shrubs in \mathcal{S}_A is at most $h_2 - 2\frac{\eta^2 k}{10^3}$, and the total order of end shrubs in \mathcal{S}_B is at most $\frac{1}{2}(h_2 - 2\frac{\eta^2 k}{10^3})$ by Definition 3.3(1). The embedding is a straightforward application of Lemma 6.22. \square

The next lemma resolves Theorem 1.2 in the presence of Configuration $(\diamond 10)$.

Lemma 6.26. *Suppose we are in Setting 5.1. For every $\eta', d', \Omega > 0$ there exists $\tilde{\varepsilon} > 0$ such that for every $\nu' > 0$ with the property that*

$$\frac{\eta'\nu'}{200\Omega} > \tau \quad (6.71)$$

there exists a number k_0 such that the following holds for each $k > k_0$.

If G is a graph with Configuration $(\diamond 10)(\tilde{\varepsilon}, d', \nu'k, \Omega k, \eta')$ then $\mathbf{trees}(k) \subseteq G$.

Proof. We give a sketch of a proof, following [PS12]. The main difference is indicated in Section 6.1.5.

Suppose we have Configuration $(\diamond 10)(\tilde{\varepsilon}, d', \nu'k, \Omega k, \eta')$, and are given a rooted tree (T, r) of order k with a (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ given by Lemma 3.5. By replacing \mathcal{L}^* by $\mathcal{L}^* \setminus \mathcal{V}(\mathcal{M})$,¹⁴ we can assume that \mathcal{L}^* and $\mathcal{V}(\mathcal{M})$ are disjoint.

For each shrub $F \in \mathcal{S}_A \cup \mathcal{S}_B$, let $x_F \in V(F)$ be its root, i.e., its minimal element in the topological order. If F is internal then we also define y_F as its (unique) maximal element that neighbours W_A . We can partition the semiregular matching \mathcal{M} and the set \mathcal{L}^* into two parts: $\mathcal{M}_A \cup \mathcal{L}_A^*$ and $\mathcal{M}_B \cup \mathcal{L}_B^*$ so that the partition satisfies

$$\deg_{\tilde{G}}(v, V(\mathcal{M}_A) \cup \bigcup \mathcal{L}_A^*) \geq v(\mathcal{S}_A) + \frac{\eta'k}{4} \quad \text{and} \quad (6.72)$$

$$\deg_{\tilde{G}}(w, V(\mathcal{M}_B) \cup \bigcup \mathcal{L}_B^*) \geq v(\mathcal{S}_B) + \frac{\eta'k}{4}, \quad (6.73)$$

for all but at most $2\tilde{\varepsilon}|A|$ vertices $v \in A$ and for all but at most $2\tilde{\varepsilon}|B|$ vertices $w \in B$. To see this, observe that the nature of the regularized graph allows us to treat¹⁵ conditions (6.72), (6.73), or that of Definition 5.21(b) in terms of average degrees of vertices in A and B , rather than in terms of individual degrees.¹⁶ If A and B were connected to each cluster $X \in \mathcal{L}^* \cup \mathcal{V}(\mathcal{M})$ by regular pairs of the same density, say d_X , it would suffice to split \mathcal{L}^* and \mathcal{M} in the ratio $v(\mathcal{S}_A) : v(\mathcal{S}_B)$. In the general setting, this can also be achieved, as was done in [PS12, Lemma 9]. Let $h_{A, \mathcal{L}^*}, h_{B, \mathcal{L}^*}, h_{A, \mathcal{M}}, h_{B, \mathcal{M}}$ be the average degrees of vertices of A and B into $\mathcal{L}_A^*, \mathcal{L}_B^*, \mathcal{M}_A, \mathcal{M}_B$.

We will now use the regularity to embed the shrubs and the seeds in \tilde{G} . We start with mapping r to A or B (depending whether $r \in W_A$ or $r \in W_B$), and proceed along a topological order on T . We denote the partial embedding of T at any particular stage as ϕ . The vertices of W_A are mapped to A , the vertices of W_B are mapped to B . As for embedding the shrubs, initially we start with embedding the shrubs of \mathcal{S}_A to \mathcal{M}_A (we say that A is in the \mathcal{M} -mode), and embedding the shrubs of \mathcal{S}_B to \mathcal{M}_B (B is in the \mathcal{M} -mode). By filling up the \mathcal{M} -edges with the shrubs as balanced as possible we can guarantee that we do not run out of space in \mathcal{M}_A before embedding \mathcal{S}_A -shrubs of total order at least $h_{A, \mathcal{M}} - \eta'k/100$. An analogous property holds for embedding \mathcal{S}_B -shrubs. We omit details and instead refer to a very similar procedure in Lemma 6.25.¹⁷

At some moment we may run out of space in \mathcal{M}_A , or in \mathcal{M}_B . Say that this happens first with the matching \mathcal{M}_A . Let $\mathcal{S}_A^* \subseteq \mathcal{S}_A$ be the set of shrubs not embedded so far. We now describe how to proceed when A is in the \mathcal{L}^* -mode. In this mode, we will not embed an upcoming shrub $F \in \mathcal{S}_A^*$, but only reserve a set U_F , $|U_F| \leq v(F)$ which serves a reminder that we want to accommodate F

¹⁴This does not change validity of the conditions in Definition 5.21.

¹⁵up to a small error

¹⁶This property is also key in the classical dense setting of the regularity lemma.

¹⁷In Lemma 6.25 it was shown how to utilize (5.51) for embedding shrubs of order up to $\approx h_1$ in regular pairs.

later on. Suppose that the parent $\text{Par}(F) \in W_A$ of F has been mapped to a typical¹⁸ vertex $z \in A$ already. We have

$$\deg_{\tilde{G}}(z, \bigcup \mathcal{L}_A^*) \geq v(\mathcal{S}_A^*) + \frac{\eta'k}{100} \geq \sum_{F'} |U_{F'}| + \frac{\eta'k}{100},$$

where the sum ranges over the already processed \mathcal{S}_A^* -shrubs F' . Consequently, there is a cluster $X \in \mathcal{V}$ such that

$$\deg_{\tilde{G}}\left(z, X \setminus \bigcup_{F'} U_{F'}\right) > \frac{\eta'|X|}{100\Omega}. \quad (6.74)$$

Let us view F as a bipartite graph, and let a_F be the size of its color class that contains x_F . Let U_F be an arbitrary set of $(N_{\tilde{G}}(z) \cap X) \setminus \bigcup_{F'} U_{F'}$ of size a_F , and also let us fix an image $\phi(x_F) \in U_F$ arbitrarily. If F is an internal shrub, we further define $\phi(y_F) \in U_F \setminus \{\phi(x_F)\}$ arbitrarily. At this stage we consider F as processed.

Later, of course, also B can switch in the \mathcal{L}^* -mode as well. At that moment, we define \mathcal{S}_B^* , and start to only make reservations U_K in clusters of \mathcal{L}_B^* instead of embedding shrubs $K \in \mathcal{S}_B^*$.

After all shrubs of $\mathcal{S}_A^* \cup \mathcal{S}_B^*$ have been processed we finalize the embedding. Consider a shrub $F \in \mathcal{S}_A^* \cup \mathcal{S}_B^*$. Suppose that $U_F \subseteq X$ for some $X \in \mathcal{V}$. We use Definition 5.21(c) to find a cluster Y such that

$$d(X, Y) \geq \frac{|Y \cap (\text{im}(\phi) \cup \bigcup_{F' \text{ yet unembedded}} U_{F'})|}{|Y|} + \frac{\eta'}{100\Omega}.$$

As $\phi(x_F)$ and $\phi(y_F)$ are typical¹⁹, we can additionally require that

$$\deg_{\tilde{G}}(\phi(x_F), Y), \deg_{\tilde{G}}(\phi(y_F), Y) \geq (d(X, Y) - \sqrt{\varepsilon})|Y|.$$

Therefore, the regularity method allows us to embed F in the pair (X, Y) avoiding the already defined image of ϕ , and the sets $U_{F'}$ corresponding to yet unembedded shrubs F' . The fact that the threshold in (6.74) was taken quite high (compared to the size of the shrubs, see (6.71)) allows us to avoid atypical vertices. We also need this embedding to be compatible with the existing placements $\phi(x_F)$ and $\phi(y_F)$. In particular, we need to find a path of length $\text{dist}_F(x_F, y_F)$ from $\phi(x_F)$ to $\phi(y_F)$. Here, it is crucial that $\text{dist}_F(x_F, y_F) \geq 4$ (cf. Definition 3.3(i)).²⁰ We remark, that in general we cannot guarantee that $X \cap \phi(F) = U_F$. So the set U_F should be regarded merely as a measure of future occupation of X , rather than an indication of exact future placement. \square

7 Proof of Theorem 1.2

The proof builds on the main results from [HKP⁺a, HKP⁺b, HKP⁺c]. We extend our subscript notation to allow referencing to parameters from [HKP⁺a, HKP⁺b, HKP⁺c]. For example, $\eta_{\text{LL3.13}}$ refers to the parameter η from Lemma 3.13 from the Ist paper, that is, from [HKP⁺a].

Let $\alpha > 0$ be given. We set

$$\eta := \min\left\{\frac{1}{30}, \frac{\alpha}{2}\right\}.$$

¹⁸in the sense of Definition 5.21(b)

¹⁹in the sense of Definition 5.21(c)

²⁰Indeed, it could be that $N(\phi(x_F)) \cap N(\phi(y_F)) = \emptyset$, which would make it impossible to find a path of length 2 from $\phi(x_F)$ to $\phi(y_F)$. If, on the other hand $\text{dist}_F(x_F, y_F) \geq 4$, then we can always find such a path using a look-ahead embedding in the regular pair (X, Y) .

We wish to fix further constants satisfying (5.1). A trouble is that we do not know the right choice of Ω^* and Ω^{**} yet. Therefore we take $g := \lfloor \frac{100}{\eta^2} \rfloor + 1$ and fix suitable constants

$$\eta \gg \frac{1}{\Omega_1} \gg \frac{1}{\Omega_2} \gg \dots \gg \frac{1}{\Omega_{g+1}} \gg \rho \gg \gamma \gg d \geq \frac{1}{\Lambda} \geq \varepsilon \geq \pi \geq \hat{\alpha} \geq \varepsilon' \geq \nu \gg \tau \gg \frac{1}{k_0} > 0.$$

where the relations between the parameters are more exactly as follows:

$$\begin{aligned} \frac{1}{\Omega_1} &\leq \frac{\eta^9}{10^{25}}, \\ \frac{1}{\Omega_{j+1}} &\leq \frac{\eta^{27}}{10^{67}\Omega_j^{36}} \quad \text{for each } j = 1, \dots, g, \\ \rho &\leq \frac{\eta^9}{10^{25}\Omega_{g+1}^5}, \\ \gamma &\leq \frac{\eta^{18}\rho^{24}}{10^{90}\Omega_{g+1}^{28}}, \\ d &\leq \min \left\{ \frac{\gamma^2\eta^2}{10^8\Omega_{g+1}^2}, \beta_{\text{II.L5.1}}(\eta_{\text{II.L5.1}} := \eta, \Omega_{\text{II.L5.1}} := \Omega_{g+1}, \gamma_{\text{II.L5.1}} := \gamma) \right\}, \\ \frac{1}{\Lambda} &\leq \min \left\{ d, \frac{\eta^{24}\gamma^{24}\rho}{10^{96}\Omega_{g+1}^{36}} \right\}, \\ \varepsilon &\leq \min \left\{ \frac{1}{\Lambda}, \frac{\gamma^2\eta^3 d \rho}{10^{13}\Omega_{g+1}^4}, \tilde{\varepsilon}_{\text{L6.26}}(\eta'_{\text{L6.26}} := \eta/40, d'_{\text{L6.26}} := \gamma^2 d/2, \Omega_{\text{L6.26}} := \frac{(\Omega_{g+1})^2}{\gamma^2}) \right\}, \\ \pi &\leq \min \left\{ \varepsilon, \pi_{\text{II.L5.1}}(\eta_{\text{II.L5.1}} := \eta, \Omega_{\text{II.L5.1}} := \Omega_{g+1}, \gamma_{\text{II.L5.1}} := \gamma, \varepsilon_{\text{II.L5.1}} := \varepsilon) \right\}, \\ \hat{\alpha} &\leq \min \left\{ \pi, \alpha_{\text{II.L4.4}} \left(\Omega_{\text{II.L4.4}} := \Omega_{g+1}, \rho_{\text{II.L4.4}} := \frac{\gamma^2}{4}, \varepsilon_{\text{II.L4.4}} := \pi, \tau_{\text{II.L4.4}} := 2\rho \right) \right\}, \\ \varepsilon' &\leq \min \left\{ \frac{\hat{\alpha}^2\gamma^6\rho^2}{10^3\Omega_{g+1}^4}, \varepsilon'_{\text{II.L5.1}}(\eta_{\text{II.L5.1}} := \eta, \Omega_{\text{II.L5.1}} := \Omega_{g+1}, \gamma_{\text{II.L5.1}} := \gamma, \varepsilon_{\text{II.L5.1}} := \varepsilon) \right\}, \\ \nu &\leq \min \left\{ \frac{\hat{\alpha}\rho}{\Omega_{g+1}}, \varepsilon', \nu_{\text{I.L3.13}}(\eta_{\text{I.L3.13}} := \alpha, \Lambda_{\text{I.L3.13}} := \Lambda, \gamma_{\text{I.L3.13}} := \gamma, \varepsilon_{\text{I.L3.13}} := \varepsilon', \rho_{\text{I.L3.13}} := \rho) \right\}, \\ \tau &\leq 2\varepsilon\pi\nu, \\ \frac{1}{k_0} &\leq \min \left\{ \frac{\gamma^3\rho\eta^8\tau\nu}{10^3\Omega_{g+1}^3}, \frac{1}{k_0^*} \right\}, \end{aligned}$$

with

$$\begin{aligned} k_0^* &:= \max \left\{ k_{0,\text{I.L3.13}}(\eta_{\text{I.L3.13}} := \alpha, \Lambda_{\text{I.L3.13}} := \Lambda, \gamma_{\text{I.L3.13}} := \gamma, \varepsilon_{\text{I.L3.13}} := \varepsilon', \rho_{\text{I.L3.13}} := \rho), \right. \\ &\quad k_{0,\text{II.L4.4}}(\Omega_{\text{II.L4.4}} := \Omega_{g+1}, \rho_{\text{II.L4.4}} := \frac{\gamma^2}{4}, \varepsilon_{\text{II.L4.4}} := \pi, \tau_{\text{II.L4.4}} := 2\rho, \alpha_{\text{II.L4.4}} := \hat{\alpha}, \nu_{\text{II.L4.4}} := \frac{2\rho}{\Omega_{g+1}}), \\ &\quad k_{0,\text{II.L5.1}}(\eta_{\text{II.L5.1}} := \eta, \Omega_{\text{II.L5.1}} := \Omega_{g+1}, \gamma_{\text{II.L5.1}} := \gamma, \varepsilon_{\text{II.L5.1}} := \varepsilon, \nu_{\text{II.L5.1}} := \nu), \\ &\quad k_{0,\text{L5.2}}(p_{\text{L5.2}} := 10, \alpha_{\text{L5.2}} := \eta/100), \\ &\quad \left. k_{0,\text{L6.26}}(\eta'_{\text{L6.26}} := \eta/40, d'_{\text{L6.26}} := \gamma^2 d/2, \tilde{\varepsilon}_{\text{L6.26}} := \varepsilon, \Omega_{\text{L6.26}} := \frac{(\Omega_{g+1})^2}{\gamma^2}, \nu'_{\text{L6.26}} := \pi\sqrt{\varepsilon'\nu}) \right\}. \end{aligned}$$

In particular, this gives a relation between α and k_0 .

Suppose now that $k > k_0$, and $G \in \mathbf{LKS}(n, k, \alpha)$ is a graph, and $T \in \mathbf{trees}(k)$ is a tree. It is our goal to show that $T \subseteq G$.

Let us now turn to the proof. First, we process the tree T by considering any (τk) -fine partition $(W_A, W_B, \mathcal{S}_A, \mathcal{S}_B)$ of T rooted at an arbitrary root r . Such a partition exists by Lemma 3.5. Let m_1 and m_2 be the total order of internal shrubs and the end shrubs, respectively. Set

$$\mathfrak{p}_0 := \frac{\eta}{100} \quad \text{and} \quad \mathfrak{p}_i := \frac{\eta}{100} + \frac{m_i}{(1 + \frac{\eta}{30})k}, \text{ for } i = 1, 2.$$

In particular we have $\mathfrak{p}_i \in [\frac{\eta}{100}, 1]$ for $i = 1, 2, 3$.

To find a suitable structure in the graph G we proceed as follows. We apply [HKP⁺a, Lemma 3.13] with input graph $G_{\text{ILL3.13}} := G$ and parameters $\eta_{\text{ILL3.13}} := \alpha$, $\Lambda_{\text{ILL3.13}} := \Lambda$, $\gamma_{\text{ILL3.13}} := \gamma$, $\varepsilon_{\text{ILL3.13}} := \varepsilon'$, $\rho_{\text{ILL3.13}} := \rho$, the sequence $(\Omega_j)_{j=1}^{g+1}$, $k_{\text{ILL3.13}} := k$ and $b_{\text{ILL3.13}} := \frac{\rho k}{100\Omega^*}$. The lemma gives a graph $G'_{\text{ILL3.13}} \in \mathbf{LKSsmall}(n, k, \eta)$, and an index $i \in [g]$. Slightly abusing notation, we call this graph still G . Set $\Omega^* := \Omega_i$ and $\Omega^{**} := \Omega_{i+1}$. Now, [HKP⁺a, Lemma 3.13(c)] yields a $(k, \Omega^{**}, \Omega^*, \Lambda, \gamma, \varepsilon', \nu, \rho)$ -sparse decomposition $\nabla = (\Psi, \mathbf{V}, \mathcal{D}, G_{\text{reg}}, G_{\text{exp}}, \mathfrak{A})$. Let \mathfrak{c} be the size of any cluster in \mathbf{V} .

We now apply [HKP⁺b, Lemma 5.1] with parameters $\eta_{\text{IIL5.1}} := \eta$, $\Omega_{\text{IIL5.1}} := \Omega_{g+1}$, $\gamma_{\text{IIL5.1}} := \gamma$, $\varepsilon_{\text{IIL5.1}} := \varepsilon$, $k_{\text{IIL5.1}} := k$, and $\Omega_{\text{IIL5.1}}^* := \Omega^*$. Given the graph G with its sparse decomposition ∇ the lemma gives three $(\varepsilon, d, \pi\mathfrak{c})$ -semiregular matchings \mathcal{M}_A , \mathcal{M}_B , and $\mathcal{M}_{\text{good}} \subseteq \mathcal{M}_A$ which fulfill the assertion either of case **(K1)**, or of **(K2)**. The matchings \mathcal{M}_A and \mathcal{M}_B also define the sets $\mathbb{X}\mathbb{A}$ and $\mathbb{X}\mathbb{B}$.

The additional features provided by [HKP⁺a, Lemma 3.13] and [HKP⁺b, Lemma 5.1] guarantee that we are in the situation described in Setting 5.1. We apply Lemma 5.2 as described in Definition 5.3; the numbers $\mathfrak{p}_0, \mathfrak{p}_1, \mathfrak{p}_2$ are as defined above. This puts us in the setting described in Setting 5.4. We now use [HKP⁺c, Lemma 4.17] to obtain one of the following configurations.

- $(\diamond 1)$,
- $(\diamond 2) \left(\frac{\eta^{27}\Omega^{**}}{4 \cdot 10^{66}(\Omega^*)^{11}}, \frac{\sqrt[4]{\Omega^{**}}}{2}, \frac{\eta^9 \rho^2}{128 \cdot 10^{22} \cdot (\Omega^*)^5} \right)$,
- $(\diamond 3) \left(\frac{\eta^{27}\Omega^{**}}{4 \cdot 10^{66}(\Omega^*)^{11}}, \frac{\sqrt[4]{\Omega^{**}}}{2}, \frac{\gamma}{2}, \frac{\eta^9 \gamma^2}{128 \cdot 10^{22} \cdot (\Omega^*)^5} \right)$,
- $(\diamond 4) \left(\frac{\eta^{27}\Omega^{**}}{4 \cdot 10^{66}(\Omega^*)^{11}}, \frac{\sqrt[4]{\Omega^{**}}}{2}, \frac{\gamma}{2}, \frac{\eta^9 \gamma^3}{384 \cdot 10^{22}(\Omega^*)^5} \right)$,
- $(\diamond 5) \left(\frac{\eta^{27}\Omega^{**}}{4 \cdot 10^{66}(\Omega^*)^{11}}, \frac{\sqrt[4]{\Omega^{**}}}{2}, \frac{\eta^9}{128 \cdot 10^{22} \cdot (\Omega^*)^3}, \frac{\eta}{2}, \frac{\eta^9}{128 \cdot 10^{22} \cdot (\Omega^*)^4} \right)$,
- $(\diamond 6) \left(\frac{\eta^3 \rho^4}{10^{14}(\Omega^*)^4}, 4\pi, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta^2 \nu}{2 \cdot 10^4}, \frac{3\eta^3}{2000}, \mathfrak{p}_2 \left(1 + \frac{\eta}{20}\right)k \right)$,
- $(\diamond 7) \left(\frac{\eta^3 \gamma^3 \rho}{10^{12}(\Omega^*)^4}, \frac{\eta\gamma}{400}, 4\pi, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta^2 \nu}{2 \cdot 10^4}, \frac{3\eta^3}{2 \cdot 10^3}, \mathfrak{p}_2 \left(1 + \frac{\eta}{20}\right)k \right)$,
- $(\diamond 8) \left(\frac{\eta^4 \gamma^4 \rho}{10^{15}(\Omega^*)^5}, \frac{\eta\gamma}{400}, \frac{400\varepsilon}{\eta}, 4\pi, \frac{d}{2}, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta\pi\mathfrak{c}}{200k}, \frac{\eta^2 \nu}{2 \cdot 10^4}, \mathfrak{p}_1 \left(1 + \frac{\eta}{20}\right)k, \mathfrak{p}_2 \left(1 + \frac{\eta}{20}\right)k \right)$,
- $(\diamond 9) \left(\frac{\rho\eta^8}{10^{27}(\Omega^*)^3}, \frac{2\eta^3}{10^3}, \mathfrak{p}_1 \left(1 + \frac{\eta}{40}\right)k, \mathfrak{p}_2 \left(1 + \frac{\eta}{20}\right)k, \frac{400\varepsilon}{\eta}, \frac{d}{2}, \frac{\eta\pi\mathfrak{c}}{200k}, 4\pi, \frac{\gamma^3 \rho}{32\Omega^*}, \frac{\eta^2 \nu}{2 \cdot 10^4} \right)$,

-
- $(\diamond\mathbf{10})(\varepsilon, \frac{\gamma^2 d}{2}, \pi\sqrt{\varepsilon'}\nu k, \frac{(\Omega^*)^2 k}{\gamma^2}, \frac{\eta}{40})$

Depending on the actual configuration Lemma 6.16, Lemma 6.19, Lemma 6.24, Lemma 6.25, or Lemma 6.26 guarantees that $T \subseteq G$. This finishes the proof of the theorem.

8 Theorem 1.2 algorithmically

We now discuss the algorithmic aspects of our proof of Theorem 1.2, covering also its parts developed in [HKP⁺a, HKP⁺b, HKP⁺c].

That is, we would like to find an algorithm which finds a copy of a given tree $T \in \mathbf{trees}(k)$ in any given graph $G \in \mathbf{LKS}(n, k, \alpha)$ in time $O(n^C)$. Here the degree C of the polynomial is allowed to depend on α , but not on k . It can be verified that each of the steps of our proof — except the extraction of dense spots (cf. [HKP⁺a, Section 3.8]) — can be turned into a polynomial time algorithm. The two randomized steps — random splitting in [HKP⁺c, Section 3.2], and the use of the stochastic process Duplicate in Section 6 — can be also efficiently derandomized using a standard technique for derandomizing the Chernoff bound. Let us sketch how to deal with extracting dense spots.

The idea is as follows. Initially, we pretend that G_{exp} consists of the entire bounded-degree part $G - \Psi$ (cleaned for minimum degree ρk as in [HKP⁺a, (3.7)]). With such a supposed sparse decomposition ∇_1 we go through [HKP⁺b, Lemma 5.1] and [HKP⁺c, Lemma 4.17] to obtain a configuration. We now start embedding T as in Section 6. Note that at this moment G_{reg} and \mathfrak{A} are absent, and so, the only embedding techniques are those involving Ψ and G_{exp} . Now, either we embed T , or we fail. The only possible reason for failure is that we were unable to perform the one-step look-ahead strategy described in [HKP⁺a, Section 3.6], because G_{exp} was not really nowhere-dense. But then we actually localized a dense spot D_1 . We get an updated supposed sparse decomposition ∇_2 in which D_1 is removed from G_{exp} and added to \mathcal{D} (and modify G_{reg} or \mathfrak{A} accordingly). We keep iterating. Since in each step we extract at least $O(k^2)$ edges we iterate the above at most $e(G)/\Theta(k^2) = O(\frac{n}{k})$ times. We are certain to succeed eventually, since after $\Theta(\frac{n}{k})$ iterations we get an honest sparse decomposition.

It seems that this iterative method is generally applicable for problems which employ a sparse decomposition.

9 Acknowledgments

The work on this project lasted from the beginning of 2008 until 2014 and we are very grateful to the following institutions and funding bodies for their support.

During the work on this paper JH was also affiliated with Zentrum Mathematik, TU Munich and Department of Computer Science, University of Warwick. JH was funded by a BAYHOST fellowship, a DAAD fellowship, Charles University grant GAUK 202-10/258009, EPSRC award EP/D063191/1, and by an EPSRC Postdoctoral Fellowship during the work on the project. The work leading to this invention has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 259385, and was supported by the European Regional Development Fund (ERDF), project “NTIS - New Technologies for Information Society”, European Centre of Excellence, CZ.1.05/1.1.00/02.0090. JK and ESz acknowledge the support of NSF grant DMS-0902241.

DP has been also affiliated with the Institute of Theoretical Computer Science, Charles University in Prague, Zentrum Mathematik, TU Munich, the Department of Computer Science and DIMAP, University of Warwick, and the school of mathematics, University of Birmingham. DP acknowledges the support of the Marie Curie fellowship FIST, DFG grant TA 309/2-1, a DAAD fellowship, Czech Ministry of Education project 1M0545, EPSRC award EP/D063191/1, and the support of the EPSRC Additional Sponsorship, with a grant reference of EP/J501414/1 which facilitated her to travel with her young child and so she could continue to collaborate closely with her coauthors on this project. This grant was also used to host MS in Birmingham.

MS was affiliated with the Institute of Mathematics and Statistics, University of São Paulo, and the Centre for Mathematical Modeling, University of Chile. She was supported by a FAPESP fellowship, and by FAPESP travel grant PQ-EX 2008/50338-0, also CMM-Basal, and FONDECYT grant 11090141. She also received funding by EPSRC Additional Sponsorship EP/J501414/1.

We enjoyed the hospitality of the School of Mathematics of University of Birmingham, Center for Mathematical Modeling, University of Chile, Alfréd Rényi Institute of Mathematics of the Hungarian Academy of Sciences and Charles University, Prague, during our long term visits.

We are very grateful to Miklós Ajtai, János Komlós, Miklós Simonovits, and Endre Szemerédi. Their yet unpublished work on the Erdős-Sós Conjecture was the starting point for our project, and our solution crucially relies on the methods developed for the Erdős-Sós Conjecture.

JH would like to thank Maxim Sviridenko for discussion on the algorithmic aspects of the problem.

A doctoral thesis entitled *Structural graph theory* submitted by Jan Hladký in September 2012 under the supervision of Daniel Král at Charles University in Prague is based on the series of the papers [HKP^a, HKP^b, HKP^c, HKP^d]. The texts of the two works overlap greatly. We are grateful to PhD committee members Peter Keevash and Michael Krivelevich. Their valuable comments are reflected in the series.

Symbol index

- $[n]$, 1
- \preceq , 3
- (\clubsuit) , 13
- $(\diamond 1)$, 13
- $(\heartsuit 1)$, 14
- $(\diamond 10)$, 16
- $(\diamond 2)$, 13
- $(\heartsuit 2)$, 15
- $(\diamond 3)$, 13
- $(\diamond 4)$, 13
- $(\diamond 5)$, 14
- $(\diamond 6)$, 15
- $(\diamond 7)$, 15
- $(\diamond 8)$, 15
- $(\diamond 9)$, 16
- $(X_1 \hookrightarrow V_1, \dots, X_\ell \hookrightarrow V_\ell)$ -embedding, 24
- (exp)**, 15
- (reg)**, 15
- \mathfrak{c} , 10
- \mathcal{M} -cover, 14
- $\text{Ch}(U)$, 3
- $\text{Ch}(v)$, 3
- $d(G)$, 2
- $d(U, W)$, 2
- deg , 2
- deg^{\max} , 2
- deg^{\min} , 2
- $\text{dist}(v_1, v_2)$, 3
- $\text{Duplicate}(\ell)$, 23
- $E(G)$, 1
- $e(G)$, 1
- $e(X)$, 2
- $e(X, Y)$, 2
- \mathbb{F} , 12
- G_{∇} , 10
- ghost, 30
- \mathbf{G}_{reg} , 8, 10
- G_{∇} , 8
- $L_{\#}$, 11
- $\mathbf{LKS}(n, k, \eta)$, 2
- $\mathbf{LKS}_{\text{small}}(n, k, \eta)$, 2
- \mathcal{N}^i , 12
- $N(v)$, 2
- $\mathcal{N}_{\mathfrak{A}}$, 10
- \mathfrak{P}_i , 12
- P , 11
- P_1 , 11
- P_2 , 11
- $P_{\mathfrak{A}}$, 11
- \mathfrak{p}_i , 12
- $\text{Par}(U)$, 3
- $\text{Par}(v)$, 3
- shadow, 9
- Seed, 4
- $T(r, \uparrow x)$, 4
- U^i , 12
- $V_{\not\sim \Psi}$, 11
- \bar{V} , 12
- \bar{V} , 12
- \bar{V}^* , 12
- \bar{V} , 12
- $\mathbf{V}_{\rightsquigarrow \mathfrak{A}}$, 10
- $V_{\rightsquigarrow \mathfrak{A}}$, 10
- $V(G)$, 1
- $v(G)$, 1
- V_+ , 11
- $V_{\text{even}}(T, r)$, 3
- V_{good} , 11
- $V_{\text{odd}}(T, r)$, 3
- $\mathbb{Y}\mathbb{A}$, 11
- $\mathbb{Y}\mathbb{B}$, 11

General index

- $(X_1 \hookrightarrow V_1, \dots, X_\ell \hookrightarrow V_\ell)$ -embedding, 24
- avoiding, 7
- avoiding threshold, 8
- balanced set, 26
- balanced way of embedding, 25
- bipartite density, 2
- bounded decomposition, 7
- captured edges, 8
- child, 3
- cluster, 8
- consistent matching, 16
- cover, 14
- dense cover, 7
- dense spot, 7
- density, 2
- discrepancy, 26
- end subtree, 4
- fine partition, 4
- fruit, 4
- ghost, 30
- induced tree, 4
- internal subtree, 4
- irregular, 2
- knag, 6
- matching involution, 30
- nowhere-dense, 7
- ordered skeleton, 7
- parent, 3
- peripheral subshrub, 6
- principal subshrub, 6
- proportional splitting, 12
- regular pair, 2
- regularized graph, 16
- rooted tree, 3
- seed, 4
- semiregular matching, 9
- shrub, 6
- sparse decomposition, 8
- subshrub, 6
- super-regular pair, 2
- unbalanced way of embedding, 25

References

- [AKS95] M. Ajtai, J. Komlós, and E. Szemerédi. On a conjecture of Loeb. In *Graph theory, combinatorics, and algorithms, Vol. 1, 2 (Kalamazoo, MI, 1992)*, Wiley-Intersci. Publ., pages 1135–1146. Wiley, New York, 1995.
- [EFLS95] P. Erdős, Z. Füredi, M. Loeb, and V. T. Sós. Discrepancy of trees. *Studia Sci. Math. Hungar.*, 30(1-2):47–57, 1995.
- [HKP⁺a] J. Hladký, J. Komlós, D. Piguet, M. Simonovits, M. Stein, and E. Szemerédi. The approximate Loeb–Komlós–Sós Conjecture I: The sparse decomposition. manuscript.
- [HKP⁺b] J. Hladký, J. Komlós, D. Piguet, M. Simonovits, M. Stein, and E. Szemerédi. The approximate Loeb–Komlós–Sós Conjecture II: The rough structure of LKS graphs. manuscript.
- [HKP⁺c] J. Hladký, J. Komlós, D. Piguet, M. Simonovits, M. Stein, and E. Szemerédi. The approximate Loeb–Komlós–Sós Conjecture III: The finer structure of LKS graphs. manuscript.
- [HKP⁺d] J. Hladký, J. Komlós, D. Piguet, M. Simonovits, M. Stein, and E. Szemerédi. The approximate Loeb–Komlós–Sós Conjecture IV: Embedding techniques and the proof of the main result. manuscript.
- [HP] J. Hladký and D. Piguet. Loeb–Komlós–Sós Conjecture: dense case. Manuscript (arXiv:0805:4834).
- [PS12] D. Piguet and M. J. Stein. An approximate version of the Loeb–Komlós–Sós conjecture. *J. Combin. Theory Ser. B*, 102(1):102–125, 2012.
- [Zha11] Y. Zhao. Proof of the $(n/2 - n/2 - n/2)$ conjecture for large n . *Electron. J. Combin.*, 18(1):Paper 27, 61, 2011.